

DIVISIONAL-CONTINUATION APPLICATION TRANSMITTAL FORM
UNDER RULE 1.53(b) (former Rule 1.60)

DOCKET NUMBER	ANTICIPATED CLASSIFICATION OF THIS APPLICATION:		PRIOR APPLICATION SERIAL NUMBER: 09/439,165	PRIOR APPLICATION FILING DATE: NOVEMBER 12, 1999
MNI- 062CP2DV1	CLASS:	SUBCLASS:	EXAMINER:	ART UNIT:

ASSISTANT COMMISSIONER FOR PATENTS
 BOX PATENT APPLICATION
 WASHINGTON, DC 20231


CERTIFICATION UNDER 37 CFR 1.10

Date of Deposit: June 2, 2000Mailing Label Number: EL 266 739 353 US

I hereby certify that this 37 CFR 1.53(b) request and the documents referred to as attached therein are being deposited with the United States Postal Service on the date indicated above in an envelope as "Express Mail Post Office to Addressee" service under 37 CFR 1.10 and addressed to the Assistant Commissioner for Patents, Box Patent Application, Washington, D.C. 20231.

Larry Taylor

Name of Person Mailing Paper


 Signature of Person Mailing Paper

Dear Sir:

This is a request for filing a ☐ continuation ☒ divisional application under 37 CFR 1.53(b), of pending prior application serial no. 09/439,165 filed on November 12, 1999, of Rory A.J. Curtis entitled NOVEL MEMBERS OF THE CAPSAICIN/VANILLOID RECEPTOR FAMILY OF PROTEINS AND USES THEREOF.

1. ☒ Enclosed is a copy of the latest inventor signed application, including the oath or declaration as originally filed. The copy of the enclosed papers is as follows:
- ☒ 72 page(s) of specification
 - ☒ 4 page(s) of claims
 - ☒ 1 page(s) of abstract
 - ☒ 35 sheet(s) of drawing (Figures 1-17)
 - ☒ 5 page(s) of an executed declaration and power of attorney.

I hereby verify that the attached papers are a true copy of the prior complete application serial no. 09/439,165 as originally filed on November 12, 1999.

2. ☐ A verified statement to establish small entity status under 37 CFR 1.9 and 1.27, a copy of which is enclosed, was filed in the prior application and such status is still proper and desired (37 CFR 1.28(a)).
3. ☒ The filing fee is calculated below:

	NUMBER OF CLAIMS FILED			NUMBER EXTRA
TOTAL	* 50	MINUS	** 20	= 30
INDEP.	* 6	MINUS	*** 3	= 3
<input checked="" type="checkbox"/> MULTIPLE DEPENDENT CLAIMS				

SMALL ENTITY

RATE	FEE
x 9 =	\$ 00
x 39 =	\$ 00
+130 =	\$ 00
BASIC FEE	\$ 00
TOTAL	\$0 00

OR

OTHER THAN A
SMALL ENTITY

RATE	FEE
x 18 =	\$540.00
x 78 =	\$234.00
+ 260 =	\$260.00
BASIC FEE	\$690.00
TOTAL	\$1724.00


OR

4. ☒ The Commissioner is hereby authorized to charge any additional fees which may be required, or credit any overpayment, to Deposit Account No. 12-0080. A duplicate copy of this sheet is enclosed.
5. ☒ A check in the amount of \$1724.00 is enclosed for payment of the filing fee.
6. ☐ Cancel in this application original claims _____ of the prior application before calculating the filing fee. (At least one original independent claim must be retained for filing purposes.)
7. ☒ A preliminary amendment is enclosed. (Claims added by this amendment have been properly numbered consecutively beginning with the number next following the highest numbered original claims in the prior application.)
8. ☒ Amend the specification by inserting before the first line the sentences: "This application is a divisional application of serial no. 09/439,165 filed on November 12, 1999, pending. The contents of all of the aforementioned application(s) are incorporated herein by reference."
9. ☐ Please abandon said prior application as of the filing date accorded this application. A duplicate copy of this transmittal is enclosed for filing in the prior application file. (May be used if signed by person authorized by §1.138 and before payment of base issue fee.)
10. ☐ Transfer the drawings from the pending prior application to this application.
11. ☐ Priority of application serial no. _____ filed on _____ in _____ is claimed under 35 U.S.C. §119.
- ☐ The certified copy has been filed in prior application serial no. _____ filed on _____.
- ☐ The certified copy will follow.
12. ☐ The prior application is assigned of record to _____
13. ☐ A _____ month extension of time has been submitted in the parent application Serial No. _____ in order to establish copendency with the present application.
14. ☒ Also enclosed is an Associate Power of Attorney (including Appendix A).
15. ☒ The power of attorney in the prior application is to Amy E. Mandragouras.
- a. ☒ The power appears in the original papers in the prior application.
- b. ☐ Since the power does not appear in the original papers, a copy of the power in the prior application is enclosed.
- c. ☐ A new power has been executed and is attached.
16. ☒ Address all future communications (May only be completed by applicant, or attorney or agent of record) to Amy E. Mandragouras at **Customer Number: 000959** whose address is:
- Lahive & Cockfield, LLP
28 State Street
Boston, Massachusetts 02109
17. ☒ Any requests for extensions of time necessary in a parent application for establishing copendency between this application and a parent application are hereby requested and the Commissioner is authorized to charge any fee associated with such an extension to Deposit Account No. 12-0080.

18. ☒ Pursuant to 37 CFR 1.821(e), the computer readable form of the sequence listing for this new application is to be identical with the computer readable form of application serial no. 09/439,165. Please use the computer readable form of application serial no. 09/439,165 in lieu of filing a duplicate computer readable form in this application. Pursuant to 37 CFR 1.821(f), the content of the paper copy of the sequence listing for this new application and the computer readable form of application serial no. 09/439,165 are the same.

June 2, 2000

Date



Maria C. Laccotripe, Ph.D.

Agent for Applicant

Limited Recognition Under 37 C.F.R. 10.9(b)

LAHIVE & COCKFIELD, LLP
28 State Street
Boston, Massachusetts 02109
Tel. (617) 227-7400

- ☐ inventor(s) ☐ filed under §1.34(a)
☐ assignee of complete interest
☒ attorney or agent of record

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re the application of: Rory A.J. Curtis

Serial No.: N/A

Filed: Herewith

For: *Novel Members Of The Capsaicin/Vanilloid
Receptor Family Of Proteins And Uses Thereof*

Attorney Docket No.: MNI-062CP2DV1

Group Art Unit:

Examiner:

Assistant Commissioner for Patents
Box Patent Application
Washington, D.C. 20231

CERTIFICATION UNDER 37 CFR 1.10

Date of Deposit: June 2, 2000

Mailing Label Number: EL 266 739 353 US

I hereby certify that this 37 CFR 1.53(b) request and the documents referred to as attached therein are being deposited with the United States Postal Service on the date indicated above in an envelope as "Express Mail Post Office to Addressee" service under 37 CFR 1.10 and addressed to the Assistant Commissioner for Patents, Box Patent Application, Washington, D.C. 20231.

Larry Taylor
Name of Person Mailing Paper


Signature of Person Mailing Paper

PRELIMINARY AMENDMENT

Dear Sir:

Prior to examination of the above-identified application, please amend the application as follows:

In the claims:

Please amend claims 1-4, 10, 11, 21, 24 and 25 as follows:

1. **(Amended)** An isolated nucleic acid molecule selected from the group consisting of:
 - (a) a nucleic acid molecule comprising the nucleotide sequence set forth in [SEQ ID NO:1, 3, 4, 6, 7, 9, 10, or 12] SEQ ID NO:4 or 6 or a complement thereof; and
 - (b) a nucleic acid molecule consisting of the nucleotide sequence set forth in [SEQ ID NO:1, 3, 4, 6, 7, 9, 10, or 12] SEQ ID NO:4 or 6 or a complement thereof.

2. **(Amended)** An isolated nucleic acid molecule which encodes a polypeptide selected from the group consisting of:

(a) a polypeptide comprising the amino acid sequence set forth in [SEQ ID NO:2, 5, 8, or 11] SEQ ID NO:5; and

(b) a polypeptide consisting of the amino acid sequence set forth in [SEQ ID NO:2, 5, 8, or 11] SEQ ID NO:5.

3. **(Amended)** An isolated nucleic acid molecule which encodes a naturally occurring allelic variant of a polypeptide comprising the amino acid sequence set forth in [SEQ ID NO:2, 5, 8, or 11] SEQ ID NO:5.

4. **(Amended)** An isolated nucleic acid molecule selected from the group consisting of:

a) a nucleic acid molecule comprising a nucleotide sequence which is at least 83% identical to the nucleotide sequence of [SEQ ID NO:1, 3, 4, 6, 7, 9, 10, or 12] SEQ ID NO:4 or 6, or a complement thereof;

b) a nucleic acid molecule comprising a fragment of at least 20 nucleotides of a nucleic acid comprising the nucleotide sequence of [SEQ ID NO:1, 3, 4, 6, 7, 9, 10, or 12] SEQ ID NO:4 or 6, or a complement thereof;

c) a nucleic acid molecule which encodes a polypeptide comprising an amino acid sequence at least about 87% identical to the amino acid sequence of [SEQ ID NO:2, 5, 8, or 11] SEQ ID NO:5; and

d) a nucleic acid molecule which encodes a fragment of a polypeptide comprising the amino acid sequence of [SEQ ID NO:2, 5, 8, or 11] SEQ ID NO:5, wherein the fragment comprises at least 15 contiguous amino acid residues of the amino acid sequence of [SEQ ID NO:2, 5, 8, or 11] SEQ ID NO:5.

10. **(Amended)** An isolated polypeptide selected from the group consisting of:

a) a fragment of a polypeptide comprising the amino acid sequence of [SEQ ID NO:2, 5, 8, or 11] SEQ ID NO:5, wherein the fragment comprises at least 15 contiguous amino acids of [SEQ ID NO:2, 5, 8, or 11] SEQ ID NO:5;

b) a naturally occurring allelic variant of a polypeptide comprising the amino acid sequence of [SEQ ID NO:2, 5, 8, or 11] SEQ ID NO:5, wherein the polypeptide is encoded by a nucleic acid molecule which hybridizes to a nucleic acid molecule consisting of [SEQ ID NO:1, 3, 4, 6, 7, 9, 10, or 12] SEQ ID NO:4 or 6 under stringent conditions;

c) a polypeptide which is encoded by a nucleic acid molecule comprising a nucleotide sequence which is at least 83% identical to a nucleic acid comprising the nucleotide sequence of [SEQ ID NO:1, 3, 4, 6, 7, 9, 10, or 12] SEQ ID NO:4 or 6; and

d) a polypeptide comprising an amino acid sequence which is at least 87% identical to the amino acid sequence of [SEQ ID NO:2, 5, 8, or 11] SEQ ID NO:5.

11. **(Amended)** The isolated polypeptide of claim 10 comprising the amino acid sequence of [SEQ ID NO:2, 5, 8, or 11] SEQ ID NO:5.

21. **(Amended)** The method of claim 20, wherein the binding of the test compound to the polypeptide is detected by a method selected from the group consisting of:

- a) detection of binding by direct detection of test compound/polypeptide binding;
- b) detection of binding using a competition binding assay; and
- c) detection of binding using an assay for [hVR-1,] hVR-2[, or rVR-2] activity.

24. **(Amended)** A method for treating a subject having a disorder characterized by aberrant [hVR-1 or] hVR-2 protein activity or nucleic acid expression comprising administering to the subject a [hVR-1 or] hVR-2 modulator such that treatment of the subject occurs.

25. **(Amended)** The method of claim 24, wherein the [hVR-1 or] hVR-2 modulator is a small molecule.

REMARKS

Claims 1-26 were pending in the present application. Claims 1-4, 10, 11, 21, 24, and 25 have been amended. Accordingly, claims 1-26 are currently pending. For the Examiner's convenience the currently pending claims are set forth in Appendix A.

No new matter has been added. Any amendments to and/or cancellation of the claims should in no way be construed as an acquiescence to any of the Examiner's rejections and was done solely to expedite the prosecution of the application. Applicant reserves the right to pursue the claims as originally filed in this or a separate application(s).

CONCLUSION

In view of the amendments and remarks set forth above, it is respectfully submitted that this application is in condition for allowance. If there are any remaining issues or the Examiner believes that a telephone conversation with Applicant's Agent would be helpful in expediting prosecution of this application, the Examiner is invited to call the undersigned at (617) 227-7400.

Respectfully submitted,
LAHIVE & COCKFIELD, LLP



Maria C. Laccotripe, Ph.D.

Agent for Applicant

Limited Recognition Under 37 C.F.R. §10.9(b)

28 State Street
Boston, Massachusetts 02109
telephone: (617) 227-7400
facsimile: (617) 742-4214
Date: June 2, 2000

CONFIDENTIAL

APPENDIX A

1. An isolated nucleic acid molecule selected from the group consisting of:
 - (a) a nucleic acid molecule comprising the nucleotide sequence set forth in SEQ ID NO:4 or 6 or a complement thereof; and
 - (b) a nucleic acid molecule consisting of the nucleotide sequence set forth in SEQ ID NO:4 or 6 or a complement thereof.
2. An isolated nucleic acid molecule which encodes a polypeptide selected from the group consisting of:
 - (a) a polypeptide comprising the amino acid sequence set forth in SEQ ID NO:5; and
 - (b) a polypeptide consisting of the amino acid sequence set forth in SEQ ID NO:5.
3. An isolated nucleic acid molecule which encodes a naturally occurring allelic variant of a polypeptide comprising the amino acid sequence set forth in SEQ ID NO:5.
4. An isolated nucleic acid molecule selected from the group consisting of:
 - a) a nucleic acid molecule comprising a nucleotide sequence which is at least 83% identical to the nucleotide sequence of SEQ ID NO:4 or 6, or a complement thereof;
 - b) a nucleic acid molecule comprising a fragment of at least 20 nucleotides of a nucleic acid comprising the nucleotide sequence of SEQ ID NO:4 or 6, or a complement thereof;
 - c) a nucleic acid molecule which encodes a polypeptide comprising an amino acid sequence at least about 87% identical to the amino acid sequence of SEQ ID NO:5; and
 - d) a nucleic acid molecule which encodes a fragment of a polypeptide comprising the amino acid sequence of SEQ ID NO:5, wherein the fragment comprises at least 15 contiguous amino acid residues of the amino acid sequence of SEQ ID NO:5.
5. An isolated nucleic acid molecule comprising the nucleic acid molecule of any one of claims 1, 2, 3, or 4, and a nucleotide sequence encoding a heterologous polypeptide.
6. A vector comprising the nucleic acid molecule of any one of claims 1, 2, 3, or 4.

7. The vector of claim 6, which is an expression vector.
8. A host cell transfected with the expression vector of claim 7.
9. A method of expressing a polypeptide comprising culturing the host cell of claim 8 in an appropriate culture medium to, thereby, express the polypeptide.
10. An isolated polypeptide selected from the group consisting of:
 - a) a fragment of a polypeptide comprising the amino acid sequence of SEQ ID NO:5, wherein the fragment comprises at least 15 contiguous amino acids of SEQ ID NO:5;
 - b) a naturally occurring allelic variant of a polypeptide comprising the amino acid sequence of SEQ ID NO:5, wherein the polypeptide is encoded by a nucleic acid molecule which hybridizes to a nucleic acid molecule consisting of SEQ ID NO:4 or 6 under stringent conditions;
 - c) a polypeptide which is encoded by a nucleic acid molecule comprising a nucleotide sequence which is at least 83% identical to a nucleic acid comprising the nucleotide sequence of SEQ ID NO:4 or 6; and
 - d) a polypeptide comprising an amino acid sequence which is at least 87% identical to the amino acid sequence of SEQ ID NO:5.
11. The isolated polypeptide of claim 10 comprising the amino acid sequence of SEQ ID NO:5.
12. The polypeptide of claim 10, further comprising heterologous amino acid sequences.
13. An antibody which selectively binds to a polypeptide of claim 10.
14. A method for detecting the presence of a polypeptide of claim 10 in a sample comprising:
 - a) contacting the sample with a compound which selectively binds to the polypeptide; and
 - b) determining whether the compound binds to the polypeptide in the sample to thereby detect the presence of a polypeptide of claim 10 in the sample.

15. The method of claim 14, wherein the compound which binds to the polypeptide is an antibody.

16. A kit comprising a compound which selectively binds to a polypeptide of claim 10 and instructions for use.

17. A method for detecting the presence of a nucleic acid molecule of any one of claims 1, 2, 3, or 4 in a sample comprising:

- a) contacting the sample with a nucleic acid probe or primer which selectively hybridizes to the nucleic acid molecule; and
- b) determining whether the nucleic acid probe or primer binds to a nucleic acid molecule in the sample to thereby detect the presence of a nucleic acid molecule of any one of claims 1, 2, 3, or 4 in the sample.

18. The method of claim 17, wherein the sample comprises mRNA molecules and is contacted with a nucleic acid probe.

19. A kit comprising a compound which selectively hybridizes to a nucleic acid molecule of any one of claims 1, 2, 3, or 4 and instructions for use.

20. A method for identifying a compound which binds to a polypeptide of claim 10 comprising:

- a) contacting the polypeptide, or a cell expressing the polypeptide with a test compound; and
- b) determining whether the polypeptide binds to the test compound.

21. The method of claim 20, wherein the binding of the test compound to the polypeptide is detected by a method selected from the group consisting of:

- a) detection of binding by direct detection of test compound/polypeptide binding;
- b) detection of binding using a competition binding assay; and
- c) detection of binding using an assay for hVR-2 activity.

22. A method for modulating the activity of a polypeptide of claim 10 comprising contacting the polypeptide or a cell expressing the polypeptide with a compound which binds to the polypeptide in a sufficient concentration to modulate the activity of the polypeptide.

23. A method for identifying a compound which modulates the activity of a polypeptide of claim 10 comprising:

- a) contacting a polypeptide of claim 10 with a test compound; and
- b) determining the effect of the test compound on the activity of the polypeptide to thereby identify a compound which modulates the activity of the polypeptide.

24. A method for treating a subject having a disorder characterized by aberrant hVR-2 protein activity or nucleic acid expression comprising administering to the subject a hVR-2 modulator such that treatment of the subject occurs.

25. The method of claim 24, wherein the hVR-2 modulator is a small molecule.

26. The method of claim 24, wherein the disorder is a pain disorder.

NOVEL MEMBERS OF THE CAPSAICIN/VANILLOID RECEPTOR FAMILY OF PROTEINS AND USES THEREOF

5 Related Applications

This application claims priority to U.S. provisional Application No. 60/108,322, filed on November 13, 1998, U.S. provisional Application No. 60/114,078 filed on December 28, 1998, U.S. Patent Application Serial No.: 09/258,633 filed on February 26, 1999, and U.S. Patent Application Serial No.: 09/421,134 filed on October 19, 1999, incorporated herein in their entirety
10 by this reference.

Background of the Invention

Pain is initiated when the peripheral terminals of a subgroup of sensory neurons are activated by noxious chemical, mechanical or thermal stimuli. These neurons, called nociceptors, transmit information regarding tissue damage to pain-processing centres in the spinal chord and brain (Fields, H.L. *Pain*, McGraw-Hill, New York, 1987). Nociceptors are characterized in part, by their sensitivity to capsaicin, a vanilloid-containing compound, and a natural product of capsicum peppers that is the active ingredient of many "hot" and spicy foods. In mammals, exposure of nociceptor terminals to capsaicin leads initially to excitation of the neuron and the consequent perception of pain and local release of inflammatory mediators. With prolonged exposure, nociceptor terminals become insensitive to capsaicin, as well as to other noxious stimuli (Szolcsanyi, J. in *Capsaicin in the Study of Pain* (ed. Wood, J.) 1-26 (Academic, London, 1993). This latter phenomenon of nociceptor desensitization underlies the seemingly paradoxical use of capsaicin as an analgesic agent in the treatment of painful disorders ranging from viral and diabetic neuropathies to rheumatoid arthritis (Campbell, E. in *Capsaicin and the Study of Pain* (ed. Wood, J.) 255-272 (Academic, London, 1993); Szallasi, A. *et al.* (1996) *Pain* 68, 195-208). Some of this decreased sensitivity to noxious stimuli may result from reversible changes in the nociceptor, but the long-term loss of responsiveness can be explained by death of the nociceptor or destruction of its peripheral terminals following exposure to capsaicin (Jancso, G. *et al.* (1977) *Nature* 270, 741-743).

The cellular specificity of capsaicin action and its ability to evoke the sensation of burning pain have led to speculation that the target of capsaicin action plays an important physiological role in the detection of painful stimuli. Indeed, capsaicin may elicit the perception of pain by mimicking the actions of a physiological stimulus or an endogenous ligand produced during tissue injury (James, I.F., Kinkina, N.N. & Wood, J.N. in *Capsaicin in the Study of Pain* (ed. Wood, J.N.) 83-104 (Academic, London, 1993).

Caterina M.J. *et al.* have recently determined the molecular basis underlying this phenomenon by characterizing a functional cDNA that encodes a vanilloid receptor (VR-1) in rat

sensory ganglia (Caterina M. J. *et al.*, (1997) *Nature* 389:816-824). VR-1 is a vanilloid-gated, nonselective cation channel that resembles members of the transient receptor potential (TRP) channel family, first identified as components of the *Drosophila* phototransduction pathway (Montell *et al.* (1989) *Neuron* 2:1313-1323).

5

Table of Contents

	A.	Summary of the Invention	-2-
	B.	Brief Description of the Drawings	-6-
10	C.	Detailed Description of the Invention	-8-
	I.	Isolated Nucleic Acid Molecules	-12-
	II.	Isolated hVR-1, hVR-2, and rVR-2 Proteins and Anti- hVR-1, Anti-hVR-2, and Anti-rVR-2 Antibodies	-22-
	III.	Recombinant Expression Vectors and Host Cells	-31-
15	IV.	Pharmaceutical Compositions	-39-
	V.	Uses and Methods of the Invention	-43-
	A.	Screening Assays	-44-
	B.	Detection Assays	-49-
	1.	Chromosome Mapping	-50-
20	2.	Tissue Typing	-51-
	3.	Use of Partial hVR-1, hVR-2, and rVR-2 Sequences in Forensic Biology	-52-
	C.	Predictive Medicine	-53-
	1.	Diagnostic Assays	-53-
	2.	Prognostic Assays	-55-
	3.	Monitoring of Effects During Clinical Trials	-60-
	D.	Methods of Treatment	-61-
	1.	Prophylactic Methods	-61-
	2.	Therapeutic Methods	-62-
30	3.	Pharmacogenomics	-63-
	D.	Examples	-65-

Summary of the Invention

The present invention is based, at least in part, on the discovery of novel members of the Capsaicin/Vanilloid family of receptors. Described herein is the isolation of the human orthologue of rat VR-1 (rVR-1), referred to herein as hVR-1, as well as another previously unknown member of the VR family of receptors, referred to herein as VR-2, and specifically as human VR-2 (hVR-2, including an alternate form which contains a deletion) and rat VR-2 (rVR-2) nucleic acid and

35

protein molecules. The hVR-1, hVR-2, and rVR-2 molecules of the present invention are useful as targets for developing modulating agents to regulate a variety of cellular processes, *e.g.*, cellular processes involved in pain. Accordingly, in one aspect, this invention provides isolated nucleic acid molecules encoding hVR-1, hVR-2, and rVR-2 proteins and fragments thereof, as well as
5 nucleic acid fragments suitable as primers or hybridization probes for the detection of hVR-1, hVR-2, and rVR-2-encoding nucleic acids.

In one embodiment, an hVR-1, hVR-2, or rVR-2 nucleic acid molecule of the invention is at least 60%, 65%, 70%, 75%, 80%, 83%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more identical to the nucleotide sequence (*e.g.*, to the entire length
10 of the nucleotide sequence) shown in SEQ ID NO:1, 3, 4, 6, 7, 9, 10, or 12 or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number _____, or a complement thereof.

In another embodiment, the isolated nucleic acid molecule includes the nucleotide sequence shown SEQ ID NO:1, 3, 4, 6, 7, 9, 10, or 12, or a complement thereof. In another embodiment, the nucleic acid molecule includes at least 10, 15, 20, or more contiguous nucleotides of SEQ ID NO:1, 3, 4, 6, 7, 9, 10, or 12.

In another embodiment, an hVR-1, hVR-2, and rVR-2 nucleic acid molecule includes a nucleotide sequence encoding a protein having an amino acid sequence sufficiently homologous to the amino acid sequence of SEQ ID NO:2, 5, 8, or 11. In one embodiment, an hVR-1, hVR-2, and rVR-2 nucleic acid molecule includes a nucleotide sequence encoding a protein having an amino acid sequence at least 60%, 65%, 70%, 75%, 80%, 85%, 87%, 90%, 95%, 98% or more identical to the entire length of the amino acid sequence of SEQ ID NO:2, 5, 8, or 11.

Another embodiment of the invention features nucleic acid molecules, preferably hVR-1, hVR-2, and rVR-2 nucleic acid molecules, which specifically detect hVR-1, hVR-2, and rVR-2 nucleic acid molecules relative to nucleic acid molecules encoding non-hVR-1, non-hVR-2, and non-hVR-2 proteins. For example, in one embodiment, such a nucleic acid molecule is at least 100-150, 1150-200, 200-250, 250-300, 300-350, 350-400, 400-450, 450-500, 500-550, 550-600, 600-700, 700-800, 800-900, 900-1000, 1088, or more nucleotides in length and hybridizes under stringent conditions to a nucleic acid molecule comprising the nucleotide sequence shown in SEQ
30 ID NO:1, 3, 4, 6, 7, 9, 10, or 12. In preferred embodiments, the nucleic acid molecules are at least 15 (*e.g.*, contiguous) nucleotides in length and hybridize under stringent conditions to nucleotides 1-17, 3696-3863, or 3901-3909 of SEQ ID NO:1. In other preferred embodiments, the nucleic acid molecules comprise nucleotides 1-17, 3696-3863, or 3901-3909 of SEQ ID NO:1. In yet other preferred embodiments, the nucleic acid molecules consist of nucleotides 1-17, 3696-3863, or
35 3901-3909 of SEQ ID NO:1. In preferred embodiments, the nucleic acid molecules are at least 15 (*e.g.*, contiguous) nucleotides in length and hybridize under stringent conditions to nucleotides 1944-2003 of SEQ ID NO:4. In other preferred embodiments, the nucleic acid molecules comprise

nucleotides 1944-2003 of SEQ ID NO:4. In yet other preferred embodiments, the nucleic acid molecules consist of nucleotides 1944-2003 of SEQ ID NO:4.

In other embodiments, the nucleic acid molecule encodes a naturally occurring allelic variant of a polypeptide comprising the amino acid sequence of SEQ ID NO:2, 5, 8, or 11, wherein the nucleic acid molecule hybridizes to a nucleic acid molecule consisting of SEQ ID NO:1, 3, 4, 6, 7, 9, 10, or 12 under stringent conditions and is encoded by the same locus as hVR-1, hVR-2 or rVR-2.

Another embodiment of the invention provides a nucleic acid molecule that encodes a naturally occurring orthologue of a polypeptide comprising the amino acid sequence of SEQ ID NO:2, 5, 8, or 11, wherein the nucleic acid molecule hybridizes to a nucleic acid molecule consisting of SEQ ID NO:1, 3, 4, 6, 7, 9, 10, or 12 under stringent conditions.

Another embodiment of the invention provides an isolated nucleic acid molecule which is antisense to an hVR-1, hVR-2, and rVR-2 nucleic acid molecule, *e.g.*, the coding strand of an hVR-1, hVR-2, and rVR-2 nucleic acid molecule.

Since the hVR2 (the alternate form) and rVR2 sequences represent fragments of the entire coding regions of these genes, another embodiment of the invention provides the complete gene sequences. A skilled artisan can readily isolate such molecule using the sequences disclosed herein.

Another aspect of the invention provides a vector comprising an hVR-1, an hVR-2, or a rVR-2 nucleic acid molecule. In certain embodiments, the vector is a recombinant expression vector. In another embodiment, the invention provides a host cell containing a vector of the invention. In yet another embodiment, the invention provides a host cell containing a nucleic acid molecule of the invention. The invention also provides a method for producing a protein, preferably an hVR-1, hVR-2, and rVR-2 protein, by culturing in a suitable medium, a host cell, *e.g.*, a mammalian host cell such as a non-human mammalian cell, of the invention containing a recombinant expression vector, such that the protein is produced.

Another aspect of this invention features isolated or recombinant hVR-1, hVR-2, and rVR-2 proteins and polypeptides. In one embodiment, the isolated protein, preferably an hVR-1, hVR-2, or rVR-2 protein, includes at least one transmembrane domain. In another embodiment, the isolated protein, preferably an hVR-1, hVR-2, or rVR-2 protein, includes at least one transmembrane domain and at least one proline rich domain. In yet another embodiment, the isolated protein, preferably an hVR-1, hVR-2, or rVR-2 protein, includes at least one transmembrane domain, at least one proline rich domain, and at least one ankyrin repeat domain. In yet another embodiment, the protein, preferably an hVR-1, hVR-2, or rVR-2 protein, includes at least one transmembrane domain, at least one proline rich domain, and at least one ankyrin repeat domain and has an amino acid sequence at least about 60%, 65%, 70%, 75%, 80%, 85%, 87%, 90%, 95%, 98% or more homologous to the amino acid sequence of SEQ ID NO:2, 5, 8, or 11. In another embodiment, the protein, preferably an hVR-1, hVR-2, or rVR-2 protein, includes at least

one transmembrane domain, at least one proline rich domain, and at least one ankyrin repeat domain and plays a role in the development and regulation of pain. In yet another embodiment, the protein, preferably an hVR-1, hVR-2, and rVR-2 protein, includes at least one transmembrane domain, at least one proline rich domain, and at least one ankyrin repeat domain and is encoded by a nucleic acid molecule having a nucleotide sequence which hybridizes under stringent hybridization conditions to a nucleic acid molecule comprising the nucleotide sequence of SEQ ID NO:1, 3, 4, 6, 7, 9, 10, or 12.

In another embodiment, the invention features fragments of the protein having the amino acid sequence of SEQ ID NO:2, 5, 8, or 11, wherein the fragment comprises at least 15, 30, 40, 50, 60, 70, 80, 90, or 100 amino acids (*e.g.*, contiguous amino acids).

In another embodiment, the invention features an isolated protein, preferably an hVR-1, hVR-2, and rVR-2 protein, which is encoded by a nucleic acid molecule consisting of a nucleotide sequence at least about 60%, 65%, 70%, 75%, 80%, 83%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more homologous to a nucleotide sequence of SEQ ID NO:1, 3, 4, 6, 7, 9, 10, or 12, or a complement thereof. This invention further features an isolated protein, preferably an hVR-1, hVR-2, or rVR-2 protein, which is encoded by a nucleic acid molecule consisting of a nucleotide sequence which hybridizes under stringent hybridization conditions to a nucleic acid molecule consisting of the nucleotide sequence of SEQ ID NO:1, 3, 4, 6, 7, 9, 10, or 12, or a complement thereof.

The proteins of the present invention or portions thereof, *e.g.*, biologically active portions thereof, can be operatively linked to a non-hVR-1, non-hVR-2, or non-rVR-2 polypeptide (*e.g.*, heterologous amino acid sequences) to form fusion proteins. The invention further features antibodies, such as monoclonal or polyclonal antibodies, that specifically bind proteins of the invention, preferably hVR-1, hVR-2, and rVR-2 proteins. In addition, the hVR-1, hVR-2, and rVR-2 proteins or biologically active portions thereof can be incorporated into pharmaceutical compositions, which optionally include pharmaceutically acceptable carriers.

In another aspect, the present invention provides a method for detecting the presence of an hVR-1, hVR-2, and rVR-2 nucleic acid molecule, protein or polypeptide in a biological sample by contacting the biological sample with an agent capable of detecting an hVR-1, hVR-2, and rVR-2 nucleic acid molecule, protein or polypeptide such that the presence of an hVR-1, hVR-2, and rVR-2 nucleic acid molecule, protein or polypeptide is detected in the biological sample.

In another aspect, the present invention provides a method for detecting the presence of hVR-1, hVR-2, and rVR-2 activity in a biological sample by contacting the biological sample with an agent capable of detecting an indicator of hVR-1, hVR-2, and rVR-2 activity such that the presence of hVR-1, hVR-2, and rVR-2 activity is detected in the biological sample.

In another aspect, the invention provides a method for modulating hVR-1, hVR-2, and rVR-2 activity comprising contacting a cell capable of expressing hVR-1, hVR-2, and rVR-2 with an agent that modulates hVR-1, hVR-2, and rVR-2 activity such that hVR-1, hVR-2, and rVR-2

activity in the cell is modulated. In one embodiment, the agent inhibits hVR-1, hVR-2, and rVR-2 activity. In another embodiment, the agent stimulates hVR-1, hVR-2, and rVR-2 activity. In one embodiment, the agent is an antibody that specifically binds to an hVR-1, hVR-2, and rVR-2 protein. In another embodiment, the agent modulates expression of hVR-1, hVR-2, and rVR-2 by modulating transcription of an hVR-1, hVR-2, and rVR-2 gene or translation of an hVR-1, hVR-2, and rVR-2 mRNA. In yet another embodiment, the agent is a nucleic acid molecule having a nucleotide sequence that is antisense to the coding strand of an hVR-1, hVR-2, and rVR-2 mRNA or an hVR-1, hVR-2, and rVR-2 gene.

In one embodiment, the methods of the present invention are used to treat a subject having a disorder characterized by aberrant hVR-1, hVR-2, and rVR-2 protein or nucleic acid expression or activity by administering an agent which is an hVR-1, hVR-2, and rVR-2 modulator to the subject. In one embodiment, the hVR-1, hVR-2, and rVR-2 modulator is an hVR-1, hVR-2, and rVR-2 protein. In another embodiment the hVR-1, hVR-2, and rVR-2 modulator is an hVR-1, hVR-2, and rVR-2 nucleic acid molecule. In yet another embodiment, the hVR-1, hVR-2, and rVR-2 modulator is a peptide, peptidomimetic, or other small molecule. In a further embodiment, the disorder characterized by aberrant hVR-1, hVR-2, and rVR-2 protein or nucleic acid expression is a pain disorder, *e.g.*, hyperalgesia.

The present invention also provides a diagnostic assay for identifying the presence or absence of a genetic alteration characterized by at least one of (i) aberrant modification or mutation of a gene encoding an hVR-1, hVR-2, and rVR-2 protein; (ii) mis-regulation of the gene; and (iii) aberrant post-translational modification of an hVR-1, hVR-2, and rVR-2 protein, wherein a wild-type form of the gene encodes a protein with an hVR-1, hVR-2, and rVR-2 activity (as described herein).

In another aspect the invention provides a method for identifying a compound that binds to or modulates the activity of an hVR-1, hVR-2, and rVR-2 protein, by providing an indicator composition comprising an hVR-1, hVR-2, and rVR-2 protein having hVR-1, hVR-2, and rVR-2 activity, contacting the indicator composition with a test compound, and determining the effect of the test compound on hVR-1, hVR-2, and rVR-2 activity in the indicator composition to identify a compound that modulates the activity of an hVR-1, hVR-2, and rVR-2 protein.

Other features and advantages of the invention will be apparent from the following detailed description and claims.

Brief Description of the Drawings

Figure 1 depicts the full length cDNA sequence and predicted amino acid sequence of human VR-1 (hVR-1). The nucleotide sequence corresponds to nucleic acids 1 to 3909 of SEQ ID NO:1. The amino acid sequence corresponds to amino acids 1 to 839 of SEQ ID NO:2. The

coding region without the 5' and 3' untranslated regions of the human VR-1 (hVR-1) gene is shown in SEQ ID NO:3.

Figure 2 depicts the full length cDNA sequence and predicted amino acid sequence of human VR-2 (hVR-2). The nucleotide sequence corresponds to nucleic acids 1 to 2809 of SEQ ID NO:4. The amino acid sequence corresponds to amino acids 1 to 764 of SEQ ID NO:5. The coding region without the 5' and 3' untranslated regions of the human VR-2 (hVR-2) gene is shown in SEQ ID NO:6.

Figure 3 depicts the partial cDNA sequence and partial predicted amino acid sequence of an alternate form of human VR-2 (hVR-2). The nucleotide sequence corresponds to nucleic acids 1 to 1489 of SEQ ID NO:7. The amino acid sequence corresponds to amino acids 1 to 436 of SEQ ID NO:8. The coding region without the 5' and 3' untranslated regions of the alternate form of human VR-2 (hVR-2) gene is shown in SEQ ID NO:9.

Figure 4 depicts the partial cDNA sequence and partial predicted amino acid sequence of rat VR-2 (rVR-2). The nucleotide sequence corresponds to nucleic acids 1 to 1794 of SEQ ID NO:10. The amino acid sequence corresponds to amino acids 1 to 554 of SEQ ID NO:11. The coding region without the 5' and 3' untranslated regions of the rat VR-2 (rVR-2) gene is shown in SEQ ID NO:12.

Figure 5 depicts an alignment of the hVR-1 protein (SEQ ID NO:2) with the human VR-2 protein (SEQ ID NO:5) using the GAP program in the GCG software package (Blosum 62 matrix) and a gap weight of 12 and a length weight of 4.

Figure 6 depicts an alignment of the hVR-1 nucleotide sequence (SEQ ID NO:1) with the human VR-2 nucleotide sequence (SEQ ID NO:4) using the GAP program in the GCG software package (nwsgapdna matrix) and a gap weight of 50 and a length weight of 3.

Figure 7 depicts an alignment of the hVR-2 protein (SEQ ID NO:5) with the rat VR-2 protein (SEQ ID NO:11) using the CLUSTAL W (1.74) multiple sequence alignment program.

Figure 8 depicts an alignment of the hVR-2 protein (SEQ ID NO:5) with the rat VR-2 protein (SEQ ID NO:11) using the GAP program in the GCG software package (Blosum 62 matrix) and a gap weight of 12 and a length weight of 4.

Figure 9 depicts an alignment of the hVR-1 nucleotide sequence (SEQ ID NO:1) with the rat VR-1 nucleotide sequence (Accession Number:AF029310) using the GAP program in the GCG software package (nwsgapdna matrix) and a gap weight of 50 and a length weight of 3.

Figure 10 depicts an alignment of the hVR-1 protein (SEQ ID NO:2) with the rat VR-1 protein (Accession Number:AF029310) using the GAP program in the GCG software package (Blosum 62 matrix) and a gap weight of 12 and a length weight of 4.

Figure 11 depicts an alignment of the hVR-2 protein (SEQ ID NO:5) with the human VR-2 protein (alternate form) (SEQ ID NO:8) using the CLUSTAL W (1.74) multiple sequence alignment program.

Figure 12 depicts a structural, hydrophobicity, and antigenicity analysis of the hVR-1 protein.

Figure 13 depicts the results of a search using the amino acid sequence of the hVR-1 protein against the HMM database.

Figure 14 depicts a structural, hydrophobicity, and antigenicity analysis of the hVR-2 protein.

Figure 15 depicts the results of a search using the amino acid sequence of the hVR-2 protein against the HMM database.

Figure 16 depicts the predicted full length amino acid sequence of the human VR-2 protein (alternate form) (SEQ ID NO:20).

Figure 17 depicts an alignment of the hVR-2 protein (SEQ ID NO:5) with the predicted full length human VR-2 protein (alternate form) (SEQ ID NO:20) using the CLUSTAL W (1.74) multiple sequence alignment program.

Detailed Description of the Invention

The present invention is based, at least in part, on the discovery of nucleic acid and amino acid molecules which are novel members of the Capsaicin/Vanilloid family of receptors. Described herein is the isolation of the human orthologue of rat VR-1 (rVR-1), referred to herein as hVR-1, as well as another previously unknown member of the VR family of receptors, referred herein as VR-2, and specifically as human VR-2 (hVR-2) and rat VR-2 (rVR-2) nucleic acid and protein molecules. The hVR-1, hVR-2, and rVR-2 molecules were identified based on their sequence similarity to the known rat vanilloid receptor (VR-1). VR-1 is a vanilloid gated, non-selective cation channel which resembles members of the transient receptor potential (TRP) ion channel family (described in Montell *et al.* (1989) *Neuron* 2:1313-1323) that mediate the influx of extracellular calcium in response to depletion of intracellular calcium stores. The rat VR-1 cDNA contains an open reading frame of 2514 nucleotides that encodes a protein of 838 amino acids. Hydrophilicity analysis has indicated that rat VR-1 contains six transmembrane domains (predicted to be mostly α -helices) with an additional short hydrophobic stretch between transmembrane regions 5 and 6. The amino terminal hydrophilic segment contains a relatively proline rich region followed by three ankyrin repeat domains. The rat VR-1 is expressed in small diameter neurons within sensory ganglia. The present hVR-1 sequence is the human orthologue of rVR-1. As described in further detail *infra*, the human VR-1 is expressed in nodose, trigeminal sensory neurons, as well as in some, but not all, small dorsal root ganglion (DRG) neurons and in a few medium sized DRG neurons.

The hVR-1, hVR-2, and rVR-2 molecules of the present invention play a role in pain signaling mechanisms. As used herein, the term "pain signaling mechanisms" includes the cellular mechanisms involved in the development and regulation of pain, *e.g.*, pain elicited by noxious chemical, mechanical, or thermal stimuli, in a subject, *e.g.*, a mammal such as a human. In

mammals, the initial detection of noxious chemical, mechanical, or thermal stimuli, a process referred to as "nociception", occurs predominantly at the peripheral terminals of specialized, small diameter primary afferent neurons, called polymodal nociceptors. These afferent neurons transmit the information to the central nervous system, evoking a perception of pain or discomfort and initiating appropriate protective reflexes. Capsaicin/Vanilloid receptors, *e.g.*, the hVR-1, hVR-2, and rVR-2 molecules of the present invention, present on these afferent neurons, are involved in detecting these noxious chemical, mechanical, or thermal stimuli and transducing this information into membrane depolarization events. Thus, the hVR-1, hVR-2, and rVR-2 molecules by participating in pain signaling mechanisms, can modulate pain elicitation and provide novel diagnostic targets and therapeutic agents to control pain.

The hVR-1, hVR-2, and rVR-2 molecules provide novel diagnostic targets and therapeutic agents to control pain in a variety of disorders, diseases, or conditions which are characterized by a deregulated, *e.g.*, upregulated or downregulated, pain response. For example, the hVR-1, hVR-2, and rVR-2 molecules provide novel diagnostic targets and therapeutic agents to control the exaggerated pain response elicited during various forms of tissue injury, *e.g.*, inflammation, infection, and ischemia, usually referred to as hyperalgesia (described in, for example, Fields, H.L. (1987) *Pain*, New York:McGraw-Hill). Moreover, the hVR-1, hVR-2, and rVR-2 molecules provide novel diagnostic targets and therapeutic agents to control pain associated with musculoskeletal disorders, *e.g.*, joint pain; tooth pain; headaches; pain associated with surgery, or neuropathic pain.

As the hVR-1 gene maps to a region of human chromosome 17 between WI-5436 (7.7cR) and WI-6584 (18.9cR) (Example 6), which has been associated with myasthenia gravis, Smith-Magenis syndrome, CORD5, Cone-rod dysrtrophy, and breast cancer, the hVR-1 molecule may provide novel diagnostic targets and therapeutic agents to treat, diagnose, or prognose these disorders or other disorders linked to this chromosomal region. Similarly, as the hVR-2 gene maps to a region of human chromosome 17 between AFMA043ZB5 (23.3 cR) and D17S721 (29.3cR) (Example 6) which has been associated with myasthenia gravis, Smith-Magenis syndrome, CORD5, Cone-rod dysrtrophy, choroidal dystrophy, central areolar, and retinal cone dystrophy, the hVR-2 molecule may provide novel diagnostic targets and therapeutic agents to treat, diagnose, or prognose these disorders or other disorders linked to this chromosomal region.

The term "family" when referring to the protein and nucleic acid molecules of the invention is intended to mean two or more proteins or nucleic acid molecules having a common structural domain or motif and having sufficient amino acid or nucleotide sequence homology as defined herein. Such family members can be naturally or non-naturally occurring and can be from either the same or different species. For example, a family can contain a first protein of human origin, as well as other, distinct proteins of human origin or alternatively, can contain homologues of non-human origin. Members of a family may also have common functional characteristics.

For example, the family of hVR-1, hVR-2, and rVR-2 proteins comprise at least one, and preferably six "transmembrane domains." As used herein, the term "transmembrane domain" includes an amino acid sequence of about 15 amino acid residues in length which spans the plasma membrane. More preferably, a transmembrane domain includes about at least 20, 25, 30, 35, 40, or 5 45 amino acid residues and spans the plasma membrane. Transmembrane domains are rich in hydrophobic residues, and typically have a helical structure. In a embodiment, at least 50%, 60%, 70%, 80%, 90%, 95% or more of the amino acid residues of a transmembrane domain are hydrophobic, *e.g.*, leucines, isoleucines, tyrosines, or tryptophans. Transmembrane domains are described in, for example, Zagotta W.N. et al, (1996) *Annual Rev. Neurosci.* 19: 235-63, the 10 contents of which are incorporated herein by reference. Amino acid residues 434-455, 480-495, (509-531; based on homology to the rat VR-1) or 514-531, (543-569; based on homology to the rat VR-1) or 538-555, (577-596; based on homology to the rat VR-1) or 580-599, and (656-683; based on homology to the rat VR-1) or 658-682 of hVR-1 (SEQ ID NO:2) and amino acid residues 391-410, 431-448, 459-476, 486-508, 538-556, and 621-645 of hVR-2 (SEQ ID NO:5) comprise 15 transmembrane domains.

In another embodiment, an hVR-1, hVR-2, and rVR-2 of the present invention is identified based on the presence of a "proline rich domain" in the protein or corresponding nucleic acid molecule. As used herein, the term "proline rich domain" includes an amino acid sequence of about 4-6 amino acid residues in length having the general sequence X-Pro-X-X-Pro-X (where X 20 can be any amino acid). Proline rich domains are usually located in a helical structure and bind through hydrophobic interactions to SH3 domains. SH3 domains recognize proline rich domains in both forward and reverse orientations. Proline rich domains are described in, for example, Sattler M. *et al.* (1998) *Leukemia* 12:637-644, the contents of which are incorporated herein by reference.

In another embodiment, an hVR-1, hVR-2, and rVR-2 of the present invention is identified 25 based on the presence of an "ankyrin repeat domain" in the protein or corresponding nucleic acid molecule. As used herein, the term "ankyrin repeat domain" includes a protein domain having an amino acid sequence of about 30-50 amino acid residues and having a bit score for the alignment of the sequence to the ankyrin repeat domain (HMM) of at least 6. Preferably, an ankyrin repeat domain includes at least about 30-45, more preferably about 30-40 amino acid residues, or about 30 30-35 amino acids and has a bit score for the alignment of the sequence to the ankyrin repeat domain (HMM) of at least 3-10, more preferably 10-30, more preferably 30-50, even more preferably 50-75, 75-100, 100-200 or greater. The ankyrin repeat domain HMM has been assigned the PFAM Accession PF00023 (<http://genome.wustl.edu/Pfam/.html>). Ankyrin repeats are involved in protein-protein interactions and are described in, for example, Ketchum K.A *et al.* 35 (1996) *FEBS Letters* 378:19-26, the contents of which are incorporated herein by reference.

To identify the presence of an ankyrin repeat domain in an hVR-1, hVR-2, and rVR-2 protein and make the determination that a protein of interest has a particular profile, the amino acid sequence of the protein is searched against a database of HMMs (*e.g.*, the Pfam database, release

2.1) using the default parameters (http://www.sanger.ac.uk/Software/Pfam/HMM_search). A description of the Pfam database can be found in Sonhammer *et al.* (1997) *Proteins* 28(3)405-420 and a detailed description of HMMs can be found, for example, in Gribskov *et al.* (1990) *Meth. Enzymol.* 183:146-159; Gribskov *et al.* (1987) *Proc. Natl. Acad. Sci. USA* 84:4355-4358; Krogh *et al.* (1994) *J. Mol. Biol.* 235:1501-1531; and Stultz *et al.* (1993) *Protein Sci.* 2:305-314, the contents of which are incorporated herein by reference. A search was performed against the HMM database resulting in the identification of three ankyrin repeat domains in the amino acid sequence of SEQ ID NO:2 (at about residues 201-233, 248-283, and 333-361) and SEQ ID NO:5 (at about residues 162-194, 208-243, and 293-328). The results of the searches are set forth in Figures 13 and 15.

Isolated proteins of the present invention, preferably hVR-1, hVR-2, and rVR-2 proteins, have an amino acid sequence sufficiently identical to the amino acid sequence of SEQ ID NO:2, 5, 8, or 11 or are encoded by a nucleotide sequence sufficiently identical to SEQ ID NO:1, 3, 4, 6, 7, 9, 10, or 12. As used herein, the term "sufficiently identical" refers to a first amino acid or nucleotide sequence which contains a sufficient or minimum number of identical or equivalent (e.g., an amino acid residue which has a similar side chain) amino acid residues or nucleotides to a second amino acid or nucleotide sequence such that the first and second amino acid or nucleotide sequences share common structural domains or motifs and/or a common functional activity. For example, amino acid or nucleotide sequences which share common structural domains have at least 30%, 40%, or 50% identity, preferably 60% identity, more preferably 70%-80%, and even more preferably 90-95% identity across the amino acid sequences of the domains and contain at least one and preferably two structural domains or motifs, are defined herein as sufficiently identical. Furthermore, amino acid or nucleotide sequences which share at least 30%, 40%, or 50%, preferably 60%, more preferably 70-80%, or 90-95% identity and share a common functional activity are defined herein as sufficiently identical.

As used interchangeably herein, an "hVR-1, hVR-2, and rVR-2 activity", "biological activity of hVR-1, hVR-2, and rVR-2" or "functional activity of hVR-1, hVR-2, and rVR-2", refers to an activity exerted by an hVR-1, hVR-2, and rVR-2 protein, polypeptide or nucleic acid molecule on an hVR-1, hVR-2, and rVR-2 responsive cell or on an hVR-1, hVR-2, and rVR-2 protein substrate, as determined *in vivo*, or *in vitro*, according to standard techniques. In one embodiment, an hVR-1, hVR-2, and rVR-2 activity is a direct activity, such as an association with an hVR-1, hVR-2, and rVR-2-target molecule. As used herein, a "target molecule" or "binding partner" is a molecule with which an hVR-1, hVR-2, and rVR-2 protein binds or interacts in nature, such that hVR-1, hVR-2, and rVR-2-mediated function is achieved. An hVR-1, hVR-2, and rVR-2 target molecule can be a non-hVR-1, non-hVR-2, and non-rVR-2 molecule or an hVR-1, hVR-2, and rVR-2 protein or polypeptide of the present invention. In an exemplary embodiment, an hVR-1, hVR-2, and rVR-2 target molecule is an hVR-1, hVR-2, and rVR-2 ligand, e.g., capsaicin. Alternatively, an hVR-1, hVR-2, and rVR-2 activity is an indirect activity, such as a cellular signaling activity mediated by interaction of the hVR-1, hVR-2, and rVR-2 protein with an hVR-1, hVR-2, and rVR-2 ligand.

Accordingly, another embodiment of the invention features isolated hVR-1, hVR-2, and rVR-2 proteins and polypeptides having an hVR-1, hVR-2, and rVR-2 activity. Other proteins of the invention are hVR-1, hVR-2, and rVR-2 proteins having at least one, and preferably six, transmembrane domains and, preferably, an hVR-1, hVR-2, and rVR-2 activity. Yet other proteins of the invention are hVR-1, hVR-2, and rVR-2 proteins having at least one transmembrane domain, at least one proline rich domain and, preferably, an hVR-1, hVR-2, and rVR-2 activity. Other proteins of the invention are hVR-1, hVR-2, and rVR-2 proteins having at least one transmembrane domain, at least one proline rich domain, at least one ankyrin repeat domain and, preferably, an hVR-1, hVR-2, and rVR-2 activity. Additional proteins of the invention have at least one transmembrane domain, at least one proline rich domain, at least one ankyrin repeat domain, and are, preferably, encoded by a nucleic acid molecule having a nucleotide sequence which hybridizes under stringent hybridization conditions to a nucleic acid molecule comprising the nucleotide sequence of SEQ ID NO:1, 3, 4, 6, 7, 9, 10, or 12.

The nucleotide sequence of the full length hVR-1 cDNA and the predicted amino acid sequence of the hVR-1 polypeptide are shown in Figure 1 and in SEQ ID NOs:1 and 2, respectively.

The nucleotide sequence of the full length hVR-2 cDNA and the predicted amino acid sequence of the hVR-2 polypeptide are shown in Figure 2 and in SEQ ID NOs:4 and 5, respectively.

The nucleotide sequence of the partial hVR-2 (alternate form) cDNA and the predicted amino acid sequence of the hVR-2 (alternate form) polypeptide are shown in Figure 3 and in SEQ ID NOs:7 and 8, respectively.

The nucleotide sequence of the partial rVR-2 cDNA and the predicted amino acid sequence of the rVR-2 polypeptide are shown in Figure 4 and in SEQ ID NOs:10 and 11, respectively.

Various aspects of the invention are described in further detail in the following subsections:

I. Isolated Nucleic Acid Molecules

One aspect of the invention pertains to isolated nucleic acid molecules that encode hVR-1, hVR-2, and rVR-2 proteins or biologically active portions thereof, as well as nucleic acid fragments sufficient for use as hybridization probes to identify hVR-1, hVR-2, and rVR-2-encoding nucleic acid molecules (*e.g.*, hVR-1, hVR-2, and rVR-2 mRNA) and fragments for use as PCR primers for the amplification or mutation of hVR-1, hVR-2, and rVR-2 nucleic acid molecules. As used herein, the term "nucleic acid molecule" is intended to include DNA molecules (*e.g.*, cDNA or genomic DNA) and RNA molecules (*e.g.*, mRNA) and analogs of the DNA or RNA generated using nucleotide analogs. The nucleic acid molecule can be single-stranded or double-stranded, but preferably is double-stranded DNA.

The term "isolated nucleic acid molecule" includes nucleic acid molecules which are separated from other nucleic acid molecules which are present in the natural source of the nucleic acid. For example, with regards to genomic DNA, the term "isolated" includes nucleic acid molecules which are separated from the chromosome with which the genomic DNA is naturally associated. Preferably, an "isolated" nucleic acid is free of sequences which naturally flank the nucleic acid (*i.e.*, sequences located at the 5' and 3' ends of the nucleic acid) in the genomic DNA of the organism from which the nucleic acid is derived. For example, in various embodiments, the isolated hVR-1, hVR-2, and rVR-2 nucleic acid molecule can contain less than about 5 kb, 4kb, 3kb, 2kb, 1 kb, 0.5 kb or 0.1 kb of nucleotide sequences which naturally flank the nucleic acid molecule in genomic DNA of the cell from which the nucleic acid is derived. Moreover, an "isolated" nucleic acid molecule, such as a cDNA molecule, can be substantially free of other cellular material, or culture medium when produced by recombinant techniques, or substantially free of chemical precursors or other chemicals when chemically synthesized.

A nucleic acid molecule of the present invention, *e.g.*, a nucleic acid molecule having the nucleotide sequence of SEQ ID NO:1, 3, 4, 6, 7, 9, 10, or 12. Using all or portion of the nucleic acid sequence of SEQ ID NO:1, 3, 4, 6, 7, 9, 10, or 12, as a hybridization probe, hVR-1, hVR-2, and rVR-2 nucleic acid molecules can be isolated using standard hybridization and cloning techniques (*e.g.*, as described in Sambrook, J., Fritsh, E. F., and Maniatis, T. *Molecular Cloning: A Laboratory Manual. 2nd, ed., Cold Spring Harbor Laboratory, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, 1989*).

Moreover, a nucleic acid molecule encompassing all or a portion of SEQ ID NO:1, 3, 4, 6, 7, 9, 10, or 12, can be isolated by the polymerase chain reaction (PCR) using synthetic oligonucleotide primers designed based upon the sequence of SEQ ID NO:1, 3, 4, 6, 7, 9, 10, or 12.

A nucleic acid of the invention can be amplified using cDNA, mRNA or alternatively, genomic DNA, as a template and appropriate oligonucleotide primers according to standard PCR amplification techniques. The nucleic acid so amplified can be cloned into an appropriate vector and characterized by DNA sequence analysis. Furthermore, oligonucleotides corresponding to hVR-1, hVR-2, and rVR-2 nucleotide sequences can be prepared by standard synthetic techniques, *e.g.*, using an automated DNA synthesizer.

In one embodiment, an isolated nucleic acid molecule of the invention comprises the nucleotide sequence shown in SEQ ID NO:1. The sequence of SEQ ID NO:1 corresponds to the full length hVR-1 encoding cDNA.

In another embodiment, an isolated nucleic acid molecule of the invention comprises the nucleotide sequence shown in SEQ ID NO:4. The sequence of SEQ ID NO:4 corresponds to the full length hVR-2 encoding cDNA.

In another embodiment, an isolated nucleic acid molecule of the invention comprises the nucleotide sequence shown in SEQ ID NO:7. The sequence of SEQ ID NO:7 corresponds to a fragment of the hVR-2 (alternate form) encoding cDNA.

In another embodiment, an isolated nucleic acid molecule of the invention comprises the nucleotide sequence shown in SEQ ID NO:10. The sequence of SEQ ID NO:10 corresponds to a fragment of the rVR-2 cDNA.

In another embodiment, an isolated nucleic acid molecule of the invention comprises a nucleic acid molecule which is a complement of the nucleotide sequence shown in SEQ ID NO:1, 3, 4, 6, 7, 9, 10, or 12, or a portion of any of these nucleotide sequences. A nucleic acid molecule which is complementary to the nucleotide sequence shown in SEQ ID NO:1, 3, 4, 6, 7, 9, 10, or 12, is one which is sufficiently complementary to the nucleotide sequence shown in SEQ ID NO:1, 3, 4, 6, 7, 9, 10, or 12, such that it can hybridize to the nucleotide sequence shown in SEQ ID NO:1, 3, 4, 6, 7, 9, 10, or 12 thereby forming a stable duplex.

In still another embodiment, an isolated nucleic acid molecule of the present invention comprises a nucleotide sequence which is at least about 60%, 65%, 70%, 75%, 80%, 83%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more homologous to the entire length of the nucleotide sequence shown in SEQ ID NO:1, 3, 4, 6, 7, 9, 10, or 12, or a portion of any of these nucleotide sequences.

Moreover, the nucleic acid molecule of the invention can comprise only a portion of the nucleic acid sequence of SEQ ID NO:1, 3, 4, 6, 7, 9, 10, or 12, for example, a fragment which can be used as a probe or primer or a fragment encoding a portion of an hVR-1, hVR-2, and rVR-2 protein, *e.g.*, a biologically active portion of an hVR-1, hVR-2, and rVR-2 protein. The nucleotide sequence determined from the cloning of the hVR-1, hVR-2, and rVR-2 gene allows for the generation of probes and primers designed for use in identifying and/or cloning other hVR-1, hVR-2, and rVR-2 family members, as well as hVR-1, hVR-2, and rVR-2 homologues from other species. The probe/primer typically comprises a substantially purified oligonucleotide. The oligonucleotide typically comprises a region of nucleotide sequence that hybridizes under stringent conditions to at least about 12 or 15, preferably about 20 or 25, more preferably about 30, 35, 40, 45, 50, 55, 60, 65, 75, or 100 consecutive nucleotides of a sense sequence of SEQ ID NO:1, 3, 4, 6, 7, 9, 10, or 12, of an anti-sense sequence of SEQ ID NO:1, 3, 4, 6, 7, 9, 10, or 12, or of a naturally occurring allelic variant or mutant of SEQ ID NO:1, 3, 4, 6, 7, 9, 10, or 12. In an exemplary embodiment, a nucleic acid molecule of the present invention comprises a nucleotide sequence which is greater than 100-150, 150-200, 200-250, 250-300, 300-350, 350-400, 400-450, 450-500, 500-550, 550-600, 600-650, 650-700, 700-750, 750-800, 800-850, 850-900, 900-950, 950-1000, 1088, or more nucleotides in length and hybridizes under stringent hybridization conditions to a nucleic acid molecule of SEQ ID NO:1, 3, 4, 6, 7, 9, 10, or 12.

Probes based on the hVR-1, hVR-2, and rVR-2 nucleotide sequences can be used to detect transcripts or genomic sequences encoding the same or homologous proteins. In preferred embodiments, the probe further comprises a label group attached thereto, *e.g.*, the label group can be a radioisotope, a fluorescent compound, an enzyme, or an enzyme co-factor. Such probes can be used as a part of a diagnostic test kit for identifying cells or tissue which misexpress an hVR-1,

hVR-2, and rVR-2 protein, such as by measuring a level of an hVR-1, hVR-2, and rVR-2-encoding nucleic acid in a sample of cells from a subject *e.g.*, detecting hVR-1, hVR-2, and rVR-2 mRNA levels or determining whether a genomic hVR-1, hVR-2, and rVR-2 gene has been mutated or deleted.

5 A nucleic acid fragment encoding a "biologically active portion of an hVR-1, hVR-2, and rVR-2 protein" can be prepared by isolating a portion of the nucleotide sequence of SEQ ID NO:1, 3, 4, 6, 7, 9, 10, or 12, which encodes a polypeptide having an hVR-1, hVR-2, and rVR-2 biological activity (the biological activities of the hVR-1, hVR-2, and rVR-2 proteins are described herein), expressing the encoded portion of the hVR-1, hVR-2, and rVR-2 protein (*e.g.*, by
10 recombinant expression *in vitro*) and assessing the activity of the encoded portion of the hVR-1, hVR-2, and rVR-2 protein.

The invention further encompasses nucleic acid molecules that differ from the nucleotide sequence shown in SEQ ID NO:1, 3, 4, 6, 7, 9, 10, or 12, due to degeneracy of the genetic code and thus encode the same hVR-1, hVR-2, and rVR-2 proteins as those encoded by the nucleotide sequence shown in SEQ ID NO:1, 3, 4, 6, 7, 9, 10, or 12. In another embodiment, an isolated nucleic acid molecule of the invention has a nucleotide sequence encoding a protein having an amino acid sequence shown in SEQ ID NO:2, 5, 8, or 11.

In addition to the hVR-1, hVR-2, and rVR-2 nucleotide sequences shown in SEQ ID NO:1, 3, 4, 6, 7, 9, 10, or 12, it will be appreciated by those skilled in the art that DNA sequence
15 polymorphisms that lead to changes in the amino acid sequences of the hVR-1, hVR-2, and rVR-2 proteins may exist within a population (*e.g.*, the human population). Such genetic polymorphism in the hVR-1, hVR-2, and rVR-2 genes may exist among individuals within a population due to natural allelic variation. As used herein, the terms "gene" and "recombinant gene" refer to nucleic acid molecules which include an open reading frame encoding an hVR-1, hVR-2, and rVR-2
20 protein, preferably a mammalian hVR-1, hVR-2, and rVR-2 protein, and can further include non-coding regulatory sequences, and introns.

Allelic variants of hVR-1, hVR-2, and rVR-2 include both functional and non-functional hVR-1, hVR-2, and rVR-2 proteins. Functional allelic variants are naturally occurring amino acid sequence variants of the hVR-1, hVR-2, and rVR-2 protein that maintain the ability to bind an
25 hVR-1, hVR-2, and rVR-2 ligand and/or modulate a pain signaling mechanism. Functional allelic variants will typically contain only conservative substitution of one or more amino acids of SEQ ID NO:2, 5, 8, or 11, or substitution, deletion or insertion of non-critical residues in non-critical regions of the protein.

Non-functional allelic variants are naturally occurring amino acid sequence variants of the
35 hVR-1, hVR-2, and rVR-2 protein that do not have the ability to either bind an hVR-1, hVR-2, and rVR-2 ligand and/or modulate a pain signaling mechanism. Non-functional allelic variants will typically contain a non-conservative substitution, a deletion, or insertion or premature truncation of

the amino acid sequence of SEQ ID NO:2, 5, 8, or 11, or a substitution, insertion or deletion in critical residues or critical regions.

The present invention further provides non-human orthologues of the hVR-2 and rVR-2 protein. Orthologues of the hVR-2 and rVR-2 protein are proteins that are isolated from non-human and non-rat organisms and possess the same hVR-2 and rVR-2 ligand binding and/or modulation of pain signaling mechanism capabilities of the hVR-2 and rVR-2 proteins. Orthologues of the hVR-2 and rVR-2 proteins can readily be identified as comprising an amino acid sequence that is substantially homologous to SEQ ID NO: 4, 6, 8 or 10.

Moreover, nucleic acid molecules encoding other hVR-1, hVR-2, and rVR-2 family members and, thus, which have a nucleotide sequence which differs from the hVR-1, hVR-2, and rVR-2 sequences of SEQ ID NO:1, 3, 4, 6, 7, 9, 10, or 12, are intended to be within the scope of the invention. For example, another hVR-1, hVR-2, and rVR-2 cDNA can be identified based on the nucleotide sequence of hVR-1, hVR-2, and rVR-2. Moreover, nucleic acid molecules encoding VR-2 proteins from different species, and which, thus, have a nucleotide sequence which differs from the hVR-2 and rVR-2 sequences of SEQ ID NO:4, 6, 8, or 10 are intended to be within the scope of the invention. For example, a mouse hVR-2 cDNA can be identified based on the nucleotide sequence of the human VR-2 (hVR-2) or the rat VR-2 (rVR-2).

Nucleic acid molecules corresponding to natural allelic variants and homologues of the hVR-1, hVR-2, and rVR-2 cDNAs of the invention can be isolated based on their homology to the hVR-1, hVR-2, and rVR-2 nucleic acids disclosed herein using the cDNAs disclosed herein, or a portion thereof, as a hybridization probe according to standard hybridization techniques under stringent hybridization conditions. Nucleic acid molecules corresponding to natural allelic variants and homologues of the hVR-1, hVR-2, and rVR-2 cDNAs of the invention can further be isolated by mapping to the same chromosome or locus as the hVR-1, hVR-2, and rVR-2 gene.

Accordingly, in another embodiment, an isolated nucleic acid molecule of the invention is at least 15, 20, 25, 30 or more nucleotides in length and hybridizes under stringent conditions to the nucleic acid molecule comprising the nucleotide sequence of SEQ ID NO:1, 3, 4, 6, 7, 9, 10, or 12. In other embodiment, the nucleic acid is at least 30, 50, 100, 150, 200, 250, 300, 350, 400, 450, 500, 550, 600, 650, 700, 750, 800, 850, 900, or 950 nucleotides in length. As used herein, the term "hybridizes under stringent conditions" is intended to describe conditions for hybridization and washing under which nucleotide sequences at least 60% identical to each other typically remain hybridized to each other. Preferably, the conditions are such that sequences at least about 70%, more preferably at least about 80%, even more preferably at least about 85% or 90% identical to each other typically remain hybridized to each other. Such stringent conditions are known to those skilled in the art and can be found in *Current Protocols in Molecular Biology*, John Wiley & Sons, N.Y. (1989), 6.3.1-6.3.6. A preferred, non-limiting example of stringent hybridization conditions are hybridization in 6X sodium chloride/sodium citrate (SSC) at about 45°C, followed by one or more washes in 0.2 X SSC, 0.1% SDS at 50°C, preferably at 55°C, more preferably at 60°C, and

even more preferably at 65°C. Preferably, an isolated nucleic acid molecule of the invention that hybridizes under stringent conditions to the sequence of SEQ ID NO:1, 3, 4, 6, 7, 9, 10, or 12 corresponds to a naturally-occurring nucleic acid molecule. As used herein, a "naturally-occurring" nucleic acid molecule refers to an RNA or DNA molecule having a nucleotide sequence that occurs in nature (*e.g.*, encodes a natural protein).

In addition to naturally-occurring allelic variants of the hVR-1, hVR-2, and rVR-2 sequences that may exist in the population, the skilled artisan will further appreciate that changes can be introduced by mutation into the nucleotide sequences of SEQ ID NO:1, 3, 4, 6, 7, 9, 10, or 12, thereby leading to changes in the amino acid sequence of the encoded hVR-1, hVR-2, and rVR-2 proteins, without altering the functional ability of the hVR-1, hVR-2, and rVR-2 proteins. For example, nucleotide substitutions leading to amino acid substitutions at "non-essential" amino acid residues can be made in the sequence of SEQ ID NO:1, 3, 4, 6, 7, 9, 10, or 12. A "non-essential" amino acid residue is a residue that can be altered from the wild-type sequence of hVR-1, hVR-2, and rVR-2 (*e.g.*, the sequence of SEQ ID NO:2, 5, 8, or 11) without altering the biological activity, whereas an "essential" amino acid residue is required for biological activity. For example, amino acid residues that are conserved among the hVR-1, hVR-2, and rVR-2 proteins of the present invention, are predicted to be particularly unamenable to alteration. Furthermore, additional amino acid residues that are conserved between the hVR-1, hVR-2, and rVR-2 proteins of the present invention and other members of the Capsaicin/Vanilloid receptor family are not likely to be amenable to alteration.

Accordingly, another aspect of the invention pertains to nucleic acid molecules encoding hVR-1, hVR-2, and rVR-2 proteins that contain changes in amino acid residues that are not essential for activity. Such hVR-1, hVR-2, and rVR-2 proteins differ in amino acid sequence from SEQ ID NO:2, 5, 8, or 11, yet retain biological activity. In one embodiment, the isolated nucleic acid molecule comprises a nucleotide sequence encoding a protein, wherein the protein comprises an amino acid sequence at least about 60%, 65%, 70%, 75%, 80%, 85%, 87%, 90%, 95%, 98% or more homologous to SEQ ID NO:2, 5, 8, or 11.

An isolated nucleic acid molecule encoding an hVR-1, hVR-2, and rVR-2 protein homologous to the protein of SEQ ID NO:2, 5, 8, or 11 can be created by introducing one or more nucleotide substitutions, additions or deletions into the nucleotide sequence of SEQ ID NO:1, 3, 4, 6, 7, 9, 10, or 12, such that one or more amino acid substitutions, additions or deletions are introduced into the encoded protein. Mutations can be introduced into SEQ ID NO:1, 3, 4, 6, 7, 9, 10, or 12, by standard techniques, such as site-directed mutagenesis and PCR-mediated mutagenesis. Preferably, conservative amino acid substitutions are made at one or more predicted non-essential amino acid residues. A "conservative amino acid substitution" is one in which the amino acid residue is replaced with an amino acid residue having a similar side chain. Families of amino acid residues having similar side chains have been defined in the art. These families include amino acids with basic side chains (*e.g.*, lysine, arginine, histidine), acidic side chains (*e.g.*, aspartic acid,

glutamic acid), uncharged polar side chains (*e.g.*, glycine, asparagine, glutamine, serine, threonine, tyrosine, cysteine), nonpolar side chains (*e.g.*, alanine, valine, leucine, isoleucine, proline, phenylalanine, methionine, tryptophan), beta-branched side chains (*e.g.*, threonine, valine, isoleucine) and aromatic side chains (*e.g.*, tyrosine, phenylalanine, tryptophan, histidine). Thus, a predicted nonessential amino acid residue in an hVR-1, hVR-2, and rVR-2 protein is preferably replaced with another amino acid residue from the same side chain family. Alternatively, in another embodiment, mutations can be introduced randomly along all or part of an hVR-1, hVR-2, and rVR-2 coding sequence, such as by saturation mutagenesis, and the resultant mutants can be screened for hVR-1, hVR-2, and rVR-2 biological activity to identify mutants that retain activity. Following mutagenesis of SEQ ID NO:1, 3, 4, 6, 7, 9, 10, or 12.

In a embodiment, a mutant hVR-1, hVR-2, and rVR-2 protein can be assayed for the ability to (1) interact with a non-hVR-1, non-hVR-2, or non-rVR-2 protein molecule, *e.g.*, a vanilloid compound such as capsaicin; (2) modulate intracellular calcium concentration; (3) activate an hVR-1, hVR-2, and rVR-2-dependent signal transduction pathway; or (4) modulate a pain signaling mechanism.

In addition to the nucleic acid molecules encoding hVR-1, hVR-2, and rVR-2 proteins described above, another aspect of the invention pertains to isolated nucleic acid molecules which are antisense thereto. An "antisense" nucleic acid comprises a nucleotide sequence which is complementary to a "sense" nucleic acid encoding a protein, *e.g.*, complementary to the coding strand of a double-stranded cDNA molecule or complementary to an mRNA sequence. Accordingly, an antisense nucleic acid can hydrogen bond to a sense nucleic acid. The antisense nucleic acid can be complementary to an entire hVR-1, hVR-2, and rVR-2 coding strand, or to only a portion thereof. In one embodiment, an antisense nucleic acid molecule is antisense to a "coding region" of the coding strand of a nucleotide sequence encoding hVR-1, hVR-2, and rVR-2. The term "coding region" refers to the region of the nucleotide sequence comprising codons which are translated into amino acid residues (*e.g.*, the coding region of hVR-1, hVR-2, and rVR-2). In another embodiment, the antisense nucleic acid molecule is antisense to a "noncoding region" of the coding strand of a nucleotide sequence encoding hVR-1, hVR-2, and rVR-2. The term "noncoding region" refers to 5' and 3' sequences which flank the coding region that are not translated into amino acids (*i.e.*, also referred to as 5' and 3' untranslated regions).

Given the coding strand sequences encoding hVR-1, hVR-2, and rVR-2 disclosed herein, antisense nucleic acids of the invention can be designed according to the rules of Watson and Crick base pairing. The antisense nucleic acid molecule can be complementary to the entire coding region of hVR-1, hVR-2, and rVR-2 mRNA, but more preferably is an oligonucleotide which is antisense to only a portion of the coding or noncoding region of hVR-1, hVR-2, and rVR-2 mRNA. For example, the antisense oligonucleotide can be complementary to the region surrounding the translation start site of hVR-1, hVR-2, and rVR-2 mRNA. An antisense oligonucleotide can be, for example, about 5, 10, 15, 20, 25, 30, 35, 40, 45 or 50 nucleotides in length. An antisense nucleic

acid of the invention can be constructed using chemical synthesis and enzymatic ligation reactions using procedures known in the art. For example, an antisense nucleic acid (*e.g.*, an antisense oligonucleotide) can be chemically synthesized using naturally occurring nucleotides or variously modified nucleotides designed to increase the biological stability of the molecules or to increase the physical stability of the duplex formed between the antisense and sense nucleic acids, *e.g.*, phosphorothioate derivatives and acridine substituted nucleotides can be used. Examples of modified nucleotides which can be used to generate the antisense nucleic acid include 5-fluorouracil, 5-bromouracil, 5-chlorouracil, 5-iodouracil, hypoxanthine, xantine, 4-acetylcytosine, 5-(carboxyhydroxymethyl) uracil, 5-carboxymethylaminomethyl-2-thiouridine, 5-carboxymethylaminomethyluracil, dihydrouracil, beta-D-galactosylqueosine, inosine, N6-isopentenyladenine, 1-methylguanine, 1-methylinosine, 2,2-dimethylguanine, 2-methyladenine, 2-methylguanine, 3-methylcytosine, 5-methylcytosine, N6-adenine, 7-methylguanine, 5-methylaminomethyluracil, 5-methoxyaminomethyl-2-thiouracil, beta-D-mannosylqueosine, 5'-methoxycarboxymethyluracil, 5-methoxyuracil, 2-methylthio-N6-isopentenyladenine, uracil-5-oxyacetic acid (v), wybutoxosine, pseudouracil, queosine, 2-thiocytosine, 5-methyl-2-thiouracil, 2-thiouracil, 4-thiouracil, 5-methyluracil, uracil-5-oxyacetic acid methylester, uracil-5-oxyacetic acid (v), 5-methyl-2-thiouracil, 3-(3-amino-3-N-2-carboxypropyl) uracil, (acp3)w, and 2,6-diaminopurine. Alternatively, the antisense nucleic acid can be produced biologically using an expression vector into which a nucleic acid has been subcloned in an antisense orientation (*i.e.*, RNA transcribed from the inserted nucleic acid will be of an antisense orientation to a target nucleic acid of interest, described further in the following subsection).

The antisense nucleic acid molecules of the invention are typically administered to a subject or generated *in situ* such that they hybridize with or bind to cellular mRNA and/or genomic DNA encoding an hVR-1, hVR-2, and rVR-2 protein to thereby inhibit expression of the protein, *e.g.*, by inhibiting transcription and/or translation. The hybridization can be by conventional nucleotide complementarity to form a stable duplex, or, for example, in the case of an antisense nucleic acid molecule which binds to DNA duplexes, through specific interactions in the major groove of the double helix. An example of a route of administration of antisense nucleic acid molecules of the invention include direct injection at a tissue site. Alternatively, antisense nucleic acid molecules can be modified to target selected cells and then administered systemically. For example, for systemic administration, antisense molecules can be modified such that they specifically bind to receptors or antigens expressed on a selected cell surface, *e.g.*, by linking the antisense nucleic acid molecules to peptides or antibodies which bind to cell surface receptors or antigens. The antisense nucleic acid molecules can also be delivered to cells using the vectors described herein. To achieve sufficient intracellular concentrations of the antisense molecules, vector constructs in which the antisense nucleic acid molecule is placed under the control of a strong pol II or pol III promoter are preferred.

In yet another embodiment, the antisense nucleic acid molecule of the invention is an -anomeric nucleic acid molecule. An α -anomeric nucleic acid molecule forms specific double-stranded hybrids with complementary RNA in which, contrary to the usual -units, the strands run parallel to each other (Gaultier *et al.* (1987) *Nucleic Acids. Res.* 15:6625-6641). The antisense nucleic acid molecule can also comprise a 2'-o-methylribonucleotide (Inoue *et al.* (1987) *Nucleic Acids Res.* 15:6131-6148) or a chimeric RNA-DNA analogue (Inoue *et al.* (1987) *FEBS Lett.* 215:327-330).

In still another embodiment, an antisense nucleic acid of the invention is a ribozyme. Ribozymes are catalytic RNA molecules with ribonuclease activity which are capable of cleaving a single-stranded nucleic acid, such as an mRNA, to which they have a complementary region. Thus, ribozymes (*e.g.*, hammerhead ribozymes (described in Haselhoff and Gerlach (1988) *Nature* 334:585-591)) can be used to catalytically cleave hVR-1, hVR-2, and rVR-2 mRNA transcripts to thereby inhibit translation of hVR-1, hVR-2, and rVR-2 mRNA. A ribozyme having specificity for an hVR-1, hVR-2, and rVR-2-encoding nucleic acid can be designed based upon the nucleotide sequence of an hVR-1, hVR-2, and rVR-2 cDNA disclosed herein (*i.e.*, SEQ ID NO:1, 3, 4, 6, 7, 9, 10, or 12). For example, a derivative of a *Tetrahymena* L-19 IVS RNA can be constructed in which the nucleotide sequence of the active site is complementary to the nucleotide sequence to be cleaved in an hVR-1, hVR-2, and rVR-2-encoding mRNA. See, *e.g.*, Cech *et al.* U.S. Patent No. 4,987,071; and Cech *et al.* U.S. Patent No. 5,116,742. Alternatively, hVR-1, hVR-2, and rVR-2 mRNA can be used to select a catalytic RNA having a specific ribonuclease activity from a pool of RNA molecules. See, for example, Bartel, D. and Szostak, J.W. (1993) *Science* 261:1411-1418.

Alternatively, hVR-1, hVR-2, and rVR-2 gene expression can be inhibited by targeting nucleotide sequences complementary to the regulatory region of the hVR-1, hVR-2, and rVR-2 (*e.g.*, the hVR-1, hVR-2, and rVR-2 promoter and/or enhancers) to form triple helical structures that prevent transcription of the hVR-1, hVR-2, and rVR-2 gene in target cells. See generally, Helene, C. (1991) *Anticancer Drug Des.* 6(6):569-84; Helene, C. *et al.* (1992) *Ann. N.Y. Acad. Sci.* 660:27-36; and Maher, L.J. (1992) *Bioassays* 14(12):807-15.

In yet another embodiment, the hVR-1, hVR-2, and rVR-2 nucleic acid molecules of the present invention can be modified at the base moiety, sugar moiety or phosphate backbone to improve, *e.g.*, the stability, hybridization, or solubility of the molecule. For example, the deoxyribose phosphate backbone of the nucleic acid molecules can be modified to generate peptide nucleic acids (see Hyrup B. *et al.* (1996) *Bioorganic & Medicinal Chemistry* 4 (1): 5-23). As used herein, the terms "peptide nucleic acids" or "PNAs" refer to nucleic acid mimics, *e.g.*, DNA mimics, in which the deoxyribose phosphate backbone is replaced by a pseudopeptide backbone and only the four natural nucleobases are retained. The neutral backbone of PNAs has been shown to allow for specific hybridization to DNA and RNA under conditions of low ionic strength. The synthesis of PNA oligomers can be performed using standard solid phase peptide synthesis

protocols as described in Hyrup B. *et al.* (1996) *supra*; Perry-O'Keefe *et al.* *Proc. Natl. Acad. Sci.* 93: 14670-675.

PNAs of hVR-1, hVR-2, and rVR-2 nucleic acid molecules can be used in therapeutic and diagnostic applications. For example, PNAs can be used as antisense or antigene agents for sequence-specific modulation of gene expression by, for example, inducing transcription or translation arrest or inhibiting replication. PNAs of hVR-1, hVR-2, and rVR-2 nucleic acid molecules can also be used in the analysis of single base pair mutations in a gene, (*e.g.*, by PNA-directed PCR clamping); as 'artificial restriction enzymes' when used in combination with other enzymes, (*e.g.*, S1 nucleases (Hyrup B. (1996) *supra*)); or as probes or primers for DNA sequencing or hybridization (Hyrup B. *et al.* (1996) *supra*; Perry-O'Keefe *supra*).

In another embodiment, PNAs of hVR-1, hVR-2, and rVR-2 can be modified, (*e.g.*, to enhance their stability or cellular uptake), by attaching lipophilic or other helper groups to PNA, by the formation of PNA-DNA chimeras, or by the use of liposomes or other techniques of drug delivery known in the art. For example, PNA-DNA chimeras of hVR-1, hVR-2, and rVR-2 nucleic acid molecules can be generated which may combine the advantageous properties of PNA and DNA. Such chimeras allow DNA recognition enzymes, (*e.g.*, RNase H and DNA polymerases), to interact with the DNA portion while the PNA portion would provide high binding affinity and specificity. PNA-DNA chimeras can be linked using linkers of appropriate lengths selected in terms of base stacking, number of bonds between the nucleobases, and orientation (Hyrup B. (1996) *supra*). The synthesis of PNA-DNA chimeras can be performed as described in Hyrup B. (1996) *supra* and Finn P.J. *et al.* (1996) *Nucleic Acids Res.* 24 (17): 3357-63. For example, a DNA chain can be synthesized on a solid support using standard phosphoramidite coupling chemistry and modified nucleoside analogs, *e.g.*, 5'-(4-methoxytrityl)amino-5'-deoxy-thymidine phosphoramidite, can be used as a between the PNA and the 5' end of DNA (Mag, M. *et al.* (1989) *Nucleic Acid Res.* 17: 5973-88). PNA monomers are then coupled in a stepwise manner to produce a chimeric molecule with a 5' PNA segment and a 3' DNA segment (Finn P.J. *et al.* (1996) *supra*). Alternatively, chimeric molecules can be synthesized with a 5' DNA segment and a 3' PNA segment (Peterser, K.H. *et al.* (1975) *Bioorganic Med. Chem. Lett.* 5: 1119-11124).

In other embodiments, the oligonucleotide may include other appended groups such as peptides (*e.g.*, for targeting host cell receptors *in vivo*), or agents facilitating transport across the cell membrane (see, *e.g.*, Letsinger *et al.* (1989) *Proc. Natl. Acad. Sci. USA* 86:6553-6556; Lemaitre *et al.* (1987) *Proc. Natl. Acad. Sci. USA* 84:648-652; PCT Publication No. W088/09810) or the blood-brain barrier (see, *e.g.*, PCT Publication No. W089/10134). In addition, oligonucleotides can be modified with hybridization-triggered cleavage agents (See, *e.g.*, Krol *et al.* (1988) *Bio-Techniques* 6:958-976) or intercalating agents. (See, *e.g.*, Zon (1988) *Pharm. Res.* 5:539-549). To this end, the oligonucleotide may be conjugated to another molecule, (*e.g.*, a peptide, hybridization triggered cross-linking agent, transport agent, or hybridization-triggered cleavage agent).

II. Isolated hVR-1, hVR-2, and rVR-2 Proteins and Anti-hVR-1, Anti-hVR-2, and Anti-rVR-2 Antibodies

One aspect of the invention pertains to isolated hVR-1, hVR-2, and rVR-2 proteins, and biologically active portions thereof, as well as polypeptide fragments suitable for use as immunogens to raise anti-hVR-1, anti-hVR-2, and anti-rVR-2 antibodies. In one embodiment, native hVR-1, hVR-2, and rVR-2 proteins can be isolated from cells or tissue sources by an appropriate purification scheme using standard protein purification techniques. In another embodiment, hVR-1, hVR-2, and rVR-2 proteins are produced by recombinant DNA techniques. Alternative to recombinant expression, an hVR-1, hVR-2, and rVR-2 protein or polypeptide can be synthesized chemically using standard peptide synthesis techniques.

An "isolated" or "purified" protein or biologically active portion thereof is substantially free of cellular material or other contaminating proteins from the cell or tissue source from which the hVR-1, hVR-2, and rVR-2 protein is derived, or substantially free from chemical precursors or other chemicals when chemically synthesized. The language "substantially free of cellular material" includes preparations of hVR-1, hVR-2, and rVR-2 protein in which the protein is separated from cellular components of the cells from which it is isolated or recombinantly produced. In one embodiment, the language "substantially free of cellular material" includes preparations of hVR-1, hVR-2, and rVR-2 protein having less than about 30% (by dry weight) of non-hVR-1, hVR-2, and rVR-2 protein (also referred to herein as a "contaminating protein"), more preferably less than about 20% of non-hVR-1, hVR-2, and rVR-2 protein, still more preferably less than about 10% of non-hVR-1, hVR-2, and rVR-2 protein, and most preferably less than about 5% non-hVR-1, non-hVR-2, and non-rVR-2 protein. When the hVR-1, hVR-2, and rVR-2 protein or biologically active portion thereof is recombinantly produced, it is also preferably substantially free of culture medium, *i.e.*, culture medium represents less than about 20%, more preferably less than about 10%, and most preferably less than about 5% of the volume of the protein preparation.

The language "substantially free of chemical precursors or other chemicals" includes preparations of hVR-1, hVR-2, and rVR-2 protein in which the protein is separated from chemical precursors or other chemicals which are involved in the synthesis of the protein. In one embodiment, the language "substantially free of chemical precursors or other chemicals" includes preparations of hVR-1, hVR-2, and rVR-2 protein having less than about 30% (by dry weight) of chemical precursors or non-hVR-1, hVR-2, and rVR-2 chemicals, more preferably less than about 20% chemical precursors or non-hVR-1, hVR-2, and rVR-2 chemicals, still more preferably less than about 10% chemical precursors or non-hVR-1, hVR-2, and rVR-2 chemicals, and most preferably less than about 5% chemical precursors or non-hVR-1, hVR-2, and rVR-2 chemicals.

As used herein, a "biologically active portion" of an hVR-1, hVR-2, and rVR-2 protein includes a fragment of an hVR-1, hVR-2, and rVR-2 protein which participates in an interaction between an hVR-1, hVR-2, and rVR-2 molecule and a non-hVR-1, non-hVR-2, and non-rVR-2

molecule, respectively. Biologically active portions of an hVR-1, hVR-2, and rVR-2 protein include peptides comprising amino acid sequences sufficiently homologous to or derived from the amino acid sequence of the hVR-1, hVR-2, and rVR-2 protein, *e.g.*, the amino acid sequence shown in SEQ ID NO:2, 5, 8, or 11, which include less amino acids than the full length hVR-1, hVR-2, and rVR-2 proteins, and exhibit at least one activity of an hVR-1, hVR-2, and rVR-2 protein. Typically, biologically active portions comprise a domain or motif with at least one activity of the hVR-1, hVR-2, and rVR-2 protein, *e.g.*, binding of an hVR-1, hVR-2, and rVR-2 ligand such as a vanilloid compound, *e.g.*, Capsaicin. A biologically active portion of an hVR-1, hVR-2, and rVR-2 protein can be a polypeptide which is, for example, 10, 20, 30, 40, 50, 60, 70, 80, 90, 100, 200 or more amino acids in length. Biologically active portions of an hVR-1, hVR-2, and rVR-2 protein can be used as targets for developing agents which modulate an hVR-1, hVR-2, and rVR-2 mediated activity, *e.g.*, a pain signaling mechanism.

In one embodiment, a biologically active portion of an hVR-1, hVR-2, and rVR-2 protein comprises at least one transmembrane domain, and/or at least one proline rich domain, and/or at least one ankyrin repeat domain. It is to be understood that a biologically active portion of an hVR-1, hVR-2, and rVR-2 protein of the present invention may contain at least one of the above-identified structural domains. A more biologically active portion of an hVR-1, hVR-2, and rVR-2 protein may contain at least two of the above-identified structural domains. Moreover, other biologically active portions, in which other regions of the protein are deleted, can be prepared by recombinant techniques and evaluated for one or more of the functional activities of a native hVR-1, hVR-2, and rVR-2 protein.

In a embodiment, the hVR-1, hVR-2, and rVR-2 protein has an amino acid sequence shown in SEQ ID NO:2, 5, 8, or 11. In other embodiments, the hVR-1, hVR-2, and rVR-2 protein is substantially homologous to SEQ ID NO:2, 5, 8, or 11, and retains the functional activity of the protein of SEQ ID NO:2, 5, 8, or 11, yet differs in amino acid sequence due to natural allelic variation or mutagenesis, as described in detail in subsection I above. Accordingly, in another embodiment, the hVR-1, hVR-2, and rVR-2 protein is a protein which comprises an amino acid sequence at least about 60%, 65%, 70%, 75%, 80%, 85%, 87%, 90%, 95%, 98% or more homologous to SEQ ID NO:2, 5, 8, or 11.

To determine the percent identity of two amino acid sequences or of two nucleic acid sequences, the sequences are aligned for optimal comparison purposes (*e.g.*, gaps can be introduced in one or both of a first and a second amino acid or nucleic acid sequence for optimal alignment and non-homologous sequences can be disregarded for comparison purposes). In a embodiment, the length of a reference sequence aligned for comparison purposes is at least 30%, preferably at least 40%, more preferably at least 50%, even more preferably at least 60%, and even more preferably at least 70%, 80%, or 90% of the length of the reference sequence (*e.g.*, when aligning a second sequence to the hVR-1, hVR-2, and rVR-2 amino acid sequence of SEQ ID NO:2, 5, 8, or 11, having 177 amino acid residues, at least 80, preferably at least 100, more preferably at least 120,

even more preferably at least 140, and even more preferably at least 150, 160 or 170 amino acid residues are aligned). The amino acid residues or nucleotides at corresponding amino acid positions or nucleotide positions are then compared. When a position in the first sequence is occupied by the same amino acid residue or nucleotide as the corresponding position in the second sequence, then the molecules are identical at that position (as used herein amino acid or nucleic acid "identity" is equivalent to amino acid or nucleic acid "homology"). The percent identity between the two sequences is a function of the number of identical positions shared by the sequences, taking into account the number of gaps, and the length of each gap, which need to be introduced for optimal alignment of the two sequences.

The comparison of sequences and determination of percent identity between two sequences can be accomplished using a mathematical algorithm. In a embodiment, the percent identity between two amino acid sequences is determined using the Needleman and Wunsch (*J. Mol. Biol.* (48):444-453 (1970)) algorithm which has been incorporated into the GAP program in the GCG software package (available at <http://www.gcg.com>), using either a Blossum 62 matrix or a PAM250 matrix, and a gap weight of 16, 14, 12, 10, 8, 6, or 4 and a length weight of 1, 2, 3, 4, 5, or 6. In yet another embodiment, the percent identity between two nucleotide sequences is determined using the GAP program in the GCG software package (available at <http://www.gcg.com>), using a NWSgapdna.CMP matrix and a gap weight of 40, 50, 60, 70, or 80 and a length weight of 1, 2, 3, 4, 5, or 6. In another embodiment, the percent identity between two amino acid or nucleotide sequences is determined using the algorithm of E. Meyers and W. Miller (CABIOS, 4:11-17 (1989)) which has been incorporated into the ALIGN program (version 2.0), using a PAM120 weight residue table, a gap length penalty of 12 and a gap penalty of 4.

The nucleic acid and protein sequences of the present invention can further be used as a "query sequence" to perform a search against public databases to, for example, identify other family members or related sequences. Such searches can be performed using the NBLAST and XBLAST programs (version 2.0) of Altschul, *et al.* (1990) *J. Mol. Biol.* 215:403-10. BLAST nucleotide searches can be performed with the NBLAST program, score = 100, wordlength = 12 to obtain nucleotide sequences homologous to hVR-1, hVR-2, and rVR-2 nucleic acid molecules of the invention. BLAST protein searches can be performed with the XBLAST program, score = 50, wordlength = 3 to obtain amino acid sequences homologous to hVR-1, hVR-2, and rVR-2 protein molecules of the invention. To obtain gapped alignments for comparison purposes, Gapped BLAST can be utilized as described in Altschul *et al.*, (1997) *Nucleic Acids Res.* 25(17):3389-3402. When utilizing BLAST and Gapped BLAST programs, the default parameters of the respective programs (*e.g.*, XBLAST and NBLAST) can be used. See <http://www.ncbi.nlm.nih.gov>.

The invention also provides hVR-1, hVR-2, and rVR-2 chimeric or fusion proteins. As used herein, an hVR-1, hVR-2, and rVR-2 "chimeric protein" or "fusion protein" comprises an hVR-1, hVR-2, and rVR-2 polypeptide operatively linked to a non-hVR-1, hVR-2, and rVR-2 polypeptide. An "hVR-1, hVR-2, and rVR-2 polypeptide" refers to a polypeptide having an amino

acid sequence corresponding to hVR-1, hVR-2, and rVR-2, whereas a "non-hVR-1, non-hVR-2, and non-rVR-2 polypeptide" refers to a polypeptide having an amino acid sequence corresponding to a protein which is not substantially homologous to the hVR-1, hVR-2, and rVR-2 protein, *e.g.*, a protein which is different from the hVR-1, hVR-2, and rVR-2 protein and which is derived from the same or a different organism. Within an hVR-1, hVR-2, and rVR-2 fusion protein the hVR-1, hVR-2, and rVR-2 polypeptide can correspond to all or a portion of an hVR-1, hVR-2, and rVR-2 protein. In a embodiment, an hVR-1, hVR-2, and rVR-2 fusion protein comprises at least one biologically active portion of an hVR-1, hVR-2, and rVR-2 protein. In another embodiment, an hVR-1, hVR-2, and rVR-2 fusion protein comprises at least two biologically active portions of an hVR-1, hVR-2, and rVR-2 protein. Within the fusion protein, the term "operatively linked" is intended to indicate that the hVR-1, hVR-2, and rVR-2 polypeptide and the non-hVR-1, non-hVR-2, and non-rVR-2 polypeptide are fused in-frame to each other. The non-hVR-1, hVR-2, and rVR-2 polypeptide can be fused to the N-terminus or C-terminus of the hVR-1, hVR-2, and rVR-2 polypeptide.

For example, in one embodiment, the fusion protein is a GST-hVR-1, GST-hVR-2, and GST-rVR-2 fusion protein in which the hVR-1, hVR-2, and rVR-2 sequences are fused to the C-terminus of the GST sequences. Such fusion proteins can facilitate the purification of recombinant hVR-1, hVR-2, and rVR-2.

In another embodiment, the fusion protein is an hVR-1, hVR-2, and rVR-2 protein containing a heterologous signal sequence at its N-terminus. In certain host cells (*e.g.*, mammalian host cells), expression and/or secretion of hVR-1, hVR-2, and rVR-2 can be increased through use of a heterologous signal sequence.

The hVR-1, hVR-2, and rVR-2 fusion proteins of the invention can be incorporated into pharmaceutical compositions and administered to a subject *in vivo*. The hVR-1, hVR-2, and rVR-2 fusion proteins can be used to affect the bioavailability of an hVR-1, hVR-2, and rVR-2 substrate. Use of hVR-1, hVR-2, and rVR-2 fusion proteins may be useful therapeutically for the treatment of disorders caused by, for example, (i) aberrant modification or mutation of a gene encoding an hVR-1, hVR-2, and rVR-2 protein; (ii) mis-regulation of the hVR-1, hVR-2, and rVR-2 gene; and (iii) aberrant post-translational modification of an hVR-1, hVR-2, and rVR-2 protein.

Moreover, the hVR-1, hVR-2, and rVR-2-fusion proteins of the invention can be used as immunogens to produce anti-hVR-1, anti-hVR-2, and anti-rVR-2 antibodies in a subject, to purify hVR-1, hVR-2, and rVR-2 ligands and in screening assays to identify molecules which inhibit the interaction of hVR-1, hVR-2, and rVR-2 with an hVR-1, hVR-2, and rVR-2 substrate.

Preferably, an hVR-1, hVR-2, and rVR-2 chimeric or fusion protein of the invention is produced by standard recombinant DNA techniques. For example, DNA fragments coding for the different polypeptide sequences are ligated together in-frame in accordance with conventional techniques, for example by employing blunt-ended or stagger-ended termini for ligation, restriction enzyme digestion to provide for appropriate termini, filling-in of cohesive ends as appropriate,

alkaline phosphatase treatment to avoid undesirable joining, and enzymatic ligation. In another embodiment, the fusion gene can be synthesized by conventional techniques including automated DNA synthesizers. Alternatively, PCR amplification of gene fragments can be carried out using anchor primers which give rise to complementary overhangs between two consecutive gene fragments which can subsequently be annealed and reamplified to generate a chimeric gene sequence (see, for example, *Current Protocols in Molecular Biology*, eds. Ausubel *et al.* John Wiley & Sons: 1992). Moreover, many expression vectors are commercially available that already encode a fusion moiety (*e.g.*, a GST polypeptide). An hVR-1, hVR-2, and rVR-2-encoding nucleic acid can be cloned into such an expression vector such that the fusion moiety is linked in-frame to the hVR-1, hVR-2, and rVR-2 protein.

The present invention also pertains to variants of the hVR-1, hVR-2, and rVR-2 proteins which function as either hVR-1, hVR-2, and rVR-2 agonists (mimetics) or as hVR-1, hVR-2, and rVR-2 antagonists. Variants of the hVR-1, hVR-2, and rVR-2 proteins can be generated by mutagenesis, *e.g.*, discrete point mutation or truncation of an hVR-1, hVR-2, and rVR-2 protein. An agonist of the hVR-1, hVR-2, and rVR-2 proteins can retain substantially the same, or a subset, of the biological activities of the naturally occurring form of an hVR-1, hVR-2, and rVR-2 protein. An antagonist of an hVR-1, hVR-2, and rVR-2 protein can inhibit one or more of the activities of the naturally occurring form of the hVR-1, hVR-2, and rVR-2 protein by, for example, competitively modulating an hVR-1, hVR-2, and rVR-2-mediated activity of an hVR-1, hVR-2, and rVR-2 protein. Thus, specific biological effects can be elicited by treatment with a variant of limited function. In one embodiment, treatment of a subject with a variant having a subset of the biological activities of the naturally occurring form of the protein has fewer side effects in a subject relative to treatment with the naturally occurring form of the hVR-1, hVR-2, and rVR-2 protein.

In one embodiment, variants of an hVR-1, hVR-2, and rVR-2 protein which function as either hVR-1, hVR-2, and rVR-2 agonists (mimetics) or as hVR-1, hVR-2, and rVR-2 antagonists can be identified by screening combinatorial libraries of mutants, *e.g.*, truncation mutants, of an hVR-1, hVR-2, and rVR-2 protein for hVR-1, hVR-2, and rVR-2 protein agonist or antagonist activity. In one embodiment, a variegated library of hVR-1, hVR-2, and rVR-2 variants is generated by combinatorial mutagenesis at the nucleic acid level and is encoded by a variegated gene library. A variegated library of hVR-1, hVR-2, and rVR-2 variants can be produced by, for example, enzymatically ligating a mixture of synthetic oligonucleotides into gene sequences such that a degenerate set of potential hVR-1, hVR-2, and rVR-2 sequences is expressible as individual polypeptides, or alternatively, as a set of larger fusion proteins (*e.g.*, for phage display) containing the set of hVR-1, hVR-2, and rVR-2 sequences therein. There are a variety of methods which can be used to produce libraries of potential hVR-1, hVR-2, and rVR-2 variants from a degenerate oligonucleotide sequence. Chemical synthesis of a degenerate gene sequence can be performed in an automatic DNA synthesizer, and the synthetic gene then ligated into an appropriate expression vector. Use of a degenerate set of genes allows for the provision, in one mixture, of all of the

sequences encoding the desired set of potential hVR-1, hVR-2, and rVR-2 sequences. Methods for synthesizing degenerate oligonucleotides are known in the art (see, *e.g.*, Narang, S.A. (1983) *Tetrahedron* 39:3; Itakura *et al.* (1984) *Annu. Rev. Biochem.* 53:323; Itakura *et al.* (1984) *Science* 198:1056; Ike *et al.* (1983) *Nucleic Acid Res.* 11:477.

5 In addition, libraries of fragments of an hVR-1, hVR-2, and rVR-2 protein coding sequence can be used to generate a variegated population of hVR-1, hVR-2, and rVR-2 fragments for screening and subsequent selection of variants of an hVR-1, hVR-2, and rVR-2 protein. In one embodiment, a library of coding sequence fragments can be generated by treating a double stranded PCR fragment of an hVR-1, hVR-2, and rVR-2 coding sequence with a nuclease under conditions
10 wherein nicking occurs only about once per molecule, denaturing the double stranded DNA, renaturing the DNA to form double stranded DNA which can include sense/antisense pairs from different nicked products, removing single stranded portions from reformed duplexes by treatment with S1 nuclease, and ligating the resulting fragment library into an expression vector. By this method, an expression library can be derived which encodes N-terminal, C-terminal and internal
15 fragments of various sizes of the hVR-1, hVR-2, and rVR-2 protein.

Several techniques are known in the art for screening gene products of combinatorial libraries made by point mutations or truncation, and for screening cDNA libraries for gene products having a selected property. Such techniques are adaptable for rapid screening of the gene libraries generated by the combinatorial mutagenesis of hVR-1, hVR-2, and rVR-2 proteins. The most
20 widely used techniques, which are amenable to high through-put analysis, for screening large gene libraries typically include cloning the gene library into replicable expression vectors, transforming appropriate cells with the resulting library of vectors, and expressing the combinatorial genes under conditions in which detection of a desired activity facilitates isolation of the vector encoding the gene whose product was detected. Recursive ensemble mutagenesis (REM), a new technique
25 which enhances the frequency of functional mutants in the libraries, can be used in combination with the screening assays to identify hVR-1, hVR-2, and rVR-2 variants (Arkin and Yourvan (1992) *Proc. Natl. Acad. Sci. USA* 89:7811-7815; Delgrave *et al.* (1993) *Protein Engineering* 6(3):327-331).

In one embodiment, cell based assays can be exploited to analyze a variegated hVR-1, hVR-
30 2, and rVR-2 library. For example, a library of expression vectors can be transfected into a cell line, *e.g.*, a neuronal cell line, which ordinarily responds to a particular ligand in an hVR-1, hVR-2, and rVR-2-dependent manner. The transfected cells are then contacted with the ligand and the effect of expression of the mutant on signaling by the ligand can be detected, *e.g.*, by measuring intracellular calcium concentration, neuronal membrane depolarization, or the activity of an hVR-1,
35 hVR-2, and rVR-2-regulated transcription factor. Plasmid DNA can then be recovered from the cells which score for inhibition, or alternatively, potentiation of signaling by the ligand, and the individual clones further characterized.

An isolated hVR-1, hVR-2, and rVR-2 protein, or a portion or fragment thereof, can be used as an immunogen to generate antibodies that bind hVR-1, hVR-2, and rVR-2 using standard techniques for polyclonal and monoclonal antibody preparation. A full-length hVR-1, hVR-2, and rVR-2 protein can be used or, alternatively, the invention provides antigenic peptide fragments of hVR-1, hVR-2, and rVR-2 for use as immunogens. The antigenic peptide of hVR-1, hVR-2, and rVR-2 comprises at least 8 amino acid residues of the amino acid sequence shown in SEQ ID NO:2, 5, 8, or 11 and encompasses an epitope of hVR-1, hVR-2, and rVR-2 such that an antibody raised against the peptide forms a specific immune complex with hVR-1, hVR-2, and rVR-2. Preferably, the antigenic peptide comprises at least 10 amino acid residues, more preferably at least 15 amino acid residues, even more preferably at least 20 amino acid residues, and most preferably at least 30 amino acid residues.

Epitopes encompassed by the antigenic peptide are regions of hVR-1, hVR-2, and rVR-2 that are located on the surface of the protein, *e.g.*, hydrophilic regions, as well as regions with high antigenicity (see, for example, Figures 12 and 14).

An hVR-1, hVR-2, and rVR-2 immunogen typically is used to prepare antibodies by immunizing a suitable subject, (*e.g.*, rabbit, goat, mouse or other mammal) with the immunogen. An appropriate immunogenic preparation can contain, for example, recombinantly expressed hVR-1, hVR-2, and rVR-2 protein or a chemically synthesized hVR-1, hVR-2, and rVR-2 polypeptide. The preparation can further include an adjuvant, such as Freund's complete or incomplete adjuvant, or similar immunostimulatory agent. Immunization of a suitable subject with an immunogenic hVR-1, hVR-2, and rVR-2 preparation induces a polyclonal anti-hVR-1, anti-hVR-2, and anti-rVR-2 antibody response.

Accordingly, another aspect of the invention pertains to anti-hVR-1, anti-hVR-2, and anti-rVR-2 antibodies. The term "antibody" as used herein refers to immunoglobulin molecules and immunologically active portions of immunoglobulin molecules, *i.e.*, molecules that contain an antigen binding site which specifically binds (immunoreacts with) an antigen, such as hVR-1, hVR-2, and rVR-2. Examples of immunologically active portions of immunoglobulin molecules include F(ab) and F(ab')₂ fragments which can be generated by treating the antibody with an enzyme such as pepsin. The invention provides polyclonal and monoclonal antibodies that bind hVR-1, hVR-2, and rVR-2. The term "monoclonal antibody" or "monoclonal antibody composition", as used herein, refers to a population of antibody molecules that contain only one species of an antigen binding site capable of immunoreacting with a particular epitope of hVR-1, hVR-2, and rVR-2. A monoclonal antibody composition thus typically displays a single binding affinity for a particular hVR-1, hVR-2, and rVR-2 protein with which it immunoreacts.

Polyclonal anti-hVR-1, anti-hVR-2, and anti-rVR-2 antibodies can be prepared as described above by immunizing a suitable subject with an hVR-1, hVR-2, and rVR-2 immunogen. The anti-hVR-1, anti-hVR-2, and anti-rVR-2 antibody titer in the immunized subject can be monitored over time by standard techniques, such as with an enzyme linked immunosorbent assay (ELISA) using

immobilized hVR-1, hVR-2, and rVR-2. If desired, the antibody molecules directed against hVR-1, hVR-2, and rVR-2 can be isolated from the mammal (e.g., from the blood) and further purified by well known techniques, such as protein A chromatography to obtain the IgG fraction. At an appropriate time after immunization, e.g., when the anti-hVR-1, anti-hVR-2, and anti-rVR-2 antibody titers are highest, antibody-producing cells can be obtained from the subject and used to prepare monoclonal antibodies by standard techniques, such as the hybridoma technique originally described by Kohler and Milstein (1975) *Nature* 256:495-497) (see also, Brown *et al.* (1981) *J. Immunol.* 127:539-46; Brown *et al.* (1980) *J. Biol. Chem.* 255:4980-83; Yeh *et al.* (1976) *Proc. Natl. Acad. Sci. USA* 76:2927-31; and Yeh *et al.* (1982) *Int. J. Cancer* 29:269-75), the more recent human B cell hybridoma technique (Kozbor *et al.* (1983) *Immunol Today* 4:72), the EBV-hybridoma technique (Cole *et al.* (1985), *Monoclonal Antibodies and Cancer Therapy*, Alan R. Liss, Inc., pp. 77-96) or trioma techniques. The technology for producing monoclonal antibody hybridomas is well known (see generally R. H. Kenneth, in *Monoclonal Antibodies: A New Dimension In Biological Analyses*, Plenum Publishing Corp., New York, New York (1980); E. A. Lerner (1981) *Yale J. Biol. Med.*, 54:387402; M. L. Geftter *et al.* (1977) *Somatic Cell Genet.* 3:23136). Briefly, an immortal cell line (typically a myeloma) is fused to lymphocytes (typically splenocytes) from a mammal immunized with an hVR-1, hVR-2, and rVR-2 immunogen as described above, and the culture supernatants of the resulting hybridoma cells are screened to identify a hybridoma producing a monoclonal antibody that binds hVR-1, hVR-2, and rVR-2.

Any of the many well known protocols used for fusing lymphocytes and immortalized cell lines can be applied for the purpose of generating an anti-hVR-1, anti-hVR-2, and anti-rVR-2 monoclonal antibodies (see, e.g., G. Galfre *et al.* (1977) *Nature* 266:55052; Geftter *et al.* *Somatic Cell Genet.*, cited *supra*; Lerner, *Yale J. Biol. Med.*, cited *supra*; Kenneth, *Monoclonal Antibodies*, cited *supra*). Moreover, the ordinarily skilled worker will appreciate that there are many variations of such methods which also would be useful. Typically, the immortal cell line (e.g., a myeloma cell line) is derived from the same mammalian species as the lymphocytes. For example, murine hybridomas can be made by fusing lymphocytes from a mouse immunized with an immunogenic preparation of the present invention with an immortalized mouse cell line. immortal cell lines are mouse myeloma cell lines that are sensitive to culture medium containing hypoxanthine, aminopterin and thymidine ("HAT medium"). Any of a number of myeloma cell lines can be used as a fusion partner according to standard techniques, e.g., the P3-NS1/1-Ag4-1, P3-x63-Ag8.653 or Sp2/O-Ag14 myeloma lines. These myeloma lines are available from ATCC. Typically, HAT-sensitive mouse myeloma cells are fused to mouse splenocytes using polyethylene glycol ("PEG"). Hybridoma cells resulting from the fusion are then selected using HAT medium, which kills unfused and unproductively fused myeloma cells (unfused splenocytes die after several days because they are not transformed). Hybridoma cells producing a monoclonal antibody of the invention are detected by screening the hybridoma culture supernatants for antibodies that bind hVR-1, hVR-2, and rVR-2, e.g., using a standard ELISA assay.

Alternative to preparing monoclonal antibody-secreting hybridomas, a monoclonal anti-hVR-1, anti-hVR-2, and anti-rVR-2 antibody can be identified and isolated by screening a recombinant combinatorial immunoglobulin library (e.g., an antibody phage display library) with hVR-1, hVR-2, and rVR-2 to thereby isolate immunoglobulin library members that bind hVR-1, hVR-2, and rVR-2. Kits for generating and screening phage display libraries are commercially available (e.g., the Pharmacia *Recombinant Phage Antibody System*, Catalog No. 27-9400-01; and the Stratagene *SurfZAP™ Phage Display Kit*, Catalog No. 240612). Additionally, examples of methods and reagents particularly amenable for use in generating and screening antibody display library can be found in, for example, Ladner *et al.* U.S. Patent No. 5,223,409; Kang *et al.* PCT International Publication No. WO 92/18619; Dower *et al.* PCT International Publication No. WO 91/17271; Winter *et al.* PCT International Publication WO 92/20791; Markland *et al.* PCT International Publication No. WO 92/15679; Breitling *et al.* PCT International Publication WO 93/01288; McCafferty *et al.* PCT International Publication No. WO 92/01047; Garrard *et al.* PCT International Publication No. WO 92/09690; Ladner *et al.* PCT International Publication No. WO 90/02809; Fuchs *et al.* (1991) *Bio/Technology* 9:1370-1372; Hay *et al.* (1992) *Hum. Antibod. Hybridomas* 3:81-85; Huse *et al.* (1989) *Science* 246:1275-1281; Griffiths *et al.* (1993) *EMBO J* 12:725-734; Hawkins *et al.* (1992) *J. Mol. Biol.* 226:889-896; Clarkson *et al.* (1991) *Nature* 352:624-628; Gram *et al.* (1992) *Proc. Natl. Acad. Sci. USA* 89:3576-3580; Garrad *et al.* (1991) *Bio/Technology* 9:1373-1377; Hoogenboom *et al.* (1991) *Nuc. Acid Res.* 19:4133-4137; Barbas *et al.* (1991) *Proc. Natl. Acad. Sci. USA* 88:7978-7982; and McCafferty *et al.* *Nature* (1990) 348:552-554.

Additionally, recombinant anti-hVR-1, anti-hVR-2, and anti-rVR-2 antibodies, such as chimeric and humanized monoclonal antibodies, comprising both human and non-human portions, which can be made using standard recombinant DNA techniques, are within the scope of the invention. Such chimeric and humanized monoclonal antibodies can be produced by recombinant DNA techniques known in the art, for example using methods described in Robinson *et al.* International Application No. PCT/US86/02269; Akira, *et al.* European Patent Application 184,187; Taniguchi, M., European Patent Application 171,496; Morrison *et al.* European Patent Application 173,494; Neuberger *et al.* PCT International Publication No. WO 86/01533; Cabilly *et al.* U.S. Patent No. 4,816,567; Cabilly *et al.* European Patent Application 125,023; Better *et al.* (1988) *Science* 240:1041-1043; Liu *et al.* (1987) *Proc. Natl. Acad. Sci. USA* 84:3439-3443; Liu *et al.* (1987) *J. Immunol.* 139:3521-3526; Sun *et al.* (1987) *Proc. Natl. Acad. Sci. USA* 84:214-218; Nishimura *et al.* (1987) *Canc. Res.* 47:999-1005; Wood *et al.* (1985) *Nature* 314:446-449; and Shaw *et al.* (1988) *J. Natl. Cancer Inst.* 80:1553-1559; Morrison, S. L. (1985) *Science* 229:1202-1207; Oi *et al.* (1986) *BioTechniques* 4:214; Winter U.S. Patent 5,225,539; Jones *et al.* (1986) *Nature* 321:552-525; Verhoeyan *et al.* (1988) *Science* 239:1534; and Beidler *et al.* (1988) *J. Immunol.* 141:4053-4060.

An anti-hVR-1, anti-hVR-2, and anti-rVR-2 antibody (e.g., monoclonal antibody) can be used to isolate hVR-1, hVR-2, and rVR-2 by standard techniques, such as affinity chromatography or immunoprecipitation. An anti-hVR-1, anti-hVR-2, and anti-rVR-2 antibody can facilitate the purification of natural hVR-1, hVR-2, and rVR-2 from cells and of recombinantly produced hVR-1, hVR-2, and rVR-2 expressed in host cells. Moreover, an anti-hVR-1, anti-hVR-2, and anti-rVR-2 antibody can be used to detect hVR-1, hVR-2, and rVR-2 protein (e.g., in a cellular lysate or cell supernatant) in order to evaluate the abundance and pattern of expression of the hVR-1, hVR-2, and rVR-2 protein. Anti-hVR-1, anti-hVR-2, and anti-rVR-2 antibodies can be used diagnostically to monitor protein levels in tissue as part of a clinical testing procedure, e.g., to, for example, determine the efficacy of a given treatment regimen. Detection can be facilitated by coupling (*i.e.*, physically linking) the antibody to a detectable substance. Examples of detectable substances include various enzymes, prosthetic groups, fluorescent materials, luminescent materials, bioluminescent materials, and radioactive materials. Examples of suitable enzymes include horseradish peroxidase, alkaline phosphatase, -galactosidase, or acetylcholinesterase; examples of suitable prosthetic group complexes include streptavidin/biotin and avidin/biotin; examples of suitable fluorescent materials include umbelliferone, fluorescein, fluorescein isothiocyanate, rhodamine, dichlorotriazinylamine fluorescein, dansyl chloride or phycoerythrin; an example of a luminescent material includes luminol; examples of bioluminescent materials include luciferase, luciferin, and aequorin, and examples of suitable radioactive material include ^{125}I , ^{131}I , ^{35}S or ^3H .

III. Recombinant Expression Vectors and Host Cells

Another aspect of the invention pertains to vectors, preferably expression vectors, containing a nucleic acid encoding an hVR-1, hVR-2, and rVR-2 protein (or a portion thereof). As used herein, the term "vector" refers to a nucleic acid molecule capable of transporting another nucleic acid to which it has been linked. One type of vector is a "plasmid", which refers to a circular double stranded DNA loop into which additional DNA segments can be ligated. Another type of vector is a viral vector, wherein additional DNA segments can be ligated into the viral genome. Certain vectors are capable of autonomous replication in a host cell into which they are introduced (e.g., bacterial vectors having a bacterial origin of replication and episomal mammalian vectors). Other vectors (e.g., non-episomal mammalian vectors) are integrated into the genome of a host cell upon introduction into the host cell, and thereby are replicated along with the host genome. Moreover, certain vectors are capable of directing the expression of genes to which they are operatively linked. Such vectors are referred to herein as "expression vectors". In general, expression vectors of utility in recombinant DNA techniques are often in the form of plasmids. In the present specification, "plasmid" and "vector" can be used interchangeably as the plasmid is the most commonly used form of vector. However, the invention is intended to include such other forms of expression vectors, such as viral vectors (e.g., replication defective retroviruses, adenoviruses and adeno-associated viruses), which serve equivalent functions.

The recombinant expression vectors of the invention comprise a nucleic acid of the invention in a form suitable for expression of the nucleic acid in a host cell, which means that the recombinant expression vectors include one or more regulatory sequences, selected on the basis of the host cells to be used for expression, which is operatively linked to the nucleic acid sequence to be expressed. Within a recombinant expression vector, "operably linked" is intended to mean that the nucleotide sequence of interest is linked to the regulatory sequence(s) in a manner which allows for expression of the nucleotide sequence (*e.g.*, in an *in vitro* transcription/translation system or in a host cell when the vector is introduced into the host cell). The term "regulatory sequence" is intended to include promoters, enhancers and other expression control elements (*e.g.*, polyadenylation signals). Such regulatory sequences are described, for example, in Goeddel; *Gene Expression Technology: Methods in Enzymology* 185, Academic Press, San Diego, CA (1990). Regulatory sequences include those which direct constitutive expression of a nucleotide sequence in many types of host cells and those which direct expression of the nucleotide sequence only in certain host cells (*e.g.*, tissue-specific regulatory sequences). It will be appreciated by those skilled in the art that the design of the expression vector can depend on such factors as the choice of the host cell to be transformed, the level of expression of protein desired, and the like. The expression vectors of the invention can be introduced into host cells to thereby produce proteins or peptides, including fusion proteins or peptides, encoded by nucleic acids as described herein (*e.g.*, hVR-1, hVR-2, and rVR-2 proteins, mutant forms of hVR-1, hVR-2, and rVR-2 proteins, fusion proteins, and the like).

The recombinant expression vectors of the invention can be designed for expression of hVR-1, hVR-2, and rVR-2 proteins in prokaryotic or eukaryotic cells. For example, hVR-1, hVR-2, and rVR-2 proteins can be expressed in bacterial cells such as *E. coli*, insect cells (using baculovirus expression vectors) yeast cells or mammalian cells. Suitable host cells are discussed further in Goeddel, *Gene Expression Technology: Methods in Enzymology* 185, Academic Press, San Diego, CA (1990). Alternatively, the recombinant expression vector can be transcribed and translated *in vitro*, for example using T7 promoter regulatory sequences and T7 polymerase.

Expression of proteins in prokaryotes is most often carried out in *E. coli* with vectors containing constitutive or inducible promoters directing the expression of either fusion or non-fusion proteins. Fusion vectors add a number of amino acids to a protein encoded therein, usually to the amino terminus of the recombinant protein. Such fusion vectors typically serve three purposes: 1) to increase expression of recombinant protein; 2) to increase the solubility of the recombinant protein; and 3) to aid in the purification of the recombinant protein by acting as a ligand in affinity purification. Often, in fusion expression vectors, a proteolytic cleavage site is introduced at the junction of the fusion moiety and the recombinant protein to enable separation of the recombinant protein from the fusion moiety subsequent to purification of the fusion protein. Such enzymes, and their cognate recognition sequences, include Factor Xa, thrombin and enterokinase. Typical fusion expression vectors include pGEX (Pharmacia Biotech Inc; Smith,

D.B. and Johnson, K.S. (1988) *Gene* 67:31-40), pMAL (New England Biolabs, Beverly, MA) and pRIT5 (Pharmacia, Piscataway, NJ) which fuse glutathione S-transferase (GST), maltose E binding protein, or protein A, respectively, to the target recombinant protein.

Purified fusion proteins can be utilized in hVR-1, hVR-2, and rVR-2 activity assays, (e.g., direct assays or competitive assays described in detail below), or to, for example, generate antibodies specific for hVR-1, hVR-2, and rVR-2 proteins. In a embodiment, an hVR-1, hVR-2, and rVR-2 fusion protein expressed in a retroviral expression vector of the present invention can be utilized to infect bone marrow cells which are subsequently transplanted into irradiated recipients. The pathology of the subject recipient is then examined after sufficient time has passed (e.g., six (6) weeks).

Examples of suitable inducible non-fusion *E. coli* expression vectors include pTrc (Amann *et al.*, (1988) *Gene* 69:301-315) and pET 11d (Studier *et al.*, *Gene Expression Technology: Methods in Enzymology* 185, Academic Press, San Diego, California (1990) 60-89). Target gene expression from the pTrc vector relies on host RNA polymerase transcription from a hybrid trp-lac fusion promoter. Target gene expression from the pET 11d vector relies on transcription from a T7 gn10-lac fusion promoter mediated by a coexpressed viral RNA polymerase (T7 gn1). This viral polymerase is supplied by host strains BL21(DE3) or HMS174(DE3) from a resident prophage harboring a T7 gn1 gene under the transcriptional control of the lacUV 5 promoter.

One strategy to maximize recombinant protein expression in *E. coli* is to express the protein in a host bacteria with an impaired capacity to proteolytically cleave the recombinant protein (Gottesman, S., *Gene Expression Technology: Methods in Enzymology* 185, Academic Press, San Diego, California (1990) 119-128). Another strategy is to alter the nucleic acid sequence of the nucleic acid to be inserted into an expression vector so that the individual codons for each amino acid are those preferentially utilized in *E. coli* (Wada *et al.*, (1992) *Nucleic Acids Res.* 20:2111-2118). Such alteration of nucleic acid sequences of the invention can be carried out by standard DNA synthesis techniques.

In another embodiment, the hVR-1, hVR-2, and rVR-2 expression vector is a yeast expression vector. Examples of vectors for expression in yeast *S. cerevisiae* include pYepSec1 (Baldari, *et al.*, (1987) *Embo J.* 6:229-234), pMFa (Kurjan and Herskowitz, (1982) *Cell* 30:933-943), pJRY88 (Schultz *et al.*, (1987) *Gene* 54:113-123), pYES2 (Invitrogen Corporation, San Diego, CA), and picZ (Invitrogen Corp, San Diego, CA).

Alternatively, hVR-1, hVR-2, and rVR-2 proteins can be expressed in insect cells using baculovirus expression vectors. Baculovirus vectors available for expression of proteins in cultured insect cells (e.g., Sf9 cells) include the pAc series (Smith *et al.* (1983) *Mol. Cell Biol.* 3:2156-2165) and the pVL series (Lucklow and Summers (1989) *Virology* 170:31-39).

In yet another embodiment, a nucleic acid of the invention is expressed in mammalian cells using a mammalian expression vector. Examples of mammalian expression vectors include pCDM8 (Seed, B. (1987) *Nature* 329:840) and pMT2PC (Kaufman *et al.* (1987) *EMBO J.* 6:187-

195). When used in mammalian cells, the expression vector's control functions are often provided by viral regulatory elements. For example, commonly used promoters are derived from polyoma, Adenovirus 2, cytomegalovirus and Simian Virus 40. For other suitable expression systems for both prokaryotic and eukaryotic cells see chapters 16 and 17 of Sambrook, J., Fritsch, E. F., and Maniatis, T. *Molecular Cloning: A Laboratory Manual*. 2nd, ed., Cold Spring Harbor Laboratory, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, 1989.

In another embodiment, the recombinant mammalian expression vector is capable of directing expression of the nucleic acid preferentially in a particular cell type (e.g., tissue-specific regulatory elements are used to express the nucleic acid). Tissue-specific regulatory elements are known in the art. Non-limiting examples of suitable tissue-specific promoters include the albumin promoter (liver-specific; Pinkert *et al.* (1987) *Genes Dev.* 1:268-277), lymphoid-specific promoters (Calame and Eaton (1988) *Adv. Immunol.* 43:235-275), in particular promoters of T cell receptors (Winoto and Baltimore (1989) *EMBO J.* 8:729-733) and immunoglobulins (Banerji *et al.* (1983) *Cell* 33:729-740; Queen and Baltimore (1983) *Cell* 33:741-748), neuron-specific promoters (e.g., the neurofilament promoter; Byrne and Ruddle (1989) *Proc. Natl. Acad. Sci. USA* 86:5473-5477), pancreas-specific promoters (Edlund *et al.* (1985) *Science* 230:912-916), and mammary gland-specific promoters (e.g., milk whey promoter; U.S. Patent No. 4,873,316 and European Application Publication No. 264,166). Developmentally-regulated promoters are also encompassed, for example the murine hox promoters (Kessel and Gruss (1990) *Science* 249:374-379) and the α -fetoprotein promoter (Campes and Tilghman (1989) *Genes Dev.* 3:537-546).

The expression characteristics of an endogenous hVR-1, hVR-2, and rVR-2 gene within a cell line or microorganism may be modified by inserting a heterologous DNA regulatory element into the genome of a stable cell line or cloned microorganism such that the inserted regulatory element is operatively linked with the endogenous hVR-1, hVR-2, and rVR-2 gene. For example, an endogenous hVR-1, hVR-2, and rVR-2 gene which is normally "transcriptionally silent", i.e., a hVR-1, hVR-2, and rVR-2 gene which is normally not expressed, or is expressed only at very low levels in a cell line or microorganism, may be activated by inserting a regulatory element which is capable of promoting the expression of a normally expressed gene product in that cell line or microorganism. Alternatively, a transcriptionally silent, endogenous hVR-1, hVR-2, and rVR-2 gene, may be activated by insertion of a promiscuous regulatory element that works across cell types.

A heterologous regulatory element may be inserted into a stable cell line or cloned microorganism, such that it is operatively linked with an endogenous hVR-1, hVR-2, and rVR-2 gene, using techniques, such as targeted homologous recombination, which are well known to those of skill in the art, and described e.g., in Chappel, U.S. Patent No.: 5,272,071; PCT publication No. WO 91/06667, published May 16, 1991.

The invention further provides a recombinant expression vector comprising a DNA molecule of the invention cloned into the expression vector in an antisense orientation. That is, the

DNA molecule is operatively linked to a regulatory sequence in a manner which allows for expression (by transcription of the DNA molecule) of an RNA molecule which is antisense to hVR-1, hVR-2, and rVR-2 mRNA. Regulatory sequences operatively linked to a nucleic acid cloned in the antisense orientation can be chosen which direct the continuous expression of the antisense RNA molecule in a variety of cell types, for instance viral promoters and/or enhancers, or regulatory sequences can be chosen which direct constitutive, tissue specific or cell type specific expression of antisense RNA. The antisense expression vector can be in the form of a recombinant plasmid, phagemid or attenuated virus in which antisense nucleic acids are produced under the control of a high efficiency regulatory region, the activity of which can be determined by the cell type into which the vector is introduced. For a discussion of the regulation of gene expression using antisense genes see Weintraub, H. *et al.*, Antisense RNA as a molecular tool for genetic analysis, *Reviews - Trends in Genetics*, Vol. 1(1) 1986.

Another aspect of the invention pertains to host cells into which an hVR-1, hVR-2, and rVR-2 nucleic acid molecule of the invention is introduced, *e.g.*, an hVR-1, hVR-2, and rVR-2 nucleic acid molecule within a recombinant expression vector or an hVR-1, hVR-2, and rVR-2 nucleic acid molecule containing sequences which allow it to homologously recombine into a specific site of the host cell's genome. The terms "host cell" and "recombinant host cell" are used interchangeably herein. It is understood that such terms refer not only to the particular subject cell but to the progeny or potential progeny of such a cell. Because certain modifications may occur in succeeding generations due to either mutation or environmental influences, such progeny may not, in fact, be identical to the parent cell, but are still included within the scope of the term as used herein.

A host cell can be any prokaryotic or eukaryotic cell. For example, an hVR-1, hVR-2, and rVR-2 protein can be expressed in bacterial cells such as *E. coli*, insect cells, yeast or mammalian cells (such as Chinese hamster ovary cells (CHO) or COS cells). Other suitable host cells are known to those skilled in the art.

Vector DNA can be introduced into prokaryotic or eukaryotic cells via conventional transformation or transfection techniques. As used herein, the terms "transformation" and "transfection" are intended to refer to a variety of art-recognized techniques for introducing foreign nucleic acid (*e.g.*, DNA) into a host cell, including calcium phosphate or calcium chloride co-precipitation, DEAE-dextran-mediated transfection, lipofection, or electroporation. Suitable methods for transforming or transfecting host cells can be found in Sambrook, *et al.* (*Molecular Cloning: A Laboratory Manual*. 2nd, ed., Cold Spring Harbor Laboratory, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, 1989), and other laboratory manuals.

For stable transfection of mammalian cells, it is known that, depending upon the expression vector and transfection technique used, only a small fraction of cells may integrate the foreign DNA into their genome. In order to identify and select these integrants, a gene that encodes a selectable marker (*e.g.*, resistance to antibiotics) is generally introduced into the host cells along

with the gene of interest. selectable markers include those which confer resistance to drugs, such as G418, hygromycin and methotrexate. Nucleic acid encoding a selectable marker can be introduced into a host cell on the same vector as that encoding an hVR-1, hVR-2, and rVR-2 protein or can be introduced on a separate vector. Cells stably transfected with the introduced nucleic acid can be identified by drug selection (*e.g.*, cells that have incorporated the selectable marker gene will survive, while the other cells die).

A host cell of the invention, such as a prokaryotic or eukaryotic host cell in culture, can be used to produce (*i.e.*, express) an hVR-1, hVR-2, and rVR-2 protein. Accordingly, the invention further provides methods for producing an hVR-1, hVR-2, and rVR-2 protein using the host cells of the invention. In one embodiment, the method comprises culturing the host cell of invention (into which a recombinant expression vector encoding an hVR-1, hVR-2, and rVR-2 protein has been introduced) in a suitable medium such that an hVR-1, hVR-2, and rVR-2 protein is produced. In another embodiment, the method further comprises isolating an hVR-1, hVR-2, and rVR-2 protein from the medium or the host cell.

The host cells of the invention can also be used to produce non-human transgenic animals. For example, in one embodiment, a host cell of the invention is a fertilized oocyte or an embryonic stem cell into which hVR-1, hVR-2, and rVR-2-coding sequences have been introduced. Such host cells can then be used to create non-human transgenic animals in which exogenous hVR-1, hVR-2, and rVR-2 sequences have been introduced into their genome or homologous recombinant animals in which endogenous hVR-1, hVR-2, and rVR-2 sequences have been altered. Such animals are useful for studying the function and/or activity of an hVR-1, hVR-2, and rVR-2 and for identifying and/or evaluating modulators of hVR-1, hVR-2, and rVR-2 activity. As used herein, a "transgenic animal" is a non-human animal, preferably a mammal, more preferably a rodent such as a rat or mouse, in which one or more of the cells of the animal includes a transgene. Other examples of transgenic animals include non-human primates, sheep, dogs, cows, goats, chickens, amphibians, and the like. A transgene is exogenous DNA which is integrated into the genome of a cell from which a transgenic animal develops and which remains in the genome of the mature animal, thereby directing the expression of an encoded gene product in one or more cell types or tissues of the transgenic animal. As used herein, a "homologous recombinant animal" is a non-human animal, preferably a mammal, more preferably a mouse, in which an endogenous hVR-1, hVR-2, and rVR-2 gene has been altered by homologous recombination between the endogenous gene and an exogenous DNA molecule introduced into a cell of the animal, *e.g.*, an embryonic cell of the animal, prior to development of the animal.

A transgenic animal of the invention can be created by introducing an hVR-1, hVR-2, and rVR-2-encoding nucleic acid into the male pronuclei of a fertilized oocyte, *e.g.*, by microinjection, retroviral infection, and allowing the oocyte to develop in a pseudopregnant female foster animal. The hVR-1, hVR-2, and rVR-2 cDNA sequence of SEQ ID NO:1, 3, 5, 7 or 9 can be introduced as a transgene into the genome of a non-human animal. Alternatively, a nonhuman homologue of a

hVR-2 gene, such as a mouse or rat hVR-2, *e.g.*, the rVR-2 gene, can be used as a transgene. Alternatively, an hVR-1, hVR-2, and rVR-2 gene homologue, such as another member of the Capsaicin/Vanilloid family, can be isolated based on hybridization to the hVR-1, hVR-2, and rVR-2 cDNA sequences of SEQ ID NO:1, 3, 4, 6, 7, 9, 10, or 12, (described further in subsection I above) and used as a transgene. Intronic sequences and polyadenylation signals can also be included in the transgene to increase the efficiency of expression of the transgene. A tissue-specific regulatory sequence(s) can be operably linked to an hVR-1, hVR-2, and rVR-2 transgene to direct expression of an hVR-1, hVR-2, and rVR-2 protein to particular cells. Methods for generating transgenic animals via embryo manipulation and microinjection, particularly animals such as mice, have become conventional in the art and are described, for example, in U.S. Patent Nos. 4,736,866 and 4,870,009, both by Leder *et al.*, U.S. Patent No. 4,873,191 by Wagner *et al.* and in Hogan, B., *Manipulating the Mouse Embryo*, (Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., 1986). Similar methods are used for production of other transgenic animals. A transgenic founder animal can be identified based upon the presence of an hVR-1, hVR-2, and rVR-2 transgene in its genome and/or expression of hVR-1, hVR-2, and rVR-2 mRNA in tissues or cells of the animals. A transgenic founder animal can then be used to breed additional animals carrying the transgene. Moreover, transgenic animals carrying a transgene encoding an hVR-1, hVR-2, and rVR-2 protein can further be bred to other transgenic animals carrying other transgenes.

To create a homologous recombinant animal, a vector is prepared which contains at least a portion of an hVR-1, hVR-2, and rVR-2 gene into which a deletion, addition or substitution has been introduced to thereby alter, *e.g.*, functionally disrupt, the hVR-1, hVR-2, and rVR-2 gene. The VR-1 or VR-2 gene can be a human gene (*e.g.*, the cDNA of SEQ ID NO:1, 3, 5, 4, 6, 7, or 9), but more preferably, is a non-human homologue of a hVR-1 and hVR-2 gene (*e.g.*, the cDNA of SEQ ID NO:10 or 12, or a cDNA isolated by stringent hybridization with the nucleotide sequence of SEQ ID NO: 1, 3, 5, 4, 6, 7, or 9). For example, a mouse VR-2 gene can be used to construct a homologous recombination nucleic acid molecule, *e.g.*, a vector, suitable for altering an endogenous VR-2 gene in the mouse genome. In an embodiment, the homologous recombination nucleic acid molecule is designed such that, upon homologous recombination, the endogenous hVR-1, hVR-2, and rVR-2 gene is functionally disrupted (*i.e.*, no longer encodes a functional protein; also referred to as a "knock out" vector). Alternatively, the homologous recombination nucleic acid molecule can be designed such that, upon homologous recombination, the endogenous hVR-1, hVR-2, and rVR-2 gene is mutated or otherwise altered but still encodes functional protein (*e.g.*, the upstream regulatory region can be altered to thereby alter the expression of the endogenous hVR-1, hVR-2, and rVR-2 protein). In the homologous recombination nucleic acid molecule, the altered portion of the hVR-1, hVR-2, and rVR-2 gene is flanked at its 5' and 3' ends by additional nucleic acid sequence of the hVR-1, hVR-2, and rVR-2 gene to allow for homologous recombination to occur between the exogenous hVR-1, hVR-2, and rVR-2 gene carried by the homologous recombination nucleic acid molecule and an endogenous hVR-1, hVR-2, and rVR-2

gene in a cell, *e.g.*, an embryonic stem cell. The additional flanking hVR-1, hVR-2, and rVR-2 nucleic acid sequence is of sufficient length for successful homologous recombination with the endogenous gene. Typically, several kilobases of flanking DNA (both at the 5' and 3' ends) are included in the homologous recombination nucleic acid molecule (see, *e.g.*, Thomas, K.R. and
5 Capecchi, M. R. (1987) *Cell* 51:503 for a description of homologous recombination vectors). The homologous recombination nucleic acid molecule is introduced into a cell, *e.g.*, an embryonic stem cell line (*e.g.*, by electroporation) and cells in which the introduced hVR-1, hVR-2, and rVR-2 gene has homologously recombined with the endogenous hVR-1, hVR-2, and rVR-2 gene are selected (see *e.g.*, Li, E. *et al.* (1992) *Cell* 69:915). The selected cells can then be injected into a blastocyst of
10 an animal (*e.g.*, a mouse) to form aggregation chimeras (see *e.g.*, Bradley, A. in *Teratocarcinomas and Embryonic Stem Cells: A Practical Approach*, E.J. Robertson, ed. (IRL, Oxford, 1987) pp. 113-152). A chimeric embryo can then be implanted into a suitable pseudopregnant female foster animal and the embryo brought to term. Progeny harboring the homologously recombined DNA in their germ cells can be used to breed animals in which all cells of the animal contain the homologously recombined DNA by germline transmission of the transgene. Methods for constructing homologous recombination nucleic acid molecules, *e.g.*, vectors, or homologous recombinant animals are described further in Bradley, A. (1991) *Current Opinion in Biotechnology* 2:823-829 and in PCT International Publication Nos.: WO 90/11354 by Le Mouellec *et al.*; WO 91/01140 by Smithies *et al.*; WO 92/0968 by Zijlstra *et al.*; and WO 93/04169 by Berns *et al.*

20 In another embodiment, transgenic non-humans animals can be produced which contain selected systems which allow for regulated expression of the transgene. One example of such a system is the *cre/loxP* recombinase system of bacteriophage P1. For a description of the *cre/loxP* recombinase system, see, *e.g.*, Lakso *et al.* (1992) *Proc. Natl. Acad. Sci. USA* 89:6232-6236. Another example of a recombinase system is the FLP recombinase system of *Saccharomyces cerevisiae* (O'Gorman *et al.* (1991) *Science* 251:1351-1355. If a *cre/loxP* recombinase system is used to regulate expression of the transgene, animals containing transgenes encoding both the *Cre* recombinase and a selected protein are required. Such animals can be provided through the construction of "double" transgenic animals, *e.g.*, by mating two transgenic animals, one containing a transgene encoding a selected protein and the other containing a transgene encoding a
25 recombinase.
30

Clones of the non-human transgenic animals described herein can also be produced according to the methods described in Wilmut, I. *et al.* (1997) *Nature* 385:810-813 and PCT International Publication Nos. WO 97/07668 and WO 97/07669. In brief, a cell, *e.g.*, a somatic cell, from the transgenic animal can be isolated and induced to exit the growth cycle and enter G₀
35 phase. The quiescent cell can then be fused, *e.g.*, through the use of electrical pulses, to an enucleated oocyte from an animal of the same species from which the quiescent cell is isolated. The reconstructed oocyte is then cultured such that it develops to morula or blastocyte and then

transferred to pseudopregnant female foster animal. The offspring borne of this female foster animal will be a clone of the animal from which the cell, *e.g.*, the somatic cell, is isolated.

IV. Pharmaceutical Compositions

5 The hVR-1, hVR-2, and rVR-2 nucleic acid molecules, fragments of hVR-1, hVR-2, and rVR-2 proteins, and anti-hVR-1, anti-hVR-2, and anti-rVR-2 antibodies (also referred to herein as "active compounds") of the invention can be incorporated into pharmaceutical compositions suitable for administration. Such compositions typically comprise the nucleic acid molecule, protein, or antibody and a pharmaceutically acceptable carrier. As used herein the language
10 "pharmaceutically acceptable carrier" is intended to include any and all solvents, dispersion media, coatings, antibacterial and antifungal agents, isotonic and absorption delaying agents, and the like, compatible with pharmaceutical administration. The use of such media and agents for pharmaceutically active substances is well known in the art. Except insofar as any conventional media or agent is incompatible with the active compound, use thereof in the compositions is contemplated. Supplementary active compounds can also be incorporated into the compositions.

15 A pharmaceutical composition of the invention is formulated to be compatible with its intended route of administration. Examples of routes of administration include parenteral, *e.g.*, intravenous, intradermal, subcutaneous, oral (*e.g.*, inhalation), transdermal (topical), transmucosal, and rectal administration. Solutions or suspensions used for parenteral, intradermal, or
20 subcutaneous application can include the following components: a sterile diluent such as water for injection, saline solution, fixed oils, polyethylene glycols, glycerine, propylene glycol or other synthetic solvents; antibacterial agents such as benzyl alcohol or methyl parabens; antioxidants such as ascorbic acid or sodium bisulfite; chelating agents such as ethylenediaminetetraacetic acid; buffers such as acetates, citrates or phosphates and agents for the adjustment of tonicity such as
25 sodium chloride or dextrose. pH can be adjusted with acids or bases, such as hydrochloric acid or sodium hydroxide. The parenteral preparation can be enclosed in ampoules, disposable syringes or multiple dose vials made of glass or plastic.

30 Pharmaceutical compositions suitable for injectable use include sterile aqueous solutions (where water soluble) or dispersions and sterile powders for the extemporaneous preparation of sterile injectable solutions or dispersion. For intravenous administration, suitable carriers include physiological saline, bacteriostatic water, Cremophor EL™ (BASF, Parsippany, NJ) or phosphate buffered saline (PBS). In all cases, the composition must be sterile and should be fluid to the extent that easy syringability exists. It must be stable under the conditions of manufacture and storage and must be preserved against the contaminating action of microorganisms such as bacteria and fungi.
35 The carrier can be a solvent or dispersion medium containing, for example, water, ethanol, polyol (for example, glycerol, propylene glycol, and liquid polyethylene glycol, and the like), and suitable mixtures thereof. The proper fluidity can be maintained, for example, by the use of a coating such as lecithin, by the maintenance of the required particle size in the case of dispersion

and by the use of surfactants. Prevention of the action of microorganisms can be achieved by various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, ascorbic acid, thimerosal, and the like. In many cases, it will be preferable to include isotonic agents, for example, sugars, polyalcohols such as manitol, sorbitol, sodium chloride in the composition.

- 5 Prolonged absorption of the injectable compositions can be brought about by including in the composition an agent which delays absorption, for example, aluminum monostearate and gelatin.

10 Sterile injectable solutions can be prepared by incorporating the active compound (*e.g.*, a fragment of an hVR-1, hVR-2, and rVR-2 protein or an anti-hVR-1, anti-hVR-2, and anti-rVR-2 antibody) in the required amount in an appropriate solvent with one or a combination of ingredients enumerated above, as required, followed by filtered sterilization. Generally, dispersions are prepared by incorporating the active compound into a sterile vehicle which contains a basic dispersion medium and the required other ingredients from those enumerated above. In the case of sterile powders for the preparation of sterile injectable solutions, the methods of preparation are vacuum drying and freeze-drying which yields a powder of the active ingredient plus any additional desired ingredient from a previously sterile-filtered solution thereof.

15 Oral compositions generally include an inert diluent or an edible carrier. They can be enclosed in gelatin capsules or compressed into tablets. For the purpose of oral therapeutic administration, the active compound can be incorporated with excipients and used in the form of tablets, troches, or capsules. oral compositions can also be prepared using a fluid carrier for use as a mouthwash, wherein the compound in the fluid carrier is applied orally and swished and expectorated or swallowed. Pharmaceutically compatible binding agents, and/or adjuvant materials can be included as part of the composition. The tablets, pills, capsules, troches and the like can contain any of the following ingredients, or compounds of a similar nature: a binder such as microcrystalline cellulose, gum tragacanth or gelatin; an excipient such as starch or lactose, a disintegrating agent such as alginic acid, Primogel, or corn starch; a lubricant such as magnesium stearate or Sterotes; a glidant such as colloidal silicon dioxide; a sweetening agent such as sucrose or saccharin; or a flavoring agent such as peppermint, methyl salicylate, or orange flavoring.

25 For administration by inhalation, the compounds are delivered in the form of an aerosol spray from pressured container or dispenser which contains a suitable propellant, *e.g.*, a gas such as carbon dioxide, or a nebulizer.

30 Systemic administration can also be by transmucosal or transdermal means. For transmucosal or transdermal administration, penetrants appropriate to the barrier to be permeated are used in the formulation. Such penetrants are generally known in the art, and include, for example, for transmucosal administration, detergents, bile salts, and fusidic acid derivatives.

35 Transmucosal administration can be accomplished through the use of nasal sprays or suppositories. For transdermal administration, the active compounds are formulated into ointments, salves, gels, or creams as generally known in the art.

The compounds can also be prepared in the form of suppositories (*e.g.*, with conventional suppository bases such as cocoa butter and other glycerides) or retention enemas for rectal delivery.

In one embodiment, the active compounds are prepared with carriers that will protect the compound against rapid elimination from the body, such as a controlled release formulation, including implants and microencapsulated delivery systems. Biodegradable, biocompatible polymers can be used, such as ethylene vinyl acetate, polyanhydrides, polyglycolic acid, collagen, polyorthoesters, and polylactic acid. Methods for preparation of such formulations will be apparent to those skilled in the art. The materials can also be obtained commercially from Alza Corporation and Nova Pharmaceuticals, Inc. Liposomal suspensions (including liposomes targeted to infected cells with monoclonal antibodies to viral antigens) can also be used as pharmaceutically acceptable carriers. These can be prepared according to methods known to those skilled in the art, for example, as described in U.S. Patent No. 4,522,811.

It is especially advantageous to formulate oral or parenteral compositions in dosage unit form for ease of administration and uniformity of dosage. Dosage unit form as used herein refers to physically discrete units suited as unitary dosages for the subject to be treated; each unit containing a predetermined quantity of active compound calculated to produce the desired therapeutic effect in association with the required pharmaceutical carrier. The specification for the dosage unit forms of the invention are dictated by and directly dependent on the unique characteristics of the active compound and the particular therapeutic effect to be achieved, and the limitations inherent in the art of compounding such an active compound for the treatment of individuals.

Toxicity and therapeutic efficacy of such compounds can be determined by standard pharmaceutical procedures in cell cultures or experimental animals, *e.g.*, for determining the LD50 (the dose lethal to 50% of the population) and the ED50 (the dose therapeutically effective in 50% of the population). The dose ratio between toxic and therapeutic effects is the therapeutic index and it can be expressed as the ratio LD50/ED50. Compounds which exhibit large therapeutic indices are preferred. While compounds that exhibit toxic side effects may be used, care should be taken to design a delivery system that targets such compounds to the site of affected tissue in order to minimize potential damage to uninfected cells and, thereby, reduce side effects.

The data obtained from the cell culture assays and animal studies can be used in formulating a range of dosage for use in humans. The dosage of such compounds lies preferably within a range of circulating concentrations that include the ED50 with little or no toxicity. The dosage may vary within this range depending upon the dosage form employed and the route of administration utilized. For any compound used in the method of the invention, the therapeutically effective dose can be estimated initially from cell culture assays. A dose may be formulated in animal models to achieve a circulating plasma concentration range that includes the IC50 (*i.e.*, the concentration of the test compound which achieves a half-maximal inhibition of symptoms) as determined in cell culture. Such information can be used to more accurately determine useful doses in humans. Levels in plasma may be measured, for example, by high performance liquid chromatography.

As defined herein, a therapeutically effective amount of protein or polypeptide (*i.e.*, an effective dosage) ranges from about 0.001 to 30 mg/kg body weight, preferably about 0.01 to 25 mg/kg body weight, more preferably about 0.1 to 20 mg/kg body weight, and even more preferably about 1 to 10 mg/kg, 2 to 9 mg/kg, 3 to 8 mg/kg, 4 to 7 mg/kg, or 5 to 6 mg/kg body weight. The skilled artisan will appreciate that certain factors may influence the dosage required to effectively treat a subject, including but not limited to the severity of the disease or disorder, previous treatments, the general health and/or age of the subject, and other diseases present. Moreover, treatment of a subject with a therapeutically effective amount of a protein, polypeptide, or antibody can include a single treatment or, preferably, can include a series of treatments. In a preferred example, a subject is treated with antibody, protein, or polypeptide in the range of between about 0.1 to 20 mg/kg body weight, one time per week for between about 1 to 10 weeks, preferably between 2 to 8 weeks, more preferably between about 3 to 7 weeks, and even more preferably for about 4, 5, or 6 weeks. It will also be appreciated that the effective dosage of antibody, protein, or polypeptide used for treatment may increase or decrease over the course of a particular treatment. Changes in dosage may result and become apparent from the results of diagnostic assays as described herein.

The present invention encompasses agents which modulate expression or activity. An agent may, for example, be a small molecule. For example, such small molecules include, but are not limited to, peptides, peptidomimetics, amino acids, amino acid analogs, polynucleotides, polynucleotide analogs, nucleotides, nucleotide analogs, organic or inorganic compounds (*i.e.*, including heteroorganic and organometallic compounds) having a molecular weight less than about 10,000 grams per mole, organic or inorganic compounds having a molecular weight less than about 5,000 grams per mole, organic or inorganic compounds having a molecular weight less than about 1,000 grams per mole, organic or inorganic compounds having a molecular weight less than about 500 grams per mole, and salts, esters, and other pharmaceutically acceptable forms of such compounds.

It is understood that appropriate doses of small molecule agents depends upon a number of factors within the ken of the ordinarily skilled physician, veterinarian, or researcher. The dose(s) of the small molecule will vary, for example, depending upon the identity, size, and condition of the subject or sample being treated, further depending upon the route by which the composition is to be administered, if applicable, and the effect which the practitioner desires the small molecule to have upon the nucleic acid or polypeptide of the invention.

Exemplary doses include milligram or microgram amounts of the small molecule per kilogram of subject or sample weight (*e.g.*, about 1 microgram per kilogram to about 500 milligrams per kilogram, about 100 micrograms per kilogram to about 5 milligrams per kilogram, or about 1 microgram per kilogram to about 50 micrograms per kilogram. It is furthermore understood that appropriate doses of a small molecule depend upon the potency of the small

molecule with respect to the expression or activity to be modulated. Such appropriate doses may be determined using the assays described herein.

When one or more of these small molecules is to be administered to an animal (*e.g.*, a human) in order to modulate expression or activity of a polypeptide or nucleic acid of the invention, a physician, veterinarian, or researcher may, for example, prescribe a relatively low dose at first, subsequently increasing the dose until an appropriate response is obtained. In addition, it is understood that the specific dose level for any particular animal subject will depend upon a variety of factors including the activity of the specific compound employed, the age, body weight, general health, gender, and diet of the subject, the time of administration, the route of administration, the rate of excretion, any drug combination, and the degree of expression or activity to be modulated.

The nucleic acid molecules of the invention can be inserted into vectors and used as gene therapy vectors. Gene therapy vectors can be delivered to a subject by, for example, intravenous injection, local administration (see U.S. Patent 5,328,470) or by stereotactic injection (see *e.g.*, Chen *et al.* (1994) *Proc. Natl. Acad. Sci. USA* 91:3054-3057). The pharmaceutical preparation of the gene therapy vector can include the gene therapy vector in an acceptable diluent, or can comprise a slow release matrix in which the gene delivery vehicle is imbedded. Alternatively, where the complete gene delivery vector can be produced intact from recombinant cells, *e.g.*, retroviral vectors, the pharmaceutical preparation can include one or more cells which produce the gene delivery system.

The pharmaceutical compositions can be included in a container, pack, or dispenser together with instructions for administration.

V. Uses and Methods of the Invention

The nucleic acid molecules, proteins, protein homologues, and antibodies described herein can be used in one or more of the following methods: a) screening assays; b) predictive medicine (*e.g.*, diagnostic assays, prognostic assays, monitoring clinical trials, and pharmacogenetics); and c) methods of treatment (*e.g.*, therapeutic and prophylactic). As described herein, an hVR-1, hVR-2, and rVR-2 protein of the invention has one or more of the following activities: (1) it interacts with a non-hVR-1, non-hVR-2, and non-rVR-2 protein molecule, *e.g.*, a vanilloid compound such as capsaicin; (2) it modulates intracellular calcium concentration; (3) it activates an hVR-1, hVR-2, and rVR-2-dependent signal transduction pathway; and (4) it modulates a pain signaling mechanism, and, thus, can be used to, for example, (1) modulate the interaction with a non-hVR-1, non-hVR-2, and non-rVR-2 protein molecule; (2) modulate intracellular calcium concentration; (3) activate an hVR-1, hVR-2, and rVR-2-dependent signal transduction pathway; and (4) modulate a pain signaling mechanism.

The isolated nucleic acid molecules of the invention can be used, for example, to express hVR-1, hVR-2, and rVR-2 protein (*e.g.*, via a recombinant expression vector in a host cell in gene therapy applications), to detect hVR-1, hVR-2, and rVR-2 mRNA (*e.g.*, in a biological sample) or a

genetic alteration in an hVR-1, hVR-2, and rVR-2 gene, and to modulate hVR-1, hVR-2, and rVR-2 activity, as described further below. The hVR-1, hVR-2, and rVR-2 proteins can be used to screen for naturally occurring hVR-1, hVR-2, and rVR-2 substrates, to screen for drugs or compounds which modulate hVR-1, hVR-2, and rVR-2 activity, as well as to treat disorders characterized by insufficient or excessive production of hVR-1, hVR-2, and rVR-2 protein or production of hVR-1, hVR-2, and rVR-2 protein forms which have decreased or aberrant activity compared to hVR-1, hVR-2, and rVR-2 wild type protein (*e.g.*, pain disorders). Moreover, the anti-hVR-1, anti-hVR-2, and anti-rVR-2 antibodies of the invention can be used to detect and isolate hVR-1, hVR-2, and rVR-2 proteins, regulate the bioavailability of hVR-1, hVR-2, and rVR-2 proteins, and modulate hVR-1, hVR-2, and rVR-2 activity.

A. Screening Assays:

The invention provides a method (also referred to herein as a "screening assay") for identifying modulators, *i.e.*, candidate or test compounds or agents (*e.g.*, peptides, peptidomimetics, small molecules or other drugs) which bind to hVR-1, hVR-2, and rVR-2 proteins, have a stimulatory or inhibitory effect on, for example, hVR-1, hVR-2, and rVR-2 expression or hVR-1, hVR-2, and rVR-2 activity, or have a stimulatory or inhibitory effect on, for example, the expression or activity of hVR-1, hVR-2, and rVR-2 substrate.

In one embodiment, the invention provides assays for screening candidate or test compounds which are substrates of an hVR-1, hVR-2, and rVR-2 protein or polypeptide or biologically active portion thereof. In another embodiment, the invention provides assays for screening candidate or test compounds which bind to or modulate the activity of an hVR-1, hVR-2, and rVR-2 protein or polypeptide or biologically active portion thereof. The test compounds of the present invention can be obtained using any of the numerous approaches in combinatorial library methods known in the art, including: biological libraries; spatially addressable parallel solid phase or solution phase libraries; synthetic library methods requiring deconvolution; the 'one-bead one-compound' library method; and synthetic library methods using affinity chromatography selection. The biological library approach is limited to peptide libraries, while the other four approaches are applicable to peptide, non-peptide oligomer or small molecule libraries of compounds (Lam, K.S. (1997) *Anticancer Drug Des.* 12:145).

Examples of methods for the synthesis of molecular libraries can be found in the art, for example in: DeWitt *et al.* (1993) *Proc. Natl. Acad. Sci. U.S.A.* 90:6909; Erb *et al.* (1994) *Proc. Natl. Acad. Sci. USA* 91:11422; Zuckermann *et al.* (1994). *J. Med. Chem.* 37:2678; Cho *et al.* (1993) *Science* 261:1303; Carrell *et al.* (1994) *Angew. Chem. Int. Ed. Engl.* 33:2059; Carell *et al.* (1994) *Angew. Chem. Int. Ed. Engl.* 33:2061; and in Gallop *et al.* (1994) *J. Med. Chem.* 37:1233.

Libraries of compounds may be presented in solution (*e.g.*, Houghten (1992) *Biotechniques* 13:412-421), or on beads (Lam (1991) *Nature* 354:82-84), chips (Fodor (1993) *Nature* 364:555-556), bacteria (Ladner USP 5,223,409), spores (Ladner USP '409), plasmids (Cull *et al.* (1992) *Proc Natl Acad Sci USA* 89:1865-1869) or on phage (Scott and Smith (1990) *Science* 249:386-390); (Devlin (1990) *Science* 249:404-406); (Cwirle *et al.* (1990) *Proc. Natl. Acad. Sci.* 87:6378-6382); (Felici (1991) *J. Mol. Biol.* 222:301-310); (Ladner *supra.*).

In one embodiment, an assay is a cell-based assay in which a cell, *e.g.*, a neuronal cell, which expresses an hVR-1, hVR-2, and rVR-2 protein or biologically active portion thereof is contacted with a test compound and the ability of the test compound to modulate hVR-1, hVR-2, and rVR-2 activity is determined. Determining the ability of the test compound to modulate hVR-1, hVR-2, and rVR-2 activity can be accomplished by monitoring, for example, intracellular calcium concentration or membrane depolarization by, *e.g.*, patch-clamp recordings in whole-cell, inside-out, and outside-out configurations (as described in, for example, Tominaga M. *et al.* (1998) *Neuron* 21:531-543). Determining the ability of the test compound to modulate hVR-1, hVR-2, and rVR-2 activity can further be accomplished by monitoring the activity of an hVR-1, hVR-2, and rVR-2-regulated transcription factor. The cell, for example, can be of mammalian origin, *e.g.*, a neuronal cell.

The ability of the test compound to modulate hVR-1, hVR-2, and rVR-2 binding to a substrate or to bind to hVR-1, hVR-2, and rVR-2 can also be determined. Determining the ability of the test compound to modulate hVR-1, hVR-2, and rVR-2 binding to a substrate can be accomplished, for example, by coupling the hVR-1, hVR-2, and rVR-2 substrate with a radioisotope or enzymatic label such that binding of the hVR-1, hVR-2, and rVR-2 substrate to hVR-1, hVR-2, and rVR-2 can be determined by detecting the labeled hVR-1, hVR-2, and rVR-2 substrate in a complex. Determining the ability of the test compound to bind hVR-1, hVR-2, and rVR-2 can be accomplished, for example, by coupling the compound with a radioisotope or enzymatic label such that binding of the compound to hVR-1, hVR-2, and rVR-2 can be determined by detecting the labeled hVR-1, hVR-2, and rVR-2 compound in a complex. For example, compounds (*e.g.*, hVR-1, hVR-2, and rVR-2 substrates) can be labeled with ^{125}I , ^{35}S , ^{14}C , or ^3H , either directly or indirectly, and the radioisotope detected by direct counting of radioemmission or by scintillation counting. Alternatively, compounds can be enzymatically labeled with, for example, horseradish peroxidase, alkaline phosphatase, or luciferase, and the enzymatic label detected by determination of conversion of an appropriate substrate to product.

It is also within the scope of this invention to determine the ability of a compound (*e.g.*, an hVR-1, hVR-2, and rVR-2 substrate) to interact with hVR-1, hVR-2, and rVR-2 without the labeling of any of the interactants. For example, a microphysiometer can be used to detect the interaction of a compound with hVR-1, hVR-2, and rVR-2 without the labeling of either the compound or the hVR-1, hVR-2, and rVR-2. McConnell, H. M. *et al.* (1992) *Science* 257:1906-1912. As used herein, a "microphysiometer" (*e.g.*, Cytosensor) is an analytical instrument that

measures the rate at which a cell acidifies its environment using a light-addressable potentiometric sensor (LAPS). Changes in this acidification rate can be used as an indicator of the interaction between a compound and hVR-1, hVR-2, and rVR-2.

In yet another embodiment, an assay of the present invention is a cell-free assay in which an hVR-1, hVR-2, and rVR-2 protein or biologically active portion thereof is contacted with a test compound and the ability of the test compound to bind to the hVR-1, hVR-2, and rVR-2 protein or biologically active portion thereof is determined. biologically active portions of the hVR-1, hVR-2, and rVR-2 proteins to be used in assays of the present invention include fragments which participate in interactions with non-hVR-1, non-hVR-2, and non-rVR-2 molecules, *e.g.*, fragments with high surface probability scores. Binding of the test compound to the hVR-1, hVR-2, and rVR-2 protein can be determined either directly or indirectly as described above. In a embodiment, the assay includes contacting the hVR-1, hVR-2, and rVR-2 protein or biologically active portion thereof with a known compound which binds hVR-1, hVR-2, and rVR-2 to form an assay mixture, contacting the assay mixture with a test compound, and determining the ability of the test compound to interact with an hVR-1, hVR-2, and rVR-2 protein, wherein determining the ability of the test compound to interact with an hVR-1, hVR-2, and rVR-2 protein comprises determining the ability of the test compound to preferentially bind to hVR-1, hVR-2, and rVR-2 or biologically active portion thereof as compared to the known compound.

In another embodiment, the assay is a cell-free assay in which an hVR-1, hVR-2, and rVR-2 protein or biologically active portion thereof is contacted with a test compound and the ability of the test compound to modulate (*e.g.*, stimulate or inhibit) the activity of the hVR-1, hVR-2, and rVR-2 protein or biologically active portion thereof is determined. Determining the ability of the test compound to modulate the activity of an hVR-1, hVR-2, and rVR-2 protein can be accomplished, for example, by determining the ability of the hVR-1, hVR-2, and rVR-2 protein to bind to an hVR-1, hVR-2, and rVR-2 target molecule, *e.g.*, a vanilloid compound such as capsaicin, by one of the methods described above for determining direct binding. Determining the ability of the hVR-1, hVR-2, and rVR-2 protein to bind to an hVR-1, hVR-2, and rVR-2 target molecule can also be accomplished using a technology such as real-time Biomolecular Interaction Analysis (BIA). Sjolander, S. and Urbaniczky, C. (1991) *Anal. Chem.* 63:2338-2345 and Szabo *et al.* (1995) *Curr. Opin. Struct. Biol.* 5:699-705. As used herein, "BIA" is a technology for studying biospecific interactions in real time, without labeling any of the interactants (*e.g.*, BIAcore). Changes in the optical phenomenon of surface plasmon resonance (SPR) can be used as an indication of real-time reactions between biological molecules.

In an alternative embodiment, determining the ability of the test compound to modulate the activity of an hVR-1, hVR-2, and rVR-2 protein can be accomplished by determining the ability of the hVR-1, hVR-2, and rVR-2 protein to further modulate the activity of a downstream effector of an hVR-1, hVR-2, and rVR-2 target molecule. For example, the activity of the effector molecule

on an appropriate target can be determined or the binding of the effector to an appropriate target can be determined as previously described.

In yet another embodiment, the cell-free assay involves contacting an hVR-1, hVR-2, and rVR-2 protein or biologically active portion thereof with a known compound which binds the hVR-1, hVR-2, and rVR-2 protein to form an assay mixture, contacting the assay mixture with a test compound, and determining the ability of the test compound to interact with the hVR-1, hVR-2, and rVR-2 protein, wherein determining the ability of the test compound to interact with the hVR-1, hVR-2, and rVR-2 protein comprises determining the ability of the hVR-1, hVR-2, and rVR-2 protein to preferentially bind to or modulate the activity of an hVR-1, hVR-2, and rVR-2 target molecule.

The cell-free assays of the present invention are amenable to use of both soluble and/or membrane-bound forms of isolated proteins (*e.g.*, hVR-1, hVR-2, and rVR-2 proteins or biologically active portions thereof). In the case of cell-free assays in which a membrane-bound form of an isolated protein is used it may be desirable to utilize a solubilizing agent such that the membrane-bound form of the isolated protein is maintained in solution. Examples of such solubilizing agents include non-ionic detergents such as n-octylglucoside, n-dodecylglucoside, n-dodecylmaltoside, octanoyl-N-methylglucamide, decanoyl-N-methylglucamide, Triton[®] X-100, Triton[®] X-114, Thesit[®], Isotridecypoly(ethylene glycol ether)_n, 3-[(3-cholamidopropyl)dimethylamminio]-1-propane sulfonate (CHAPS), 3-[(3-cholamidopropyl)dimethylamminio]-2-hydroxy-1-propane sulfonate (CHAPSO), or N-dodecyl=N,N-dimethyl-3-ammonio-1-propane sulfonate.

In more than one embodiment of the above assay methods of the present invention, it may be desirable to immobilize either hVR-1, hVR-2, and rVR-2 or its target molecule to facilitate separation of complexed from uncomplexed forms of one or both of the proteins, as well as to accommodate automation of the assay. Binding of a test compound to an hVR-1, hVR-2, and rVR-2 protein, or interaction of an hVR-1, hVR-2, and rVR-2 protein with a target molecule in the presence and absence of a candidate compound, can be accomplished in any vessel suitable for containing the reactants. Examples of such vessels include microtitre plates, test tubes, and micro-centrifuge tubes. In one embodiment, a fusion protein can be provided which adds a domain that allows one or both of the proteins to be bound to a matrix. For example, glutathione-S-transferase/hVR-1, hVR-2, and rVR-2 fusion proteins or glutathione-S-transferase/target fusion proteins can be adsorbed onto glutathione sepharose beads (Sigma Chemical, St. Louis, MO) or glutathione derivatized microtitre plates, which are then combined with the test compound or the test compound and either the non-adsorbed target protein or hVR-1, hVR-2, and rVR-2 protein, and the mixture incubated under conditions conducive to complex formation (*e.g.*, at physiological conditions for salt and pH). Following incubation, the beads or microtitre plate wells are washed to remove any unbound components, the matrix immobilized in the case of beads, complex determined either directly or indirectly, for example, as described above. Alternatively, the

complexes can be dissociated from the matrix, and the level of hVR-1, hVR-2, and rVR-2 binding or activity determined using standard techniques.

Other techniques for immobilizing proteins on matrices can also be used in the screening assays of the invention. For example, either an hVR-1, hVR-2, and rVR-2 protein or an hVR-1, hVR-2, and rVR-2 target molecule can be immobilized utilizing conjugation of biotin and streptavidin. Biotinylated hVR-1, hVR-2, and rVR-2 protein or target molecules can be prepared from biotin-NHS (N-hydroxy-succinimide) using techniques known in the art (*e.g.*, biotinylation kit, Pierce Chemicals, Rockford, IL), and immobilized in the wells of streptavidin-coated 96 well plates (Pierce Chemical). Alternatively, antibodies reactive with hVR-1, hVR-2, and rVR-2 protein or target molecules but which do not interfere with binding of the hVR-1, hVR-2, and rVR-2 protein to its target molecule can be derivatized to the wells of the plate, and unbound target or hVR-1, hVR-2, and rVR-2 protein trapped in the wells by antibody conjugation. Methods for detecting such complexes, in addition to those described above for the GST-immobilized complexes, include immunodetection of complexes using antibodies reactive with the hVR-1, hVR-2, and rVR-2 protein or target molecule, as well as enzyme-linked assays which rely on detecting an enzymatic activity associated with the hVR-1, hVR-2, and rVR-2 protein or target molecule.

In another embodiment, modulators of hVR-1, hVR-2, and rVR-2 expression are identified in a method wherein a cell is contacted with a candidate compound and the expression of hVR-1, hVR-2, and rVR-2 mRNA or protein in the cell is determined. The level of expression of hVR-1, hVR-2, and rVR-2 mRNA or protein in the presence of the candidate compound is compared to the level of expression of hVR-1, hVR-2, and rVR-2 mRNA or protein in the absence of the candidate compound. The candidate compound can then be identified as a modulator of hVR-1, hVR-2, and rVR-2 expression based on this comparison. For example, when expression of hVR-1, hVR-2, and rVR-2 mRNA or protein is greater (statistically significantly greater) in the presence of the candidate compound than in its absence, the candidate compound is identified as a stimulator of hVR-1, hVR-2, and rVR-2 mRNA or protein expression. Alternatively, when expression of hVR-1, hVR-2, and rVR-2 mRNA or protein is less (statistically significantly less) in the presence of the candidate compound than in its absence, the candidate compound is identified as an inhibitor of hVR-1, hVR-2, and rVR-2 mRNA or protein expression. The level of hVR-1, hVR-2, and rVR-2 mRNA or protein expression in the cells can be determined by methods described herein for detecting hVR-1, hVR-2, and rVR-2 mRNA or protein.

In yet another aspect of the invention, the hVR-1, hVR-2, and rVR-2 proteins can be used as "bait proteins" in a two-hybrid assay or three-hybrid assay (see, *e.g.*, U.S. Patent No. 5,283,317; Zervos *et al.* (1993) *Cell* 72:223-232; Madura *et al.* (1993) *J. Biol. Chem.* 268:12046-12054; Bartel *et al.* (1993) *Biotechniques* 14:920-924; Iwabuchi *et al.* (1993) *Oncogene* 8:1693-1696; and Brent WO94/10300), to identify other proteins, which bind to or interact with hVR-1, hVR-2, and rVR-2 ("hVR-1-binding proteins", "hVR-2-binding proteins", and "rVR-2-binding proteins" or "hVR-1-bp", "hVR-2-bp", and "rVR-2-bp") and are involved in hVR-1, hVR-2, and rVR-2 activity. Such

hVR-1, hVR-2, and rVR-2-binding proteins are also likely to be involved in the propagation of signals by the hVR-1, hVR-2, and rVR-2 proteins or hVR-1, hVR-2, and rVR-2 targets as, for example, downstream elements of an hVR-1, hVR-2, and rVR-2-mediated signaling pathway, *e.g.*, a pain signaling pathway. Alternatively, such hVR-1, hVR-2, and rVR-2-binding proteins are likely to be hVR-1, hVR-2, and rVR-2 inhibitors.

The two-hybrid system is based on the modular nature of most transcription factors, which consist of separable DNA-binding and activation domains. Briefly, the assay utilizes two different DNA constructs. In one construct, the gene that codes for an hVR-1, hVR-2, and rVR-2 protein is fused to a gene encoding the DNA binding domain of a known transcription factor (*e.g.*, GAL-4). In the other construct, a DNA sequence, from a library of DNA sequences, that encodes an unidentified protein ("prey" or "sample") is fused to a gene that codes for the activation domain of the known transcription factor. If the "bait" and the "prey" proteins are able to interact, *in vivo*, forming an hVR-1, hVR-2, and rVR-2-dependent complex, the DNA-binding and activation domains of the transcription factor are brought into close proximity. This proximity allows transcription of a reporter gene (*e.g.*, LacZ) which is operably linked to a transcriptional regulatory site responsive to the transcription factor. Expression of the reporter gene can be detected and cell colonies containing the functional transcription factor can be isolated and used to obtain the cloned gene which encodes the protein which interacts with the hVR-1, hVR-2, and rVR-2 protein.

This invention further pertains to novel agents identified by the above-described screening assays. Accordingly, it is within the scope of this invention to further use an agent identified as described herein in an appropriate animal model. For example, an agent identified as described herein (*e.g.*, an hVR-1, hVR-2, and rVR-2 modulating agent, an antisense hVR-1, hVR-2, and rVR-2 nucleic acid molecule, an hVR-1, hVR-2, and rVR-2-specific antibody, or an hVR-1, hVR-2, and rVR-2-binding partner) can be used in an animal model to determine the efficacy, toxicity, or side effects of treatment with such an agent. Alternatively, an agent identified as described herein can be used in an animal model to determine the mechanism of action of such an agent. Furthermore, this invention pertains to uses of novel agents identified by the above-described screening assays for treatments as described herein.

B. Detection Assays

Portions or fragments of the cDNA sequences identified herein (and the corresponding complete gene sequences) can be used in numerous ways as polynucleotide reagents. For example, these sequences can be used to: (i) map their respective genes on a chromosome; and, thus, locate gene regions associated with genetic disease; (ii) identify an individual from a minute biological sample (tissue typing); and (iii) aid in forensic identification of a biological sample. These applications are described in the subsections below.

1. Chromosome Mapping

Once the sequence (or a portion of the sequence) of a gene has been isolated, this sequence can be used to map the location of the gene on a chromosome. This process is called chromosome mapping. Accordingly, portions or fragments of the hVR-1, hVR-2, and rVR-2 nucleotide sequences, described herein, can be used to map the location of the hVR-1, hVR-2, and rVR-2 genes on a chromosome. The mapping of the hVR-1, hVR-2, and rVR-2 sequences to chromosomes is an important first step in correlating these sequences with genes associated with disease.

Briefly, hVR-1, hVR-2, and rVR-2 genes can be mapped to chromosomes by preparing PCR primers (preferably 15-25 bp in length) from the hVR-1, hVR-2, and rVR-2 nucleotide sequences. Computer analysis of the hVR-1, hVR-2, and rVR-2 sequences can be used to predict primers that do not span more than one exon in the genomic DNA, thus complicating the amplification process. These primers can then be used for PCR screening of somatic cell hybrids containing individual human chromosomes. Only those hybrids containing the human gene corresponding to the hVR-1, hVR-2, and rVR-2 sequences will yield an amplified fragment.

Somatic cell hybrids are prepared by fusing somatic cells from different mammals (*e.g.*, human and mouse cells). As hybrids of human and mouse cells grow and divide, they gradually lose human chromosomes in random order, but retain the mouse chromosomes. By using media in which mouse cells cannot grow, because they lack a particular enzyme, but human cells can, the one human chromosome that contains the gene encoding the needed enzyme, will be retained. By using various media, panels of hybrid cell lines can be established. Each cell line in a panel contains either a single human chromosome or a small number of human chromosomes, and a full set of mouse chromosomes, allowing easy mapping of individual genes to specific human chromosomes. (D'Eustachio P. *et al.* (1983) *Science* 220:919-924). Somatic cell hybrids containing only fragments of human chromosomes can also be produced by using human chromosomes with translocations and deletions.

PCR mapping of somatic cell hybrids is a rapid procedure for assigning a particular sequence to a particular chromosome. Three or more sequences can be assigned per day using a single thermal cycler. Using the hVR-1, hVR-2, and rVR-2 nucleotide sequences to design oligonucleotide primers, sublocalization can be achieved with panels of fragments from specific chromosomes. Other mapping strategies which can similarly be used to map an hVR-1, hVR-2, and rVR-2 sequence to its chromosome include *in situ* hybridization (described in Fan, Y. *et al.* (1990) *Proc. Natl. Acad. Sci. USA*, 87:6223-27), pre-screening with labeled flow-sorted chromosomes, and pre-selection by hybridization to chromosome specific cDNA libraries.

Fluorescence *in situ* hybridization (FISH) of a DNA sequence to a metaphase chromosomal spread can further be used to provide a precise chromosomal location in one step. Chromosome spreads can be made using cells whose division has been blocked in metaphase by a chemical such as colcemid that disrupts the mitotic spindle. The chromosomes can be treated briefly with trypsin,

and then stained with Giemsa. A pattern of light and dark bands develops on each chromosome, so that the chromosomes can be identified individually. The FISH technique can be used with a DNA sequence as short as 500 or 600 bases. However, clones larger than 1,000 bases have a higher likelihood of binding to a unique chromosomal location with sufficient signal intensity for simple detection. Preferably 1,000 bases, and more preferably 2,000 bases will suffice to get good results at a reasonable amount of time. For a review of this technique, see Verma *et al.*, Human Chromosomes: A Manual of Basic Techniques (Pergamon Press, New York 1988).

Reagents for chromosome mapping can be used individually to mark a single chromosome or a single site on that chromosome, or panels of reagents can be used for marking multiple sites and/or multiple chromosomes. Reagents corresponding to noncoding regions of the genes actually are for mapping purposes. Coding sequences are more likely to be conserved within gene families, thus increasing the chance of cross hybridizations during chromosomal mapping.

Once a sequence has been mapped to a precise chromosomal location, the physical position of the sequence on the chromosome can be correlated with genetic map data. (Such data are found, for example, in V. McKusick, Mendelian Inheritance in Man, available on-line through Johns Hopkins University Welch Medical Library). The relationship between a gene and a disease, mapped to the same chromosomal region, can then be identified through linkage analysis (co-inheritance of physically adjacent genes), described in, for example, Egeland, J. *et al.* (1987) *Nature*, 325:783-787.

Moreover, differences in the DNA sequences between individuals affected and unaffected with a disease associated with the hVR-1, hVR-2, and rVR-2 gene, can be determined. If a mutation is observed in some or all of the affected individuals but not in any unaffected individuals, then the mutation is likely to be the causative agent of the particular disease. Comparison of affected and unaffected individuals generally involves first looking for structural alterations in the chromosomes, such as deletions or translocations that are visible from chromosome spreads or detectable using PCR based on that DNA sequence. Ultimately, complete sequencing of genes from several individuals can be performed to confirm the presence of a mutation and to distinguish mutations from polymorphisms.

2. Tissue Typing

The hVR-1, hVR-2, and rVR-2 sequences of the present invention can also be used to identify individuals from minute biological samples. The United States military, for example, is considering the use of restriction fragment length polymorphism (RFLP) for identification of its personnel. In this technique, an individual's genomic DNA is digested with one or more restriction enzymes, and probed on a Southern blot to yield unique bands for identification. This method does not suffer from the current limitations of "Dog Tags" which can be lost, switched, or stolen, making positive identification difficult. The sequences of the present invention are useful as additional DNA markers for RFLP (described in U.S. Patent 5,272,057).

Furthermore, the sequences of the present invention can be used to provide an alternative technique which determines the actual base-by-base DNA sequence of selected portions of an individual's genome. Thus, the hVR-1, hVR-2, and rVR-2 nucleotide sequences described herein can be used to prepare two PCR primers from the 5' and 3' ends of the sequences. These primers can then be used to amplify an individual's DNA and subsequently sequence it.

Panels of corresponding DNA sequences from individuals, prepared in this manner, can provide unique individual identifications, as each individual will have a unique set of such DNA sequences due to allelic differences. The sequences of the present invention can be used to obtain such identification sequences from individuals and from tissue. The hVR-1, hVR-2, and rVR-2 nucleotide sequences of the invention uniquely represent portions of the human genome. Allelic variation occurs to some degree in the coding regions of these sequences, and to a greater degree in the noncoding regions. It is estimated that allelic variation between individual humans occurs with a frequency of about once per each 500 bases. Each of the sequences described herein can, to some degree, be used as a standard against which DNA from an individual can be compared for identification purposes. Because greater numbers of polymorphisms occur in the noncoding regions, fewer sequences are necessary to differentiate individuals.

If a panel of reagents from hVR-1, hVR-2, and rVR-2 nucleotide sequences described herein is used to generate a unique identification database for an individual, those same reagents can later be used to identify tissue from that individual. Using the unique identification database, positive identification of the individual, living or dead, can be made from extremely small tissue samples.

3. Use of Partial hVR-1, hVR-2, and rVR-2 Sequences in Forensic Biology

DNA-based identification techniques can also be used in forensic biology. Forensic biology is a scientific field employing genetic typing of biological evidence found at a crime scene as a means for positively identifying, for example, a perpetrator of a crime. To make such an identification, PCR technology can be used to amplify DNA sequences taken from very small biological samples such as tissues, *e.g.*, hair or skin, or body fluids, *e.g.*, blood, saliva, or semen found at a crime scene. The amplified sequence can then be compared to a standard, thereby allowing identification of the origin of the biological sample.

The sequences of the present invention can be used to provide polynucleotide reagents, *e.g.*, PCR primers, targeted to specific loci in the human genome, which can enhance the reliability of DNA-based forensic identifications by, for example, providing another "identification marker" (*i.e.* another DNA sequence that is unique to a particular individual). As mentioned above, actual base sequence information can be used for identification as an accurate alternative to patterns formed by restriction enzyme generated fragments. Examples of polynucleotide reagents include the hVR-1, hVR-2, and rVR-2 nucleotide sequences or portions thereof, *e.g.*, fragments derived from SEQ ID NO:1, 3, 4, 6, 7, 9, 10, or 11 having a length of at least 20 bases, preferably at least 30 bases.

The hVR-1, hVR-2, and rVR-2 nucleotide sequences described herein can further be used to provide polynucleotide reagents, *e.g.*, labeled or labelable probes which can be used in, for example, an *in situ* hybridization technique, to identify a specific tissue, *e.g.*, brain tissue. This can be very useful in cases where a forensic pathologist is presented with a tissue of unknown origin.

5 Panels of such hVR-1, hVR-2, and rVR-2 probes can be used to identify tissue by species and/or by organ type.

In a similar fashion, these reagents, *e.g.*, hVR-1, hVR-2, and rVR-2 primers or probes can be used to screen tissue culture for contamination (*i.e.* screen for the presence of a mixture of different types of cells in a culture).

10 C. Predictive Medicine:

The present invention also pertains to the field of predictive medicine in which diagnostic assays, prognostic assays, and monitoring clinical trials are used for prognostic (predictive) purposes to thereby treat an individual prophylactically. Accordingly, one aspect of the present invention relates to diagnostic assays for determining hVR-1, hVR-2, and rVR-2 protein and/or nucleic acid expression as well as hVR-1, hVR-2, and rVR-2 activity, in the context of a biological sample (*e.g.*, blood, serum, cells, tissue) to thereby determine whether an individual is afflicted with a disease or disorder, or is at risk of developing a disorder, associated with aberrant hVR-1, hVR-2, and rVR-2 expression or activity. The invention also provides for prognostic (or
15
20
25 predictive) assays for determining whether an individual is at risk of developing a disorder associated with hVR-1, hVR-2, and rVR-2 protein, nucleic acid expression or activity. For example, mutations in an hVR-1, hVR-2, and rVR-2 gene can be assayed in a biological sample. Such assays can be used for prognostic or predictive purpose to thereby prophylactically treat an individual prior to the onset of a disorder characterized by or associated with hVR-1, hVR-2, and rVR-2 protein, nucleic acid expression or activity.

Another aspect of the invention pertains to monitoring the influence of agents (*e.g.*, drugs, compounds) on the expression or activity of hVR-1, hVR-2, and rVR-2 in clinical trials.

These and other agents are described in further detail in the following sections.

30 1. Diagnostic Assays

An exemplary method for detecting the presence or absence of hVR-1, hVR-2, and rVR-2 protein or nucleic acid in a biological sample involves obtaining a biological sample from a test subject and contacting the biological sample with a compound or an agent capable of detecting hVR-1, hVR-2, and rVR-2 protein or nucleic acid (*e.g.*, mRNA, genomic DNA) that encodes hVR-1, hVR-2, and rVR-2 protein such that the presence of hVR-1, hVR-2, and rVR-2 protein or nucleic acid is detected in the biological sample. A agent for detecting hVR-1, hVR-2, and rVR-2 mRNA or genomic DNA is a labeled nucleic acid probe capable of hybridizing to hVR-1, hVR-2, and rVR-2 mRNA or genomic DNA. The nucleic acid probe can be, for example, a full-length hVR-1, hVR-
35

2, and rVR-2 nucleic acid, such as the nucleic acid of SEQ ID NO:1, 3, 4, 6, 7, 9, 10, or 12, or a portion thereof, such as an oligonucleotide of at least 15, 30, 50, 100, 250 or 500 nucleotides in length and sufficient to specifically hybridize under stringent conditions to hVR-1, hVR-2, and rVR-2 mRNA or genomic DNA. Other suitable probes for use in the diagnostic assays of the invention are described herein.

An agent for detecting hVR-1, hVR-2, and rVR-2 protein is an antibody capable of binding to hVR-1, hVR-2, and rVR-2 protein, preferably an antibody with a detectable label. Antibodies can be polyclonal, or more preferably, monoclonal. An intact antibody, or a fragment thereof (e.g., Fab or F(ab')₂) can be used. The term "labeled", with regard to the probe or antibody, is intended to encompass direct labeling of the probe or antibody by coupling (i.e., physically linking) a detectable substance to the probe or antibody, as well as indirect labeling of the probe or antibody by reactivity with another reagent that is directly labeled. Examples of indirect labeling include detection of a primary antibody using a fluorescently labeled secondary antibody and end-labeling of a DNA probe with biotin such that it can be detected with fluorescently labeled streptavidin. The term "biological sample" is intended to include tissues, cells and biological fluids isolated from a subject, as well as tissues, cells and fluids present within a subject. That is, the detection method of the invention can be used to detect hVR-1, hVR-2, and rVR-2 mRNA, protein, or genomic DNA in a biological sample *in vitro* as well as *in vivo*. For example, *in vitro* techniques for detection of hVR-1, hVR-2, and rVR-2 mRNA include Northern hybridizations and *in situ* hybridizations. *In vitro* techniques for detection of hVR-1, hVR-2, and rVR-2 protein include enzyme linked immunosorbent assays (ELISAs), Western blots, immunoprecipitations and immunofluorescence. *In vitro* techniques for detection of hVR-1, hVR-2, and rVR-2 genomic DNA include Southern hybridizations. Furthermore, *in vivo* techniques for detection of hVR-1, hVR-2, and rVR-2 protein include introducing into a subject a labeled anti-hVR-1, hVR-2, and rVR-2 antibody. For example, the antibody can be labeled with a radioactive marker whose presence and location in a subject can be detected by standard imaging techniques.

In one embodiment, the biological sample contains protein molecules from the test subject. Alternatively, the biological sample can contain mRNA molecules from the test subject or genomic DNA molecules from the test subject. A biological sample is a serum sample isolated by conventional means from a subject.

In another embodiment, the methods further involve obtaining a control biological sample from a control subject, contacting the control sample with a compound or agent capable of detecting hVR-1, hVR-2, and rVR-2 protein, mRNA, or genomic DNA, such that the presence of hVR-1, hVR-2, and rVR-2 protein, mRNA or genomic DNA is detected in the biological sample, and comparing the presence of hVR-1, hVR-2, and rVR-2 protein, mRNA or genomic DNA in the control sample with the presence of hVR-1, hVR-2, and rVR-2 protein, mRNA or genomic DNA in the test sample.

The invention also encompasses kits for detecting the presence of hVR-1, hVR-2, and rVR-2 in a biological sample. For example, the kit can comprise a labeled compound or agent capable of detecting hVR-1, hVR-2, and rVR-2 protein or mRNA in a biological sample; means for determining the amount of hVR-1, hVR-2, and rVR-2 in the sample; and means for comparing the amount of hVR-1, hVR-2, and rVR-2 in the sample with a standard. The compound or agent can be packaged in a suitable container. The kit can further comprise instructions for using the kit to detect hVR-1, hVR-2, and rVR-2 protein or nucleic acid.

2. Prognostic Assays

The diagnostic methods described herein can furthermore be utilized to identify subjects having or at risk of developing a disease or disorder associated with aberrant hVR-1, hVR-2, and rVR-2 expression or activity. As used herein, the term "aberrant" includes an hVR-1, hVR-2, and rVR-2 expression or activity which deviates from the wild type hVR-1, hVR-2, and rVR-2 expression or activity. Aberrant expression or activity includes increased or decreased expression or activity, as well as expression or activity which does not follow the wild type developmental pattern of expression or the subcellular pattern of expression. For example, aberrant hVR-1, hVR-2, and rVR-2 expression or activity is intended to include the cases in which a mutation in the hVR-1, hVR-2, and rVR-2 gene causes the hVR-1, hVR-2, and rVR-2 gene to be under-expressed or over-expressed and situations in which such mutations result in a non-functional hVR-1, hVR-2, and rVR-2 protein or a protein which does not function in a wild-type fashion, *e.g.*, a protein which does not interact with an hVR-1, hVR-2, and rVR-2 ligand or one which interacts with a non-hVR-1, non-hVR-2, and non-rVR-2 ligand.

The assays described herein, such as the preceding diagnostic assays or the following assays, can be utilized to identify a subject having or at risk of developing a disorder associated with a misregulation in hVR-1, hVR-2, and rVR-2 protein activity or nucleic acid expression, such as a pain disorder. Alternatively, the prognostic assays can be utilized to identify a subject having or at risk for developing a disorder associated with a misregulation in hVR-1, hVR-2, and rVR-2 protein activity or nucleic acid expression, such as a pain disorder. Thus, the present invention provides a method for identifying a disease or disorder associated with aberrant hVR-1, hVR-2, and rVR-2 expression or activity in which a test sample is obtained from a subject and hVR-1, hVR-2, and rVR-2 protein or nucleic acid (*e.g.*, mRNA or genomic DNA) is detected, wherein the presence of hVR-1, hVR-2, and rVR-2 protein or nucleic acid is diagnostic for a subject having or at risk of developing a disease or disorder associated with aberrant hVR-1, hVR-2, and rVR-2 expression or activity. As used herein, a "test sample" refers to a biological sample obtained from a subject of interest. For example, a test sample can be a biological fluid (*e.g.*, serum), cell sample, or tissue.

Furthermore, the prognostic assays described herein can be used to determine whether a subject can be administered an agent (*e.g.*, an agonist, antagonist, peptidomimetic, protein, peptide, nucleic acid, small molecule, or other drug candidate) to treat a disease or disorder associated with

aberrant hVR-1, hVR-2, and rVR-2 expression or activity. For example, such methods can be used to determine whether a subject can be effectively treated with an agent for a pain disorder. Thus, the present invention provides methods for determining whether a subject can be effectively treated with an agent for a disorder associated with aberrant hVR-1, hVR-2, and rVR-2 expression or activity in which a test sample is obtained and hVR-1, hVR-2, and rVR-2 protein or nucleic acid expression or activity is detected (e.g., wherein the abundance of hVR-1, hVR-2, and rVR-2 protein or nucleic acid expression or activity is diagnostic for a subject that can be administered the agent to treat a disorder associated with aberrant hVR-1, hVR-2, and rVR-2 expression or activity).

The methods of the invention can also be used to detect genetic alterations in an hVR-1, hVR-2, and rVR-2 gene, thereby determining if a subject with the altered gene is at risk for a disorder characterized by misregulation in hVR-1, hVR-2, and rVR-2 protein activity or nucleic acid expression, such as a neurodegenerative disorder. In embodiments, the methods include detecting, in a sample of cells from the subject, the presence or absence of a genetic alteration characterized by at least one of an alteration affecting the integrity of a gene encoding an hVR-1, hVR-2, and rVR-2-protein, or the mis-expression of the hVR-1, hVR-2, and rVR-2 gene. For example, such genetic alterations can be detected by ascertaining the existence of at least one of 1) a deletion of one or more nucleotides from an hVR-1, hVR-2, and rVR-2 gene; 2) an addition of one or more nucleotides to an hVR-1, hVR-2, and rVR-2 gene; 3) a substitution of one or more nucleotides of an hVR-1, hVR-2, and rVR-2 gene, 4) a chromosomal rearrangement of an hVR-1, hVR-2, and rVR-2 gene; 5) an alteration in the level of a messenger RNA transcript of an hVR-1, hVR-2, and rVR-2 gene, 6) aberrant modification of an hVR-1, hVR-2, and rVR-2 gene, such as of the methylation pattern of the genomic DNA, 7) the presence of a non-wild type splicing pattern of a messenger RNA transcript of an hVR-1, hVR-2, and rVR-2 gene, 8) a non-wild type level of an hVR-1, hVR-2, and rVR-2-protein, 9) allelic loss of an hVR-1, hVR-2, and rVR-2 gene, and 10) inappropriate post-translational modification of an hVR-1, hVR-2, and rVR-2-protein. As described herein, there are a large number of assays known in the art which can be used for detecting alterations in an hVR-1, hVR-2, and rVR-2 gene. A biological sample is a tissue or serum sample isolated by conventional means from a subject.

In certain embodiments, detection of the alteration involves the use of a probe/primer in a polymerase chain reaction (PCR) (see, e.g., U.S. Patent Nos. 4,683,195 and 4,683,202), such as anchor PCR or RACE PCR, or, alternatively, in a ligation chain reaction (LCR) (see, e.g., Landegran *et al.* (1988) *Science* 241:1077-1080; and Nakazawa *et al.* (1994) *Proc. Natl. Acad. Sci. USA* 91:360-364), the latter of which can be particularly useful for detecting point mutations in the hVR-1, hVR-2, and rVR-2-gene (see Abravaya *et al.* (1995) *Nucleic Acids Res.* 23:675-682). This method can include the steps of collecting a sample of cells from a subject, isolating nucleic acid (e.g., genomic, mRNA or both) from the cells of the sample, contacting the nucleic acid sample with one or more primers which specifically hybridize to an hVR-1, hVR-2, and rVR-2 gene under conditions such that hybridization and amplification of the hVR-1, hVR-2, and rVR-2-gene (if

present) occurs, and detecting the presence or absence of an amplification product, or detecting the size of the amplification product and comparing the length to a control sample. It is anticipated that PCR and/or LCR may be desirable to use as a preliminary amplification step in conjunction with any of the techniques used for detecting mutations described herein.

Alternative amplification methods include: self sustained sequence replication (Guatelli, J.C. *et al.*, (1990) *Proc. Natl. Acad. Sci. USA* 87:1874-1878), transcriptional amplification system (Kwoh, D.Y. *et al.*, (1989) *Proc. Natl. Acad. Sci. USA* 86:1173-1177), Q-Beta Replicase (Lizardi, P.M. *et al.* (1988) *Bio-Technology* 6:1197), or any other nucleic acid amplification method, followed by the detection of the amplified molecules using techniques well known to those of skill in the art. These detection schemes are especially useful for the detection of nucleic acid molecules if such molecules are present in very low numbers.

In an alternative embodiment, mutations in an hVR-1, hVR-2, and rVR-2 gene from a sample cell can be identified by alterations in restriction enzyme cleavage patterns. For example, sample and control DNA is isolated, amplified (optionally), digested with one or more restriction endonucleases, and fragment length sizes are determined by gel electrophoresis and compared. Differences in fragment length sizes between sample and control DNA indicates mutations in the sample DNA. Moreover, the use of sequence specific ribozymes (see, for example, U.S. Patent No. 5,498,531) can be used to score for the presence of specific mutations by development or loss of a ribozyme cleavage site.

In other embodiments, genetic mutations in hVR-1, hVR-2, and rVR-2 can be identified by hybridizing a sample and control nucleic acids, *e.g.*, DNA or RNA, to high density arrays containing hundreds or thousands of oligonucleotides probes (Cronin, M.T. *et al.* (1996) *Human Mutation* 7: 244-255; Kozal, M.J. *et al.* (1996) *Nature Medicine* 2: 753-759). For example, genetic mutations in hVR-1, hVR-2, and rVR-2 can be identified in two dimensional arrays containing light-generated DNA probes as described in Cronin, M.T. *et al. supra*. Briefly, a first hybridization array of probes can be used to scan through long stretches of DNA in a sample and control to identify base changes between the sequences by making linear arrays of sequential overlapping probes. This step allows the identification of point mutations. This step is followed by a second hybridization array that allows the characterization of specific mutations by using smaller, specialized probe arrays complementary to all variants or mutations detected. Each mutation array is composed of parallel probe sets, one complementary to the wild-type gene and the other complementary to the mutant gene.

In yet another embodiment, any of a variety of sequencing reactions known in the art can be used to directly sequence the hVR-1, hVR-2, and rVR-2 gene and detect mutations by comparing the sequence of the sample hVR-1, hVR-2, and rVR-2 with the corresponding wild-type (control) sequence. Examples of sequencing reactions include those based on techniques developed by Maxam and Gilbert ((1977) *Proc. Natl. Acad. Sci. USA* 74:560) or Sanger ((1977) *Proc. Natl. Acad. Sci. USA* 74:5463). It is also contemplated that any of a variety of automated sequencing

procedures can be utilized when performing the diagnostic assays ((1995) *Biotechniques* 19:448), including sequencing by mass spectrometry (see, e.g., PCT International Publication No. WO 94/16101; Cohen *et al.* (1996) *Adv. Chromatogr.* 36:127-162; and Griffin *et al.* (1993) *Appl. Biochem. Biotechnol.* 38:147-159).

5 Other methods for detecting mutations in the hVR-1, hVR-2, and rVR-2 gene include methods in which protection from cleavage agents is used to detect mismatched bases in RNA/RNA or RNA/DNA heteroduplexes (Myers *et al.* (1985) *Science* 230:1242). In general, the art technique of "mismatch cleavage" starts by providing heteroduplexes of formed by hybridizing (labeled) RNA or DNA containing the wild-type hVR-1, hVR-2, and rVR-2 sequence with
10 potentially mutant RNA or DNA obtained from a tissue sample. The double-stranded duplexes are treated with an agent which cleaves single-stranded regions of the duplex such as which will exist due to basepair mismatches between the control and sample strands. For instance, RNA/DNA duplexes can be treated with RNase and DNA/DNA hybrids treated with S1 nuclease to enzymatically digesting the mismatched regions. In other embodiments, either DNA/DNA or
15 RNA/DNA duplexes can be treated with hydroxylamine or osmium tetroxide and with piperidine in order to digest mismatched regions. After digestion of the mismatched regions, the resulting material is then separated by size on denaturing polyacrylamide gels to determine the site of mutation. See, for example, Cotton *et al.* (1988) *Proc. Natl Acad Sci USA* 85:4397; Saleeba *et al.* (1992) *Methods Enzymol.* 217:286-295. In a embodiment, the control DNA or RNA can be labeled for detection.
20

In still another embodiment, the mismatch cleavage reaction employs one or more proteins that recognize mismatched base pairs in double-stranded DNA (so called "DNA mismatch repair" enzymes) in defined systems for detecting and mapping point mutations in hVR-1, hVR-2, and rVR-2 cDNAs obtained from samples of cells. For example, the mutY enzyme of *E. coli* cleaves A at G/A mismatches and the thymidine DNA glycosylase from HeLa cells cleaves T at G/T mismatches (Hsu *et al.* (1994) *Carcinogenesis* 15:1657-1662). According to an exemplary embodiment, a probe based on an hVR-1, hVR-2, and rVR-2 sequence, e.g., a wild-type hVR-1, hVR-2, and rVR-2 sequence, is hybridized to a cDNA or other DNA product from a test cell(s). The duplex is treated with a DNA mismatch repair enzyme, and the cleavage products, if any, can
25 be detected from electrophoresis protocols or the like. See, for example, U.S. Patent No. 5,459,039.
30

In other embodiments, alterations in electrophoretic mobility will be used to identify mutations in hVR-1, hVR-2, and rVR-2 genes. For example, single strand conformation polymorphism (SSCP) may be used to detect differences in electrophoretic mobility between
35 mutant and wild type nucleic acids (orita *et al.* (1989) *Proc Natl. Acad. Sci USA*: 86:2766, see also Cotton (1993) *Mutat. Res.* 285:125-144; and Hayashi (1992) *Genet. Anal. Tech. Appl.* 9:73-79). Single-stranded DNA fragments of sample and control hVR-1, hVR-2, and rVR-2 nucleic acids will be denatured and allowed to renature. The secondary structure of single-stranded nucleic acids

varies according to sequence, the resulting alteration in electrophoretic mobility enables the detection of even a single base change. The DNA fragments may be labeled or detected with labeled probes. The sensitivity of the assay may be enhanced by using RNA (rather than DNA), in which the secondary structure is more sensitive to a change in sequence. In a embodiment, the subject method utilizes heteroduplex analysis to separate double stranded heteroduplex molecules on the basis of changes in electrophoretic mobility (Keen *et al.* (1991) *Trends Genet* 7:5).

In yet another embodiment the movement of mutant or wild-type fragments in polyacrylamide gels containing a gradient of denaturant is assayed using denaturing gradient gel electrophoresis (DGGE) (Myers *et al.* (1985) *Nature* 313:495). When DGGE is used as the method of analysis, DNA will be modified to insure that it does not completely denature, for example by adding a GC clamp of approximately 40 bp of high-melting GC-rich DNA by PCR. In a further embodiment, a temperature gradient is used in place of a denaturing gradient to identify differences in the mobility of control and sample DNA (Rosenbaum and Reissner (1987) *Biophys Chem* 265:12753).

Examples of other techniques for detecting point mutations include, but are not limited to, selective oligonucleotide hybridization, selective amplification, or selective primer extension. For example, oligonucleotide primers may be prepared in which the known mutation is placed centrally and then hybridized to target DNA under conditions which permit hybridization only if a perfect match is found (Saiki *et al.* (1986) *Nature* 324:163); Saiki *et al.* (1989) *Proc. Natl Acad. Sci USA* 86:6230). Such allele specific oligonucleotides are hybridized to PCR amplified target DNA or a number of different mutations when the oligonucleotides are attached to the hybridizing membrane and hybridized with labeled target DNA.

Alternatively, allele specific amplification technology which depends on selective PCR amplification may be used in conjunction with the instant invention. Oligonucleotides used as primers for specific amplification may carry the mutation of interest in the center of the molecule (so that amplification depends on differential hybridization) (Gibbs *et al.* (1989) *Nucleic Acids Res.* 17:2437-2448) or at the extreme 3' end of one primer where, under appropriate conditions, mismatch can prevent, or reduce polymerase extension (Prossner (1993) *Tibtech* 11:238). In addition it may be desirable to introduce a novel restriction site in the region of the mutation to create cleavage-based detection (Gasparini *et al.* (1992) *Mol. Cell Probes* 6:1). It is anticipated that in certain embodiments amplification may also be performed using Taq ligase for amplification (Barany (1991) *Proc. Natl. Acad. Sci USA* 88:189). In such cases, ligation will occur only if there is a perfect match at the 3' end of the 5' sequence making it possible to detect the presence of a known mutation at a specific site by looking for the presence or absence of amplification.

The methods described herein may be performed, for example, by utilizing pre-packaged diagnostic kits comprising at least one probe nucleic acid or antibody reagent described herein, which may be conveniently used, *e.g.*, in clinical settings to diagnose patients exhibiting symptoms or family history of a disease or illness involving an hVR-1, hVR-2, and rVR-2 gene.

Furthermore, any cell type or tissue in which hVR-1, hVR-2, and rVR-2 is expressed may be utilized in the prognostic assays described herein.

3. Monitoring of Effects During Clinical Trials

Monitoring the influence of agents (*e.g.*, drugs) on the expression or activity of an hVR-1, hVR-2, and rVR-2 protein can be applied not only in basic drug screening, but also in clinical trials. For example, the effectiveness of an agent determined by a screening assay as described herein to increase hVR-1, hVR-2, and rVR-2 gene expression, protein levels, or upregulate hVR-1, hVR-2, and rVR-2 activity, can be monitored in clinical trials of subjects exhibiting decreased hVR-1, hVR-2, and rVR-2 gene expression, protein levels, or downregulated hVR-1, hVR-2, and rVR-2 activity. Alternatively, the effectiveness of an agent determined by a screening assay to decrease hVR-1, hVR-2, and rVR-2 gene expression, protein levels, or downregulate hVR-1, hVR-2, and rVR-2 activity, can be monitored in clinical trials of subjects exhibiting increased hVR-1, hVR-2, and rVR-2 gene expression, protein levels, or upregulated hVR-1, hVR-2, and rVR-2 activity. In such clinical trials, the expression or activity of an hVR-1, hVR-2, and rVR-2 gene, and preferably, other genes that have been implicated in, for example, an hVR-1, hVR-2, and rVR-2-associated disorder can be used as a "read out" or markers of the phenotype of a particular cell.

For example, and not by way of limitation, genes, including hVR-1, hVR-2, and rVR-2, that are modulated in cells by treatment with an agent (*e.g.*, compound, drug or small molecule) which modulates hVR-1, hVR-2, and rVR-2 activity (*e.g.*, identified in a screening assay as described herein) can be identified. Thus, to study the effect of agents on hVR-1, hVR-2, and rVR-2-associated disorders (*e.g.*, pain disorders), for example, in a clinical trial, cells can be isolated and RNA prepared and analyzed for the levels of expression of hVR-1, hVR-2, and rVR-2 and other genes implicated in the hVR-1, hVR-2, and rVR-2-associated disorder, respectively. The levels of gene expression (*e.g.*, a gene expression pattern) can be quantified by northern blot analysis or RT-PCR, as described herein, or alternatively by measuring the amount of protein produced, by one of the methods as described herein, or by measuring the levels of activity of hVR-1, hVR-2, and rVR-2 or other genes. In this way, the gene expression pattern can serve as a marker, indicative of the physiological response of the cells to the agent. Accordingly, this response state may be determined before, and at various points during treatment of the individual with the agent.

In a embodiment, the present invention provides a method for monitoring the effectiveness of treatment of a subject with an agent (*e.g.*, an agonist, antagonist, peptidomimetic, protein, peptide, nucleic acid, small molecule, or other drug candidate identified by the screening assays described herein) including the steps of (i) obtaining a pre-administration sample from a subject prior to administration of the agent; (ii) detecting the level of expression of an hVR-1, hVR-2, and rVR-2 protein, mRNA, or genomic DNA in the preadministration sample; (iii) obtaining one or more post-administration samples from the subject; (iv) detecting the level of expression or activity of the hVR-1, hVR-2, and rVR-2 protein, mRNA, or genomic DNA in the post-administration

samples; (v) comparing the level of expression or activity of the hVR-1, hVR-2, and rVR-2 protein, mRNA, or genomic DNA in the pre-administration sample with the hVR-1, hVR-2, and rVR-2 protein, mRNA, or genomic DNA in the post administration sample or samples; and (vi) altering the administration of the agent to the subject accordingly. For example, increased administration of the agent may be desirable to increase the expression or activity of hVR-1, hVR-2, and rVR-2 to higher levels than detected, *i.e.*, to increase the effectiveness of the agent. Alternatively, decreased administration of the agent may be desirable to decrease expression or activity of hVR-1, hVR-2, and rVR-2 to lower levels than detected, *i.e.* to decrease the effectiveness of the agent. According to such an embodiment, hVR-1, hVR-2, and rVR-2 expression or activity may be used as an indicator of the effectiveness of an agent, even in the absence of an observable phenotypic response.

D. Methods of Treatment:

The present invention provides for both prophylactic and therapeutic methods of treating a subject at risk of (or susceptible to) a disorder or having a disorder associated with aberrant hVR-1, hVR-2, and rVR-2 expression or activity. With regards to both prophylactic and therapeutic methods of treatment, such treatments may be specifically tailored or modified, based on knowledge obtained from the field of pharmacogenomics. "Pharmacogenomics", as used herein, refers to the application of genomics technologies such as gene sequencing, statistical genetics, and gene expression analysis to drugs in clinical development and on the market. More specifically, the term refers the study of how a patient's genes determine his or her response to a drug (*e.g.*, a patient's "drug response phenotype", or "drug response genotype".) Thus, another aspect of the invention provides methods for tailoring an individual's prophylactic or therapeutic treatment with either the hVR-1, hVR-2, and rVR-2 molecules of the present invention or hVR-1, hVR-2, and rVR-2 modulators according to that individual's drug response genotype. Pharmacogenomics allows a clinician or physician to target prophylactic or therapeutic treatments to patients who will most benefit from the treatment and to avoid treatment of patients who will experience toxic drug-related side effects.

1. Prophylactic Methods

In one aspect, the invention provides a method for preventing in a subject, a disease or condition associated with an aberrant hVR-1, hVR-2, and rVR-2 expression or activity, by administering to the subject an hVR-1, hVR-2, and rVR-2 or an agent which modulates hVR-1, hVR-2, and rVR-2 expression or at least one hVR-1, hVR-2, and rVR-2 activity. Subjects at risk for a disease which is caused or contributed to by aberrant hVR-1, hVR-2, and rVR-2 expression or activity can be identified by, for example, any or a combination of diagnostic or prognostic assays

as described herein. Administration of a prophylactic agent can occur prior to the manifestation of symptoms characteristic of the hVR-1, hVR-2, and rVR-2 aberrancy, such that a disease or disorder is prevented or, alternatively, delayed in its progression. Depending on the type of hVR-1, hVR-2, and rVR-2 aberrancy, for example, an hVR-1, hVR-2, and rVR-2, hVR-1, hVR-2, and rVR-2 agonist or hVR-1, hVR-2, and rVR-2 antagonist agent can be used for treating the subject. The appropriate agent can be determined based on screening assays described herein.

2. Therapeutic Methods

Another aspect of the invention pertains to methods of modulating hVR-1, hVR-2, and rVR-2 expression or activity for therapeutic purposes. Accordingly, in an exemplary embodiment, the modulatory method of the invention involves contacting a cell with an hVR-1, hVR-2, and rVR-2 or agent that modulates one or more of the activities of hVR-1, hVR-2, and rVR-2 protein activity associated with the cell. An agent that modulates hVR-1, hVR-2, and rVR-2 protein activity can be an agent as described herein, such as a nucleic acid or a protein, a naturally-occurring target molecule of an hVR-1, hVR-2, and rVR-2 protein (e.g., an hVR-1, hVR-2, and rVR-2 substrate), an hVR-1, hVR-2, and rVR-2 antibody, an hVR-1, hVR-2, and rVR-2 agonist or antagonist, a peptidomimetic of an hVR-1, hVR-2, and rVR-2 agonist or antagonist, or other small molecule. In one embodiment, the agent stimulates one or more hVR-1, hVR-2, and rVR-2 activities. Examples of such stimulatory agents include active hVR-1, hVR-2, and rVR-2 protein and a nucleic acid molecule encoding hVR-1, hVR-2, and rVR-2 that has been introduced into the cell. In another embodiment, the agent inhibits one or more hVR-1, hVR-2, and rVR-2 activities. Examples of such inhibitory agents include antisense hVR-1, hVR-2, and rVR-2 nucleic acid molecules, anti-hVR-1, hVR-2, and rVR-2 antibodies, and hVR-1, hVR-2, and rVR-2 inhibitors. These modulatory methods can be performed *in vitro* (e.g., by culturing the cell with the agent) or, alternatively, *in vivo* (e.g., by administering the agent to a subject). As such, the present invention provides methods of treating an individual afflicted with a disease or disorder characterized by aberrant expression or activity of an hVR-1, hVR-2, and rVR-2 protein or nucleic acid molecule. In one embodiment, the method involves administering an agent (e.g., an agent identified by a screening assay described herein), or combination of agents that modulates (e.g., upregulates or downregulates) hVR-1, hVR-2, and rVR-2 expression or activity. In another embodiment, the method involves administering an hVR-1, hVR-2, and rVR-2 protein or nucleic acid molecule as therapy to compensate for reduced or aberrant hVR-1, hVR-2, and rVR-2 expression or activity.

Stimulation of hVR-1, hVR-2, and rVR-2 activity is desirable in situations in which hVR-1, hVR-2, and rVR-2 is abnormally downregulated and/or in which increased hVR-1, hVR-2, and rVR-2 activity is likely to have a beneficial effect. For example, stimulation of hVR-1, hVR-2, and rVR-2 activity is desirable in situations in which an hVR-1, hVR-2, and rVR-2 is downregulated and/or in which increased hVR-1, hVR-2, and rVR-2 activity is likely to have a beneficial effect. Likewise, inhibition of hVR-1, hVR-2, and rVR-2 activity is desirable in situations in which hVR-

1, hVR-2, and rVR-2 is abnormally upregulated and/or in which decreased hVR-1, hVR-2, and rVR-2 activity is likely to have a beneficial effect.

3. Pharmacogenomics

5 The hVR-1, hVR-2, and rVR-2 molecules of the present invention, as well as agents, or modulators which have a stimulatory or inhibitory effect on hVR-1, hVR-2, and rVR-2 activity (e.g., hVR-1, hVR-2, and rVR-2 gene expression) as identified by a screening assay described herein can be administered to individuals to treat (prophylactically or therapeutically) hVR-1, hVR-2, and rVR-2-associated disorders (e.g., pain disorders) associated with aberrant hVR-1, hVR-2, and rVR-2 activity. In conjunction with such treatment, pharmacogenomics (i.e., the study of the relationship between an individual's genotype and that individual's response to a foreign compound or drug) may be considered. Differences in metabolism of therapeutics can lead to severe toxicity or therapeutic failure by altering the relation between dose and blood concentration of the pharmacologically active drug. Thus, a physician or clinician may consider applying knowledge obtained in relevant pharmacogenomics studies in determining whether to administer an hVR-1, hVR-2, and rVR-2 molecule or hVR-1, hVR-2, and rVR-2 modulator as well as tailoring the dosage and/or therapeutic regimen of treatment with an hVR-1, hVR-2, and rVR-2 molecule or hVR-1, hVR-2, and rVR-2 modulator.

Pharmacogenomics deals with clinically significant hereditary variations in the response to drugs due to altered drug disposition and abnormal action in affected persons. See, for example, Eichelbaum, M. *et al.* (1996) *Clin. Exp. Pharmacol. Physiol.* 23(10-11):983-985 and Linder, M.W. *et al.* (1997) *Clin. Chem.* 43(2):254-266. In general, two types of pharmacogenetic conditions can be differentiated. Genetic conditions transmitted as a single factor altering the way drugs act on the body (altered drug action) or genetic conditions transmitted as single factors altering the way the body acts on drugs (altered drug metabolism). These pharmacogenetic conditions can occur either as rare genetic defects or as naturally-occurring polymorphisms. For example, glucose-6-phosphate dehydrogenase deficiency (G6PD) is a common inherited enzymopathy in which the main clinical complication is haemolysis after ingestion of oxidant drugs (anti-malarials, sulfonamides, analgesics, nitrofurans) and consumption of fava beans.

One pharmacogenomics approach to identifying genes that predict drug response, known as "a genome-wide association", relies primarily on a high-resolution map of the human genome consisting of already known gene-related markers (e.g., a "bi-allelic" gene marker map which consists of 60,000-100,000 polymorphic or variable sites on the human genome, each of which has two variants.) Such a high-resolution genetic map can be compared to a map of the genome of each of a statistically significant number of patients taking part in a Phase II/III drug trial to identify markers associated with a particular observed drug response or side effect. Alternatively, such a high resolution map can be generated from a combination of some ten-million known single nucleotide polymorphisms (SNPs) in the human genome. As used herein, a "SNP" is a common

alteration that occurs in a single nucleotide base in a stretch of DNA. For example, a SNP may occur once per every 1000 bases of DNA. A SNP may be involved in a disease process, however, the vast majority may not be disease-associated. Given a genetic map based on the occurrence of such SNPs, individuals can be grouped into genetic categories depending on a particular pattern of SNPs in their individual genome. In such a manner, treatment regimens can be tailored to groups of genetically similar individuals, taking into account traits that may be common among such genetically similar individuals.

Alternatively, a method termed the "candidate gene approach", can be utilized to identify genes that predict drug response. According to this method, if a gene that encodes a drug's target is known (*e.g.*, an hVR-1, hVR-2, and rVR-2 protein of the present invention), all common variants of that gene can be fairly easily identified in the population and it can be determined if having one version of the gene versus another is associated with a particular drug response.

As an illustrative embodiment, the activity of drug metabolizing enzymes is a major determinant of both the intensity and duration of drug action. The discovery of genetic polymorphisms of drug metabolizing enzymes (*e.g.*, N-acetyltransferase 2 (NAT 2) and cytochrome P450 enzymes CYP2D6 and CYP2C19) has provided an explanation as to why some patients do not obtain the expected drug effects or show exaggerated drug response and serious toxicity after taking the standard and safe dose of a drug. These polymorphisms are expressed in two phenotypes in the population, the extensive metabolizer (EM) and poor metabolizer (PM). The prevalence of PM is different among different populations. For example, the gene coding for CYP2D6 is highly polymorphic and several mutations have been identified in PM, which all lead to the absence of functional CYP2D6. Poor metabolizers of CYP2D6 and CYP2C19 quite frequently experience exaggerated drug response and side effects when they receive standard doses. If a metabolite is the active therapeutic moiety, PM show no therapeutic response, as demonstrated for the analgesic effect of codeine mediated by its CYP2D6-formed metabolite morphine. The other extreme are the so called ultra-rapid metabolizers who do not respond to standard doses. Recently, the molecular basis of ultra-rapid metabolism has been identified to be due to CYP2D6 gene amplification.

Alternatively, a method termed the "gene expression profiling", can be utilized to identify genes that predict drug response. For example, the gene expression of an animal dosed with a drug (*e.g.*, an hVR-1, hVR-2, and rVR-2 molecule or hVR-1, hVR-2, and rVR-2 modulator of the present invention) can give an indication whether gene pathways related to toxicity have been turned on.

Information generated from more than one of the above pharmacogenomics approaches can be used to determine appropriate dosage and treatment regimens for prophylactic or therapeutic treatment of an individual. This knowledge, when applied to dosing or drug selection, can avoid adverse reactions or therapeutic failure and thus enhance therapeutic or prophylactic efficiency when treating a subject with an hVR-1, hVR-2, and rVR-2 molecule or hVR-1, hVR-2, and rVR-2

modulator, such as a modulator identified by one of the exemplary screening assays described herein.

This invention is further illustrated by the following examples which should not be construed as limiting. The contents of all references, patents and published patent applications cited throughout this application, as well as the Figures and the Sequence Listing are incorporated herein by reference.

EXAMPLES

EXAMPLE 1: IDENTIFICATION AND CHARACTERIZATION OF hVR-1, hVR-2, and rVR-2 cDNA

In this example, the identification and characterization of the genes encoding hVR-1 (clone Fchrb87a6), hVR-2 (clone flh21e11), hVR-2 alternate form (clone frhob12c4), and rVR-2 (clone flrxb147g11) are described.

Isolation of the hVR-1, hVR-2, and the rVR-2 cDNA

The invention is based, at least in part, on the discovery of two human genes and one rat gene encoding novel members of the Capsaicin/Vanilloid receptor family, referred to herein as hVR-1, hVR-2, and rVR-2, respectively. These clones were identified from a human heart library and a rat dorsal root ganglion (DRG) library, based on sequence homology to the known rat VR-1 (Accession Number AF029310). The sequence of the two human clones and the rat clone was determined and found to contain open reading frames.

The nucleotide sequence of the full length hVR-1 cDNA and the predicted amino acid sequence of the hVR-1 polypeptide are shown in Figure 1 and in SEQ ID NOs:1 and 2, respectively.

The nucleotide sequence of the full length hVR-2 cDNA and the predicted amino acid sequence of the hVR-2 polypeptide are shown in Figure 2 and in SEQ ID NOs:4 and 5, respectively.

The nucleotide sequence of the partial hVR-2 (alternate form) cDNA and the predicted amino acid sequence of the hVR-2 (alternate form) polypeptide are shown in Figure 3 and in SEQ ID NOs:7 and 8, respectively.

The amino acid sequence of the predicted full length human VR-2 protein (alternate form) is shown in Figure 16 and in SEQ ID NO:20.

The nucleotide sequence of the partial rVR-2 cDNA and the predicted amino acid sequence of the rVR-2 polypeptide are shown in Figure 4 and in SEQ ID NOs:10 and 11, respectively.

Analysis of the hVR-1, hVR-2, and rVR-2 Molecules

The hVR-1 protein (SEQ ID NO:2) was aligned with the human VR-2 protein (SEQ ID NO:5) using the GAP program in the GCG software package (Blosum 62 matrix) and a gap weight of 12 and a length weight of 4. The results showed a 46.348% identity and 55.378% similarity between the two sequences (see Figure 5).

The hVR-1 nucleotide sequence (SEQ ID NO:1) was aligned with the human VR-2 nucleotide sequence (SEQ ID NO:4) using the GAP program in the GCG software package (nwsgapdna matrix) and a gap weight of 50 and a length weight of 3. The results showed a 55.316% identity and 55.316% similarity between the two sequences (see Figure 6).

The hVR-2 protein (SEQ ID NO:5) was aligned with the rat VR-2 protein (SEQ ID NO:11) using the CLUSTAL W (1.74) multiple sequence alignment program (Figure 7), as well as using the GAP program in the GCG software package (Blosum 62 matrix) and a gap weight of 12 and a length weight of 4. The results showed a 79.167% identity and 81.703% similarity between the two sequences (see Figure 8).

The hVR-1 nucleotide sequence (SEQ ID NO:1) was aligned with the rat VR-1 nucleotide sequence (Accession Number:AF029310) using the GAP program in the GCG software package (nwsgapdna matrix) and a gap weight of 50 and a length weight of 3. The results showed a 82.125% identity and 82.125% similarity between the two sequences (see Figure 9).

The hVR-1 protein (SEQ ID NO:2) was aligned with the rat VR-1 protein (Accession Number:AF029310) using the GAP program in the GCG software package (Blosum 62 matrix) and a gap weight of 12 and a length weight of 4. The results showed a 86.022% identity and 89.247% similarity between the two sequences (see Figure 10).

The hVR-2 protein (SEQ ID NO:5) was aligned with the human VR-2 protein (alternate form) (SEQ ID NO:8) using the CLUSTAL W (1.74) multiple sequence alignment program (Figure 11).

Finally, the hVR-2 protein (SEQ ID NO:5) was aligned with the predicted full length human VR-2 protein (alternate form) (SEQ ID NO:20) using the CLUSTAL W (1.74) multiple sequence alignment program (Figure 17).

A search was performed against the HMM database resulting in the identification of three ankyrin repeat domains in the amino acid sequence of hVR-1 (SEQ ID NO:2) at about residues 201-233, 248-283, and 333-361, and in the amino acid sequence of hVR-2 (SEQ ID NO:5) at about residues 162-194, 208-243, and 293-328. The results of the searches are set forth in Figures 13 and 15, respectively.

Hydropathy plots have identified 6 transmembrane domains in the hVR-1 and the hVR-2 proteins (see Figures 12 and 14, respectively).

A series of searches have revealed that the hVR-1 protein matches the ProDom entry 141801 for the vanilloid receptor subtype and the ProDom entry 145518 for the vanilloid receptor subtype.

Moreover, a search was performed against the Prosite database resulting in the identification of four N-glycosylation sites in the amino acid sequence of SEQ ID NO:5 (at about residues 171-174, 192-195, 604-607, and 749-752), three cGMP-dependent protein kinase phosphorylation sites in the amino acid sequence of SEQ ID NO:5 (at about residues 2-5, 368-371, and 499-502), a series of protein kinase C and Casein kinase II phosphorylation sites in the amino acid sequence of SEQ ID NO:5, two tyrosine kinase phosphorylation sites in the amino acid sequence of SEQ ID NO:5 (at about residues 368-375 and 622-628), and two myristoylation sites in the amino acid sequence of SEQ ID NO:5 (at about residues 169-174 and 765-770).

Tissue Distribution of hVR-1 and hVR-2 mRNA

This Example describes the tissue distribution of hVR-1 and hVR-2 mRNA as determined by *in situ* hybridization.

For *in situ* analysis, tissues, such as brain regions and whole brain, obtained from human and monkey were first frozen on dry ice. Ten-micrometer-thick coronal sections of the tissues were postfixed with 4% formaldehyde in DEPC treated 1X phosphate- buffered saline at room temperature for 10 minutes before being rinsed twice in DEPC 1X phosphate-buffered saline and once in 0.1 M triethanolamine-HCl (pH 8.0). Following incubation in 0.25% acetic anhydride-0.1 M triethanolamine-HCl for 10 minutes, sections were rinsed in DEPC 2X SSC (1X SSC is 0.15M NaCl plus 0.015M sodium citrate). Tissue was then dehydrated through a series of ethanol washes, incubated in 100% chloroform for 5 minutes, and then rinsed in 100% ethanol for 1 minute and 95% ethanol for 1 minute and allowed to air dry.

Hybridizations were performed with ³⁵S-radiolabeled (5 X 10⁷ cpm/ml) cRNA probes. Probes were incubated in the presence of a solution containing 600 mM NaCl, 10 mM Tris (pH 7.5), 1 mM EDTA, 0.01% sheared salmon sperm DNA, 0.01% yeast tRNA, 0.05% yeast total RNA type X1, 1 X Denhardt's solution, 50% formamide, 10% dextran sulfate, 100 mM dithiothreitol, 0.1% sodium dodecyl sulfate (SDS), and 0.1% sodium thiosulfate for 18 hours at 55°C.

After hybridization, slides were washed with 2 X SSC. Sections were then sequentially incubated at 37°C in TNE (a solution containing 10 mM Tris-HCl (pH 7.6), 500 mM NaCl, and 1 mM EDTA), for 10 minutes, in TNE with 10µg of RNase A per ml for 30 minutes, and finally in TNE for 10 minutes. Slides were then rinsed with 2 X SSC at room temperature, washed with 2 X SSC at 50°C for 1 hour, washed with 0.2 X SSC at 55°C for 1 hour, and 0.2 X SSC at 60°C for 1 hour. Sections were then dehydrated rapidly through serial ethanol-0.3 M sodium acetate concentrations before being air dried and exposed to Kodak Biomax MR scientific imaging film for 24 hours and subsequently dipped in NB-2 photoemulsion and exposed at 4°C for 7 days before being developed and counter stained.

The data indicate that the hVR-1 molecule is not expressed in human nor monkey brain. The hVR-1 molecule is expressed in nodose, trigeminal sensory neurons, but is not expressed in sympathetic neurons. Within the nodose sensory neurons and trigeminal sensory neurons, expression was seen in distinct sub-populations. Moreover, hVR1 is expressed in some, but not all, small dorsal root ganglion (DRG) neurons and in a few medium sized DRG neurons. The hVR-1 molecule is partially co-expressed with the neuropeptide CGRP and with substance P which are present in nociceptive neurons.

The data further indicate that the VR-2 molecule is expressed in both human and monkey brain, primarily in cortical neurons. The VR2 molecule is also expressed in other brain regions, for example, the thalamus, striatum, hippocampus, hypothalamus, midbrain, medula and brain stem. In addition, the VR-2 molecule is expressed in parasympathetic neurons of the monkey heart (atrium), nodose sensory neurons, trigeminal (TRG) sensory neurons, dorsal root ganglion sensory neurons, sympathetic neurons, and motor neurons of the spinal cord. The VR2 molecule is widely expressed in TRG and DRG neurons, being present in most small and medium sized neurons and also in a few of the large neurons. VR2, like VR-1, partially co-localizes with CGRP and substance P.

Trigeminal sensory neurons are recognized pain centers while sympathetic neurons have been shown to be involved in neuropathic pain.

EXAMPLE 2: EXPRESSION OF RECOMBINANT hVR-1, hVR-2, AND rVR-2 PROTEIN IN BACTERIAL CELLS

In this example, hVR-1, hVR-2, and rVR-2 is expressed as a recombinant glutathione-S-transferase (GST) fusion polypeptide in *E. coli* and the fusion polypeptide is isolated and characterized. Specifically, hVR-1, hVR-2, and rVR-2 is fused to GST and this fusion polypeptide is expressed in *E. coli*, e.g., strain PEB199. Expression of the GST-hVR-1, GST-hVR-2, and GST-rVR-2 fusion protein in PEB199 is induced with IPTG. The recombinant fusion polypeptide is purified from crude bacterial lysates of the induced PEB199 strain by affinity chromatography on glutathione beads. Using polyacrylamide gel electrophoretic analysis of the polypeptide purified from the bacterial lysates, the molecular weight of the resultant fusion polypeptide is determined.

EXAMPLE 3: EXPRESSION OF RECOMBINANT hVR-1, hVR-2, AND rVR-2 PROTEIN IN COS CELLS

To express the hVR-1, hVR-2, and rVR-2 gene in COS cells, the pcDNA/Amp vector from Invitrogen Corporation (San Diego, CA) is used. This vector contains an SV40 origin of replication, an ampicillin resistance gene, an *E. coli* replication origin, a CMV promoter followed by a polylinker region, and an SV40 intron and polyadenylation site. A DNA fragment encoding the entire hVR-1, hVR-2, and rVR-2 protein and an HA tag (Wilson *et al.* (1984) *Cell* 37:767) or a

FLAG tag fused in-frame to its 3' end of the fragment is cloned into the polylinker region of the vector, thereby placing the expression of the recombinant protein under the control of the CMV promoter.

To construct the plasmid, the hVR-1, hVR-2, and rVR-2 DNA sequence is amplified by PCR using two primers. The 5' primer contains the restriction site of interest followed by approximately twenty nucleotides of the hVR-1, hVR-2, and rVR-2 coding sequence starting from the initiation codon; the 3' end sequence contains complementary sequences to the other restriction site of interest, a translation stop codon, the HA tag or FLAG tag and the last 20 nucleotides of the hVR-1, hVR-2, and rVR-2 coding sequence. The PCR amplified fragment and the pCDNA/Amp vector are digested with the appropriate restriction enzymes and the vector is dephosphorylated using the CIAP enzyme (New England Biolabs, Beverly, MA). Preferably the two restriction sites chosen are different so that the hVR-1, hVR-2, and rVR-2 gene is inserted in the correct orientation. The ligation mixture is transformed into *E. coli* cells (strains HB101, DH5a, SURE, available from Stratagene Cloning Systems, La Jolla, CA, can be used), the transformed culture is plated on ampicillin media plates, and resistant colonies are selected. Plasmid DNA is isolated from transformants and examined by restriction analysis for the presence of the correct fragment.

COS cells are subsequently transfected with the hVR-1, hVR-2, and rVR-2-pcDNA/Amp plasmid DNA using the calcium phosphate or calcium chloride co-precipitation methods, DEAE-dextran-mediated transfection, lipofection, or electroporation. Other suitable methods for transfecting host cells can be found in Sambrook, J., Fritsh, E. F., and Maniatis, T. *Molecular Cloning: A Laboratory Manual*. 2nd, ed., Cold Spring Harbor Laboratory, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, 1989. The expression of the hVR-1, hVR-2, and rVR-2 polypeptide is detected by radiolabelling (³⁵S-methionine or ³⁵S-cysteine available from NEN, Boston, MA, can be used) and immunoprecipitation (Harlow, E. and Lane, D. *Antibodies: A Laboratory Manual*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, 1988) using an HA specific monoclonal antibody. Briefly, the cells are labelled for 8 hours with ³⁵S-methionine (or ³⁵S-cysteine). The culture media are then collected and the cells are lysed using detergents (RIPA buffer, 150 mM NaCl, 1% NP-40, 0.1% SDS, 0.5% DOC, 50 mM Tris, pH 7.5). Both the cell lysate and the culture media are precipitated with an HA specific monoclonal antibody. Precipitated polypeptides are then analyzed by SDS-PAGE.

Alternatively, DNA containing the hVR-1, hVR-2, and rVR-2 coding sequence is cloned directly into the polylinker of the pCDNA/Amp vector using the appropriate restriction sites. The resulting plasmid is transfected into COS cells in the manner described above, and the expression of the hVR-1, hVR-2, and rVR-2 polypeptide is detected by radiolabelling and immunoprecipitation using an hVR-1, hVR-2, and rVR-2 specific monoclonal antibody.

EXAMPLE 4: ELECTROPHYSIOLOGICAL STUDIES OF VR2

Human VR2 was functionally characterized in both HEK293 cells and *Xenopus* oocytes using electrophysiological methods. VR2 (in the pcDNA3.1 vector purchased by Invitrogen) was transiently expressed in HEK293 cells (ATCC) and recordings were performed 48 hours after transfection of cells using the whole-cell patch-clamp method (described in Bertil Hille, *Ion Channels of excitable membranes*, 1992; Hammill *et al.* (1981) *Plüger Arch.* 391:85-100). The results indicate that heat stimulation (>50 °C) induces a rapid inactivating inward current (1-2 nA). Heat-evoked currents of VR2 displayed profound desensitization and could be reversibly blocked by the VR1 inhibitor capsazepin (at a 10 µM concentration). In contrast to rat VR1, Capsaicin (at a 1-10 µM concentration), resiniferatoxin (at a 0.1-3 µM concentration), and low pH (5.0-6.0) do not induce any currents from VR2. Binding studies of [³H]-resiniferatoxin (NEN) to both human VR1 and VR2 in membranes isolated from HEK293 cell homogenates also indicate that resiniferatoxin (at a 0.1-10 nM concentration) has no specific binding to VR2 while it binds to human VR1 with high affinities.

For the oocyte studies, human VR2 was subcloned into an oocyte expression vector containing 5'- and 3'-UTR of *Xenopus* β-globin (Chiara *et al.* (1999) *Biochemistry* 38(20)6689-6698). *In vitro* transcription was carried out as described in Chiara *et al.* (*supra*) and cRNA (10-100 ng) was then injected into the oocytes. VR2 function was characterized in the oocytes 48 hours after cRNA injection using a standard two-electrode voltage-clamp. Consistent with the data from the HEK293 studies, VR2 can only be activated by heat stimulation (48-50 °C) but not by vanilloid receptor agonists, capsaicin, or resiniferatoxin. The vanilloid receptor antagonist capsazepine (at a 1-10 µM concentration) blocks the heat response of VR2 reversibly.

EXAMPLE 5: GENERATION OF ANTI-hVR-2 ANTIBODIES AND hVR-2 PROTEIN LOCALIZATION BY IMMUNOSTAINING

Polyclonal antisera were raised in rabbits against the following three peptides derived from the human VR2 amino acid sequence, using the techniques described in Ed Harlow and David Lane (1988) "Antibodies; A Laboratory Manual" Cold Spring harbor Laboratory Press.

Antibody PEPTIDE 1: AFHCKSPHRHRMVVLE (SEQ ID NO:13)

Antibody PEPTIDE 2: RPEAPTGNATESVQPMEGQEDEGN (SEQ ID NO:14)

Antibody PEPTIDE 3: SVLEMENGYWWCRKKQRAG (SEQ ID NO:15)

Antisera were subsequently affinity purified using the peptide immunogen.

The polyclonal antisera were tested for immunostaining of both monkey and rat dorsal root ganglion sensory neurons. Peptides 1 and 3 gave specific staining of subpopulations of sensory neurons that was competed with the corresponding peptide. This pattern of expression was very similar to the one observed using a VR-2 riboprobe.

EXAMPLE 6: CHROMOSOMAL LOCALIZATION OF hVR-1 AND hVR-2

To chromosomally map the hVR-1 gene, primers were designed based on the sequence of hVR-1 (clone Fchrb87a6) (amplifying a 177 bp product from a human control cell line DNA and multiple faint larger products from a control Hamster cell line DNA by PCR). These primers were used to amplify 93 DNAs in duplicate from the Genebridge 4 Radiation Hybrid Panel (Research Genetics, Inc., Huntsville, AL).

The hVR-1 primers used in the PCR mapping studies were: forward - TAGGAGACCCCGTTGCCACG (SEQ ID NO:16) and reverse - GATTCACTTGGGGACAGTGACG (SEQ ID NO:17) and the PCR reactions were performed as follows: 5 µl Template DNA (10ng/µl), 1.5µl 10X Perkin Elmer PCR Buffer, 1.2µl Pharmacia dNTP mix 2.5 mM, 1.15µl Forward primer 6.6µM, 1.15µl Reverse primer 6.6µM, 5µl Gibco/BRL Platinum Taq .05U/µl (Hot Start), using an amplification profile of: 95°C for 10 minutes followed by 35 Cycles of 94°C for 40 seconds, 55°C for 40 seconds, 72°C for 40 seconds, and 72°C for 5 minutes. The PCR products were run on 2% agarose gels, post-stained with SYBR Gold (1:10,000 dilution in 1X TBE), and scanned on a Molecular Dynamics 595 Fluorimager.

The following is the vector data for the 93 Genebridge4 hybrid DNAs. These are in order 1-93. A "1" is a positive result, a "-" is a negative result, a "?" is an ambiguous result.

hVR1 1 - - 1 ? - 1 - 1 - 1 1 - - - 1 - - 1 - 1 1 - - 1 - 1 - 1 - - 1 1 - 1 - - - - - 1 - 1 1 1 - - - 1 - - - 1 1 -
- - - 1 - - - - 1 - 1 - - 1 1 1 - 1 - 1 - - - 1 1 - 1 - - - - - 1

RH linkage analysis was performed using the Map Manager QTb28 software package.

hVR1 was found to map to the p arm of human chromosome 17, 18.9 cR₃₀₀₀ telomeric to the Whitehead Institute framework marker WI-6584, and 7.7 cR₃₀₀₀ centromeric of the Whitehead framework marker WI-5436. LOD scores for linkage were 14.5 for WI-6584 and 19.3 for WI-5436. This region corresponds to the cytogenetic location 17p12-13. This region is syntenic to mouse chromosome 11.

To chromosomally map the hVR-2 gene, primers were designed from 5' UTR sequence of human VR2 (clone Flh21e11) (amplifying a 166 bp product from a human control cell line DNA and 2 much larger faint bands from a control Hamster cell line DNA by PCR). These primers were used to amplify 93 DNAs in duplicate from the Genebridge 4 Radiation Hybrid Panel (Research Genetics, Inc., Huntsville, AL).

The hVR-2 primers used in the PCR mapping studies were: forward - TTAAGCTCCCGTTCTCACCG (SEQ ID NO:18) and reverse -GCTGCGGGAGGAAGTGAAGC (SEQ ID NO:19) and the PCR reactions were performed as follows: 5µl Template DNA (10ng/µl), 1.5µl 10X Perkin Elmer PCR Buffer, 1.2µl Pharmacia dNTP mix 2.5mM, 1.15µl Forward primer 6.6µM, 1.15µl Reverse primer 6.6µM, 5µl Gibco/BRL Platinum Taq .05U/µl (Hot Start), using an amplification profile of 95°C for 10 minutes, followed by 35 Cycles of 94°C for 40 seconds, 55°C

for 40 seconds, 72°C for 40 seconds, and 72°C for 5 minutes. The PCR products were run on 2% agarose gels, post-stained with SYBR Gold (1:10,000 dilution in 1X TBE), and scanned on a Molecular Dynamics 595 Fluorimager.

The following is the vector data for the 93 Genebridge4 hybrid DNAs. These are in order 1-93. A "1" is a positive result, a "-" is a negative result, a "?" is an ambiguous result.

hVR2 1 - - 1 1 - ? 1 1 - - - - - 1 - - 1 - 1 1 - - - - 1 - 1 1 1 - - 1 - - - - - 1 1 - 1 1 1 - - 1 1 - - - 1 1
- - - - 1 1 1 - - - 1 - 1 1 - 1 - 1 1 1 1 1 - - - 1 1 - 1 - 1 - ? - - - ?

RH linkage analysis was performed using the Map Manager QTb28 software package.

hVR2 was found to map to the p arm of human chromosome 17, 29.3cR cR₃₀₀₀ telomeric to the Whitehead Institute framework marker D17S721, and 23.3 cR₃₀₀₀ centromeric of the Whitehead framework marker AFMA043ZB5. LOD scores for linkage were 11.9 for D17S721 and 13.6 for AFMA043ZB5. This region corresponds to the cytogenetic location 17p11-12. This region is syntenic to mouse chromosome 11.

Equivalents

Those skilled in the art will recognize, or be able to ascertain using no more than routine experimentation, many equivalents to the specific embodiments of the invention described herein. Such equivalents are intended to be encompassed by the following claims.

What is claimed:

1. An isolated nucleic acid molecule selected from the group consisting of:
 - (a) a nucleic acid molecule comprising the nucleotide sequence set forth in SEQ ID NO:1, 3, 4, 6, 7, 9, 10, or 12 or a complement thereof; and
 - (b) a nucleic acid molecule consisting of the nucleotide sequence set forth in SEQ ID NO:1, 3, 4, 6, 7, 9, 10, or 12 or a complement thereof.
2. An isolated nucleic acid molecule which encodes a polypeptide selected from the group consisting of:
 - (a) a polypeptide comprising the amino acid sequence set forth in SEQ ID NO:2, 5, 8, or 11; and
 - (b) a polypeptide consisting of the amino acid sequence set forth in SEQ ID NO:2, 5, 8, or 11.
3. An isolated nucleic acid molecule which encodes a naturally occurring allelic variant of a polypeptide comprising the amino acid sequence set forth in SEQ ID NO:2, 5, 8, or 11.
4. An isolated nucleic acid molecule selected from the group consisting of:
 - a) a nucleic acid molecule comprising a nucleotide sequence which is at least 83% identical to the nucleotide sequence of SEQ ID NO:1, 3, 4, 6, 7, 9, 10, or 12, or a complement thereof;
 - b) a nucleic acid molecule comprising a fragment of at least 20 nucleotides of a nucleic acid comprising the nucleotide sequence of SEQ ID NO:1, 3, 4, 6, 7, 9, 10, or 12, or a complement thereof;
 - c) a nucleic acid molecule which encodes a polypeptide comprising an amino acid sequence at least about 87% identical to the amino acid sequence of SEQ ID NO:2, 5, 8, or 11; and
 - d) a nucleic acid molecule which encodes a fragment of a polypeptide comprising the amino acid sequence of SEQ ID NO:2, 5, 8, or 11, wherein the fragment comprises at least 15 contiguous amino acid residues of the amino acid sequence of SEQ ID NO:2, 5, 8, or 11.
5. An isolated nucleic acid molecule comprising the nucleic acid molecule of any one of claims 1, 2, 3, or 4, and a nucleotide sequence encoding a heterologous polypeptide.
6. A vector comprising the nucleic acid molecule of any one of claims 1, 2, 3, or 4.

7. The vector of claim 6, which is an expression vector.

8. A host cell transfected with the expression vector of claim 7.

5 9. A method of expressing a polypeptide comprising culturing the host cell of claim 8 in an appropriate culture medium to, thereby, express the polypeptide.

10. An isolated polypeptide selected from the group consisting of:

10 a) a fragment of a polypeptide comprising the amino acid sequence of SEQ ID NO:2, 5, 8, or 11, wherein the fragment comprises at least 15 contiguous amino acids of SEQ ID NO:2, 5, 8, or 11;

15 b) a naturally occurring allelic variant of a polypeptide comprising the amino acid sequence of SEQ ID NO:2, 5, 8, or 11, wherein the polypeptide is encoded by a nucleic acid molecule which hybridizes to a nucleic acid molecule consisting of SEQ ID NO:1, 3, 4, 6, 7, 9, 10, or 12 under stringent conditions;

20 c) a polypeptide which is encoded by a nucleic acid molecule comprising a nucleotide sequence which is at least 83% identical to a nucleic acid comprising the nucleotide sequence of SEQ ID NO:1, 3, 4, 6, 7, 9, 10, or 12; and

25 d) a polypeptide comprising an amino acid sequence which is at least 87% identical to the amino acid sequence of SEQ ID NO:2, 5, 8, or 11.

11. The isolated polypeptide of claim 10 comprising the amino acid sequence of SEQ ID NO:2, 5, 8, or 11.

12. The polypeptide of claim 10, further comprising heterologous amino acid sequences.

13. An antibody which selectively binds to a polypeptide of claim 10.

30 14. A method for detecting the presence of a polypeptide of claim 10 in a sample comprising:

a) contacting the sample with a compound which selectively binds to the polypeptide; and

35 b) determining whether the compound binds to the polypeptide in the sample to thereby detect the presence of a polypeptide of claim 10 in the sample.

15. The method of claim 14, wherein the compound which binds to the polypeptide is an antibody.

16. A kit comprising a compound which selectively binds to a polypeptide of claim 10 and instructions for use.

17. A method for detecting the presence of a nucleic acid molecule of any one of claims 1, 2, 3, or 4 in a sample comprising:

a) contacting the sample with a nucleic acid probe or primer which selectively hybridizes to the nucleic acid molecule; and

b) determining whether the nucleic acid probe or primer binds to a nucleic acid molecule in the sample to thereby detect the presence of a nucleic acid molecule of any one of claims 1, 2, 3, or 4 in the sample.

18. The method of claim 17, wherein the sample comprises mRNA molecules and is contacted with a nucleic acid probe.

19. A kit comprising a compound which selectively hybridizes to a nucleic acid molecule of any one of claims 1, 2, 3, or 4 and instructions for use.

20. A method for identifying a compound which binds to a polypeptide of claim 10 comprising:

a) contacting the polypeptide, or a cell expressing the polypeptide with a test compound; and

b) determining whether the polypeptide binds to the test compound.

21. The method of claim 20, wherein the binding of the test compound to the polypeptide is detected by a method selected from the group consisting of:

a) detection of binding by direct detection of test compound/polypeptide binding;

b) detection of binding using a competition binding assay; and

c) detection of binding using an assay for hVR-1, hVR-2, or rVR-2 activity.

22. A method for modulating the activity of a polypeptide of claim 10 comprising contacting the polypeptide or a cell expressing the polypeptide with a compound which binds to the polypeptide in a sufficient concentration to modulate the activity of the polypeptide.

23. A method for identifying a compound which modulates the activity of a polypeptide of claim 10 comprising:

a) contacting a polypeptide of claim 10 with a test compound; and

b) determining the effect of the test compound on the activity of the polypeptide to thereby identify a compound which modulates the activity of the polypeptide.

24. A method for treating a subject having a disorder characterized by aberrant hVR-1 or hVR-2 protein activity or nucleic acid expression comprising administering to the subject a hVR-1 or hVR-2 modulator such that treatment of the subject occurs.

25. The method of claim 24, wherein the hVR-1 or hVR-2 modulator is a small molecule.

26. The method of claim 24, wherein the disorder is a pain disorder.

**NOVEL MEMBERS OF THE CAPSAICIN/VANILLOID RECEPTOR FAMILY OF
PROTEINS AND USES THEREOF**

Abstract of the Disclosure

5

10

The invention provides isolated nucleic acids molecules, designated hVR-1, hVR-2, and rVR-2 nucleic acid molecules, which encode novel members of the Capsaicin/Vanilloid receptor family. The invention also provides antisense nucleic acid molecules, recombinant expression vectors containing hVR-1, hVR-2, and rVR-2 nucleic acid molecules, host cells into which the expression vectors have been introduced, and nonhuman transgenic animals in which an hVR-1, hVR-2, and rVR-2 gene has been introduced or disrupted. The invention still further provides isolated hVR-1, hVR-2, and rVR-2 proteins, fusion proteins, antigenic peptides and anti-hVR-1, anti-hVR-2, and anti-rVR-2 antibodies. Diagnostic methods utilizing compositions of the invention are also provided.

US
Patent
Attorney

humanVR1 gene with translation of open reading frame

Input file Fchrb87a6.seq; Output File Fchrb87a6.tra
Sequence length 3909

GTGAGCGCAACGCACTGCGGGCAGTGAGCGCAACGCACTGCGGGCAGTGAGCGCAACGCACTGCGGGCAGTGAGCGCAA
GCACTGCGGGCAGTGAGCGCAACGCACTGCGGGCAGTGAGCGCAACGCACTGCGGGCAGTGAGCGCAACGCACTGCGG
CAGTGAGCGCAACGCACTTTCGGGGCAGTGAGCGCAACGCACTGCGGGCAGTGAGCGCAACGCACTGCGGGCAGTGAGC
GCAACGCACTGCGGGCAGTGAGCGCAACGCACTGCGGGCAGTGAGCGCAACGCACTGCGGGCAGTGAGCGCAACGCACT
GCGGGCAGTGAGCGCAACGCACTGCGGGCAGTGAGCGCAACGCACTGCGGGCAGTGAGCGCAACGCACTGCGGGCAGTG
AGCGCAACGCACTGCGGGCAGTGAGCGCAACGCACTTAATGTGAGTTAGCTCACTCATTAGGCACCCCAGGCTTTACAC
TTTATGCTTCCGGCTCGTATGTTGTGTGGAATTGTGAGCGGATAACAATTTACACAGGAAACAGCTATGACCATGATT
ACGCCAAGCTCTAATACGACTCACTATAGGGAAAGCTGGTACGCCTGCAGGTACCGGTCCGGAATTCCCGGGTTCGACCC
ACGCGTCCGAAAACACACCTCTCTGCTGTGGGAAGACTGTGCAATGGCACAGCCGCAGAGCTTGGTTTGGGAGGTTGAA
GTGCTCTGGGGAGAATTTCGTAGATCATCCTCAGAAAAGCCTTGCCCTGGTGTCTTACCAGAAAAACGTCTCCCAATCAC
CCAGAAAAGCTGTCCACAGTAGTCCCCCTTATCCACGGGTGTCACTTTCCATGGGTTCACTTATTTGCGGTCAACCAC
GCTCTGCCAATATTAAATGGAAAATTCCTCAAACAGTTCCCAAGTTTCCCTTGTGCATTGTTCTGAGCAGTGTGATGA
AGAGTCTCTGCCGTGCCATCTGGGATGCAAACCGTCCCTGTGTCCCCCACGTCCAGGCCGTAGATGCTCCCCGCCGGTC
AGTCACTTAGTTCGTGATCGCCCCCTCTGGTATCACAGTGCTTCTGTTCAAGTTGCACACTGGGCCACAGAGGATCCA

M	K	K	W	S	S	T	D	L	G	T	A	A	D	P	L	Q	K		18	
GCAAGG	ATG	AAG	AAA	TGG	AGC	AGC	ACA	GAC	TTG	GGG	ACA	GCT	GCG	GAC	CCA	CTC	CAA	AAG	54	
D	T	C	P	D	P	L	D	G	D	P	N	S	R	P	P	P	A	K	P	38
3AC	ACC	TGC	CCA	GAC	CCC	CTG	GAT	GGA	GAC	CCT	AAC	TCC	AGG	CCA	CCT	CCA	GCC	AAG	CCC	114
Q	L	P	T	A	K	S	R	T	R	L	F	G	K	G	D	S	E	E	A	58
CAG	CTC	CCC	ACG	GCC	AAG	AGC	CGC	ACC	CGG	CTC	TTT	GGG	AAG	GGT	GAC	TCG	GAG	GAG	GCT	174
F	P	V	D	C	P	H	E	E	G	E	L	D	S	C	P	T	I	T	V	78
ETC	CCG	GTG	GAT	TGC	CCC	CAC	GAG	GAA	GGT	GAG	TTG	GAC	TCC	TGC	CCG	ACC	ATC	ACA	GTC	234
S	P	V	I	T	I	Q	R	P	G	D	G	P	T	G	A	R	L	L	S	98
AGC	CCT	GTT	ATC	ACC	ATC	CAG	AGG	CCA	GGA	GAC	GGC	CCC	ACC	GGT	GCC	AGG	CTG	CTG	TCC	294
Q	D	S	V	A	A	S	T	E	K	T	L	R	L	Y	D	R	R	S	I	118
AG	GAC	TCT	GTC	GCC	GCC	AGC	ACC	GAG	AAG	ACC	CTC	AGG	CTC	TAT	GAT	CGC	AGG	AGT	ATC	354
F	E	A	V	A	Q	N	N	C	Q	D	L	E	S	L	L	L	F	L	Q	138
ET	GAA	GCC	GTT	GCT	CAG	AAT	AAC	TGC	CAG	GAT	CTG	GAG	AGC	CTG	CTG	CTC	TTC	CTG	CAG	414

FIGURE 1

K	S	K	K	H	L	T	D	N	E	F	K	D	P	E	T	G	K	T	C	158
AAG	AGC	AAG	AAG	CAC	CTC	ACA	GAC	AAC	GAG	TTC	AAA	GAC	CCT	GAG	ACA	GGG	AAG	ACC	TGT	474
L	L	K	A	M	L	N	L	H	D	G	Q	N	T	T	I	P	L	L	L	178
CTG	CTG	AAA	GCC	ATG	CTC	AAC	CTG	CAC	GAC	GGA	CAG	AAC	ACC	ACC	ATC	CCC	CTG	CTC	CTG	534
E	I	A	R	Q	T	D	S	L	K	E	L	V	N	A	S	Y	T	D	S	198
BAG	ATC	GCG	CGG	CAA	ACG	GAC	AGC	CTG	AAG	GAG	CTT	GTC	AAC	GCC	AGC	TAC	ACG	GAC	AGC	594
Y	Y	K	G	Q	T	A	L	H	I	A	I	E	R	R	N	M	A	L	V	218
TAC	TAC	AAG	GGC	CAG	ACA	GCA	CTG	CAC	ATC	GCC	ATC	GAG	AGA	CGC	AAC	ATG	GCC	CTG	GTG	654
T	L	L	V	E	N	G	A	D	V	Q	A	A	A	H	G	D	F	F	K	238
ACC	CTC	CTG	GTG	GAG	AAC	GGA	GCA	GAC	GTC	CAG	GCT	GCG	GCC	CAT	GGG	GAC	TTC	TTT	AAG	714
K	T	K	G	R	P	G	F	Y	F	G	E	L	P	L	S	L	A	A	C	258
AAA	ACC	AAA	GGG	CGG	CCT	GGA	TTC	TAC	TTC	GGT	GAA	CTG	CCC	CTG	TCC	CTG	GCC	GCG	TGC	774
T	N	Q	L	G	I	V	K	F	L	L	Q	N	S	W	Q	T	A	D	I	278
ACC	AAC	CAG	CTG	GGC	ATC	GTG	AAG	TTC	CTG	CTG	CAG	AAC	TCC	TGG	CAG	ACG	GCC	GAC	ATC	834
S	A	R	D	S	V	G	N	T	V	L	H	A	L	V	E	V	A	D	N	298
AGC	GCC	AGG	GAC	TCG	GTG	GGC	AAC	ACG	GTG	CTG	CAC	GCC	CTG	GTG	GAG	GTG	GCC	GAC	AAC	894
T	A	D	N	T	K	F	V	T	S	M	Y	N	E	I	L	M	L	G	A	318
ACG	GCC	GAC	AAC	ACG	AAG	TTT	GTG	ACG	AGC	ATG	TAC	AAT	GAG	ATT	CTG	ATG	CTG	GGG	GCC	954
K	L	H	P	T	L	K	L	E	E	L	T	N	K	K	G	M	T	P	L	338
AAA	CTG	CAC	CCG	ACG	CTG	AAG	CTG	GAG	GAG	CTC	ACC	AAC	AAG	AAG	GGA	ATG	ACG	CCG	CTG	1014
A	L	A	A	G	T	G	K	I	G	V	L	A	Y	I	L	Q	R	E	I	358
GCT	CTG	GCA	GCT	GGG	ACC	GGG	AAG	ATC	GGG	GTC	TTG	GCC	TAT	ATT	CTC	CAG	CGG	GAG	ATC	1074
Q	E	P	E	C	R	H	L	S	R	K	F	T	E	W	A	Y	G	P	V	378
CAG	GAG	CCC	GAG	TGC	AGG	CAC	CTG	TCC	AGG	AAG	TTC	ACC	GAG	TGG	GCC	TAC	GGG	CCC	GTG	1134
H	S	S	L	Y	D	L	S	C	I	D	T	C	E	K	N	S	V	L	E	398
CAC	TCC	TCG	CTG	TAC	GAC	CTG	TCC	TGC	ATC	GAC	ACC	TGC	GAG	AAG	AAC	TCG	GTG	CTG	GAG	1194
V	I	A	Y	S	S	S	E	T	P	N	R	H	D	M	L	L	V	E	P	418
GTG	ATC	GCC	TAC	AGC	AGC	AGC	GAG	ACC	CCT	AAT	CGC	CAC	GAC	ATG	CTC	TTG	GTG	GAG	CCG	1254
L	N	R	L	L	Q	D	K	W	D	R	F	V	K	R	I	F	Y	F	N	438
CTG	AAC	CGA	CTC	CTG	CAG	GAC	AAG	TGG	GAC	AGA	TTC	GTC	AAG	CGC	ATC	TTC	TAC	TTC	AAC	1314
F	L	V	Y	C	L	Y	M	I	I	F	T	M	A	A	Y	Y	R	P	V	458
TTC	CTG	GTC	TAC	TGC	CTG	TAC	ATG	ATC	ATC	TTC	ACC	ATG	GCT	GCC	TAC	TAC	AGG	CCC	GTG	1374
D	G	L	P	P	F	K	M	E	K	I	G	D	Y	F	R	V	T	G	E	478
TAT	GGC	TTG	CCT	CCC	TTT	AAG	ATG	GAA	AAA	ATT	GGA	GAC	TAT	TTC	CGA	GTT	ACT	GGA	GAG	1434
I	L	S	V	L	G	G	V	Y	F	F	F	R	G	I	Q	Y	F	L	Q	498
TC	CTG	TCT	GTG	TTA	GGA	GGA	GTC	TAC	TTC	TTT	TTC	CGA	GGG	ATT	CAG	TAT	TTC	CTG	CAG	1494

FIGURE 1 (cont'd)

R	R	P	S	M	K	T	L	F	V	D	S	Y	S	E	M	L	F	F	L	518
GG	CGG	CCG	TCG	ATG	AAG	ACC	CTG	TTT	GTG	GAC	AGC	TAC	AGT	GAG	ATG	CTT	TTC	TTT	CTG	1554
Q	S	L	F	M	L	A	T	V	V	L	Y	F	S	H	L	K	E	Y	V	538
AG	TCA	CTG	TTC	ATG	CTG	GCC	ACC	GTG	GTG	CTG	TAC	TTC	AGC	CAC	CTC	AAG	GAG	TAT	GTG	1614
A	S	M	V	F	S	L	A	L	G	W	T	N	M	L	Y	Y	T	R	G	558
3CT	TCC	ATG	GTA	TTC	TCC	CTG	GCC	TTG	GGC	TGG	ACC	AAC	ATG	CTC	TAC	TAC	ACC	CGC	GGT	1674
F	Q	Q	M	G	I	Y	A	V	M	I	E	K	M	I	L	R	D	L	C	578
CTC	CAG	CAG	ATG	GGC	ATC	TAT	GCC	GTC	ATG	ATA	GAG	AAG	ATG	ATC	CTG	AGA	GAC	CTG	TGC	1734
R	F	M	F	V	Y	I	V	F	L	F	G	F	S	T	A	V	V	T	L	598
3GT	TTC	ATG	TTT	GTC	TAC	ATC	GTC	TTC	TTG	TTC	GGG	TTT	TCC	ACA	GCG	GTG	GTG	ACG	CTG	1794
I	E	D	G	K	N	D	S	L	P	S	E	S	T	S	H	R	W	R	G	618
ATT	GAA	GAC	GGG	AAG	AAT	GAC	TCC	CTG	CCG	TCT	GAG	TCC	ACG	TCG	CAC	AGG	TGG	CGG	GGG	1854
P	A	C	R	P	P	D	S	S	Y	N	S	L	Y	S	T	C	L	E	L	638
3CT	GCC	TGC	AGG	CCC	CCC	GAT	AGC	TCC	TAC	AAC	AGC	CTG	TAC	TCC	ACC	TGC	CTG	GAG	CTG	1914
F	K	F	T	I	G	M	G	D	L	E	F	T	E	N	Y	D	F	K	A	658
TC	AAG	TTC	ACC	ATC	GGC	ATG	GGC	GAC	CTG	GAG	TTC	ACT	GAG	AAC	TAT	GAC	TTC	AAG	GCT	1974
V	F	I	I	L	L	L	A	Y	V	I	L	T	Y	I	L	L	L	N	M	678
3TC	TTC	ATC	ATC	CTG	CTG	CTG	GCC	TAT	GTA	ATT	CTC	ACC	TAC	ATC	CTC	CTG	CTC	AAC	ATG	2034
L	I	A	L	M	G	E	T	V	N	K	I	A	Q	E	S	K	N	I	W	698
TC	ATC	GCC	CTC	ATG	GGT	GAG	ACT	GTC	AAC	AAG	ATC	GCA	CAG	GAG	AGC	AAG	AAC	ATC	TGG	2094
K	L	Q	R	A	I	T	I	L	D	T	E	K	S	F	L	K	C	M	R	718
AG	CTG	CAG	AGA	GCC	ATC	ACC	ATC	CTG	GAC	ACG	GAG	AAG	AGC	TTC	CTT	AAG	TGC	ATG	AGG	2154
K	A	F	R	S	G	K	L	L	Q	V	G	Y	T	P	D	G	K	D	D	738
AG	GCC	TTC	CGC	TCA	GGC	AAG	CTG	CTG	CAG	GTG	GGG	TAC	ACA	CCT	GAT	GGC	AAG	GAC	GAC	2214
Y	R	W	C	F	R	V	D	E	V	N	W	T	T	W	N	T	N	V	G	758
AC	CGG	TGG	TGC	TTC	AGG	GTG	GAC	GAG	GTG	AAC	TGG	ACC	ACC	TGG	AAC	ACC	AAC	GTG	GGC	2274
I	I	N	E	D	P	G	N	C	E	G	V	K	R	T	L	S	F	S	L	778
TC	ATC	AAC	GAA	GAC	CCG	GGC	AAC	TGT	GAG	GGC	GTC	AAG	CGC	ACC	CTG	AGC	TTC	TCC	CTG	2334
R	S	S	R	V	S	G	R	H	W	K	N	F	A	L	V	P	L	L	R	798
GG	TCA	AGC	AGA	GTT	TCA	GGC	AGA	CAC	TGG	AAG	AAC	TTT	GCC	CTG	GTC	CCC	CTT	TTA	AGA	2394
E	A	S	A	R	D	R	Q	S	A	Q	P	E	E	V	Y	L	R	Q	F	818
AG	GCA	AGT	GCT	CGA	GAT	AGG	CAG	TCT	GCT	CAG	CCC	GAG	GAA	GTT	TAT	CTG	CGA	CAG	TTT	2454
S	G	S	L	K	P	E	D	A	E	V	F	K	S	P	A	A	S	G	E	838
CA	GGG	TCT	CTG	AAG	CCA	GAG	GAC	GCT	GAG	GTC	TTC	AAG	AGT	CCT	GCC	GCT	TCC	GGG	GAG	2514
K	*																			840
AG	TGA																			2520

FIGURE 1 (cont'd)

GGACGTCACGCAGACAGCACTGTCAACACTGGGCCTTAGGAGACCCCGTTGCCACGGGGGGCTGCTGAGGGAACACCAG
TGCTCTGTCAGCAGCCTGGCCTGGTCTGTGCCTGCCCAGCATGTTCCCAAATCTGTGCTGGACAAGCTGTGGGAAGCGT
CTTGGAAGCATGGGGAGTGATGTACATCCAACCGTCACTGTCCCCAAGTGAATCTCCTAACAGACTTTCAGGTTTTTA
CTCACTTTACTAAAAAAAAAAAAAAAAAGGGCGGCCGCTTA

FIGURE 1 (cont'd)

Full-length human VR2

Input file Flh21e11.seq; Output File Flh21e11.tra
Sequence length 2809

```

GGCTAGCCTGTCTGACAGGGGAGAGTTAAGCTCCCGTTCTCCACCGTGC CGGCTGGCCAGGTGGGCTGAGGGTGACCG
AGAGACCAGAACTGCTGTGCTGGAGCTTAGTGCTCAGAGCTGGGGAGGGAGGTTCCGCCGCTCCTCTGCTGTCTAGCGCC
GGCAGCCCCCTCCCGCTTCACTTCTCCCGCAGCCCCCTGCTACTGAGAAGCTCCGGGATCCCAGCAGCCGCCACGCCCT
GGCCTCAGCCTGCGGGGCTCCAGTCAGGCCAACACCGACGCGCAGCTGGGAGGAAGACAGGACCCTTGACATCTCCATC

                                     M   T   S   P   S   S   S   P
TGCACAGAGGTCCTGGCTGGACGAGCAGCCTCTCTCTCTAGG ATG ACC TCA CCC TCC AGC TCT CCA      8
                                                                24

V   F   R   L   E   T   L   D   G   G   Q   E   D   G   S   E   A   D   R   G      28
GTT TTC AGG TTG GAG ACA TTA GAT GGA GGC CAA GAA GAT GGC TCT GAG GCG GAC AGA GGA      84

K   L   D   F   G   S   G   L   P   P   M   E   S   Q   F   Q   G   E   D   R      48
AAG CTG GAT TTT GGG AGC GGG CTG CCT CCC ATG GAG TCA CAG TTC CAG GGC GAG GAC CGG      144

K   F   A   P   Q   I   R   V   N   L   N   Y   R   K   G   T   G   A   S   Q      68
AAA TTC GCC CCT CAG ATA AGA GTC AAC CTC AAC TAC CGA AAG GGA ACA GGT GCC AGT CAG      204

P   D   P   N   R   F   D   R   D   R   L   F   N   A   V   S   R   G   V   P      88
CCG GAT CCA AAC CGA TTT GAC CGA GAT CGG CTC TTC AAT GCG GTC TCC CGG GGT GTC CCC      264

E   D   L   A   G   L   P   E   Y   L   S   K   T   S   K   Y   L   T   D   S      108
GAG GAT CTG GCT GGA CTT CCA GAG TAC CTG AGC AAG ACC AGC AAG TAC CTC ACC GAC TCG      324

E   Y   T   E   G   S   T   G   K   T   C   L   M   K   A   V   L   N   L   K      128
GAA TAC ACA GAG GGC TCC ACA GGT AAG ACG TGC CTG ATG AAG GCT GTG CTG AAC CTT AAG      384

D   G   V   N   A   C   I   L   P   L   L   Q   I   D   R   D   S   G   N   P      148
GAC GGA GTC AAT GCC TGC ATT CTG CCA CTG CTG CAG ATC GAC AGG GAC TCT GGC AAT CCT      444

Q   P   L   V   N   A   Q   C   T   D   D   Y   Y   R   G   H   S   A   L   H      168
CAG CCC CTG GTA AAT GCC CAG TGC ACA GAT GAC TAT TAC CGA GGC CAC AGC GCT CTG CAC      504

I   A   I   E   K   R   S   L   Q   C   V   K   L   L   V   E   N   G   A   N      188
ATC GCC ATT GAG AAG AGG AGT CTG CAG TGT GTG AAG CTC CTG GTG GAG AAT GGG GCC AAT      564

V   H   A   R   A   C   G   R   F   F   Q   K   G   Q   G   T   C   F   Y   F      208
GTG CAT GCC CGG GCC TGC GGC CGC TTC TTC CAG AAG GGC CAA GGG ACT TGC TTT TAT TTC      624

G   E   L   P   L   S   L   A   A   C   T   K   Q   W   D   V   V   S   Y   L      228
GGT GAG CTA CCC CTC TCT TTG GCC GCT TGC ACC AAG CAG TGG GAT GTG GTA AGC TAC CTC      684

L   E   N   P   H   Q   P   A   S   L   Q   A   T   D   S   Q   G   N   T   V      248
CTG GAG AAC CCA CAC CAG CCC GCC AGC CTG CAG GCC ACT GAC TCC CAG GGC AAC ACA GTC      744

L   H   A   L   V   M   I   S   D   N   S   A   E   N   I   A   L   V   T   S      268
CTG CAT GCC CTA GTG ATG ATC TCG GAC AAC TCA GCT GAG AAC ATT GCA CTG GTG ACC AGC      804

M   Y   D   G   L   L   Q   A   G   A   R   L   C   P   T   V   Q   L   E   D      288
ATG TAT GAT GGG CTC CTC CAA GCT GGG GCC CGC CTC TGC CCT ACC GTG CAG CTT GAG GAC      864

I   R   N   L   Q   D   L   T   P   L   K   L   A   A   K   E   G   K   I   E      308

```

FIGURE 2

ATC	CGC	AAC	CTG	CAG	GAT	CTC	ACG	CCT	CTG	AAG	CTG	GCC	GCC	AAG	GAG	GGC	AAG	ATC	GAG	924
I	F	R	H	I	L	Q	R	E	F	S	G	L	S	H	L	S	R	K	F	328
ATT	TTC	AGG	CAC	ATC	CTG	CAG	CGG	GAG	TTT	TCA	GGA	CTG	AGC	CAC	CTT	TCC	CGA	AAG	TTC	984
T	E	W	C	Y	G	P	V	R	V	S	L	Y	D	L	A	S	V	D	S	348
ACC	GAG	TGG	TGC	TAT	GGG	CCT	GTC	CGG	GTG	TGG	CTG	TAT	GAC	CTG	GCT	TCT	GTG	GAC	AGC	1044
C	E	E	N	S	V	L	E	I	I	A	F	H	C	K	S	P	H	R	H	368
TGT	GAG	GAG	AAC	TCA	GTG	CTG	GAG	ATC	ATT	GCC	TTT	CAT	TGC	AAG	AGC	CCG	CAC	CGA	CAC	1104
R	M	V	V	L	E	P	L	N	K	L	L	Q	A	K	W	D	L	L	I	388
CGA	ATG	GTC	GTT	TTG	GAG	CCC	CTG	AAC	AAA	CTG	CTG	CAG	GCG	AAA	TGG	GAT	CTG	CTC	ATC	1164
P	K	F	F	L	N	F	L	C	N	L	I	Y	M	F	I	F	T	A	V	408
CCC	AAG	TTC	TTC	TTA	AAC	TTC	CTG	TGT	AAT	CTG	ATC	TAC	ATG	TTC	ATC	TTC	ACC	GCT	GTT	1224
A	Y	H	Q	P	T	L	K	K	Q	A	A	P	H	L	K	A	E	V	G	428
GCC	TAC	CAT	CAG	CCT	ACC	CTG	AAG	AAG	CAG	GCC	GCC	CCT	CAC	CTG	AAA	GCG	GAG	GTT	GGA	1284
N	S	M	L	L	T	G	H	I	L	I	L	L	G	G	I	Y	L	L	V	448
AAC	TCC	ATG	CTG	CTG	ACG	GGC	CAC	ATC	CTT	ATC	CTG	CTA	GGG	GGG	ATC	TAC	CTC	CTC	GTG	1344
G	Q	L	W	Y	F	W	R	R	H	V	F	I	W	I	S	F	I	D	S	468
GGC	CAG	CTG	TGG	TAC	TTC	TGG	CGG	CGC	CAC	GTG	TTC	ATC	TGG	ATC	TCG	TTC	ATA	GAC	AGC	1404
Y	F	E	I	L	F	L	F	Q	A	L	L	T	V	V	S	Q	V	L	C	488
TAC	TTT	GAA	ATC	CTC	TTC	CTG	TTC	CAG	GCC	CTG	CTC	ACA	GTG	GTG	TCC	CAG	GTG	CTG	TGT	1464
F	L	A	I	E	W	Y	L	P	L	L	V	S	A	L	V	L	G	W	L	508
TTC	CTG	GCC	ATC	GAG	TGG	TAC	CTG	CCC	CTG	CTT	GTG	TCT	GCG	CTG	GTG	CTG	GGC	TGG	CTG	1524
N	L	L	Y	Y	T	R	G	F	Q	H	T	G	I	Y	S	V	M	I	Q	528
AAC	CTG	CTT	TAC	TAT	ACA	CGT	GGC	TTC	CAG	CAC	ACA	GGC	ATC	TAC	AGT	GTC	ATG	ATC	CAG	1584
K	V	I	L	R	D	L	L	R	F	L	L	I	Y	L	V	F	L	F	G	548
AAG	GTC	ATC	CTG	CGG	GAC	CTG	CTG	CGC	TTC	CTT	CTG	ATC	TAC	TTA	GTC	TTC	CTT	TTC	GGC	1644
F	A	V	A	L	V	S	L	S	Q	E	A	W	R	P	E	A	P	T	G	568
TTC	GCT	GTA	GCC	CTG	GTG	AGC	CTG	AGC	CAG	GAG	GCT	TGG	CGC	CCC	GAA	GCT	CCT	ACA	GGC	1704
P	N	A	T	E	S	V	Q	P	M	E	G	Q	E	D	E	G	N	G	A	588
CCC	AAT	GCC	ACA	GAG	TCA	GTG	CAG	CCC	ATG	GAG	GGA	CAG	GAG	GAC	GAG	GGC	AAC	GGG	GCC	1764
Q	Y	R	G	I	L	E	A	S	L	E	L	F	K	F	T	I	G	M	G	608
CAG	TAC	AGG	GGT	ATC	CTG	GAA	GCC	TCC	TTG	GAG	CTC	TTC	AAA	TTC	ACC	ATC	GGC	ATG	GGC	1824
E	L	A	F	Q	E	Q	L	H	F	R	G	M	V	L	L	L	L	L	A	628
GAG	CTG	GCC	TTC	CAG	GAG	CAG	CTG	CAC	TTC	CGC	GGC	ATG	GTG	CTG	CTG	CTG	CTG	CTG	GCC	1884
Y	V	L	L	T	Y	I	L	L	L	N	M	L	I	A	L	M	S	E	T	648
TAC	GTG	CTG	CTC	ACC	TAC	ATC	CTG	CTG	CTC	AAC	ATG	CTC	ATC	GCC	CTC	ATG	AGC	GAG	ACC	1944
V	N	S	V	A	T	D	S	W	S	I	W	K	L	Q	K	A	I	S	V	668
GTC	AAC	AGT	GTC	GCC	ACT	GAC	AGC	TGG	AGC	ATC	TGG	AAG	CTG	CAG	AAA	GCC	ATC	TCT	GTC	2004
L	E	M	E	N	G	Y	W	W	C	R	K	K	Q	R	A	G	V	M	L	688
CTG	GAG	ATG	GAG	AAT	GGC	TAT	TGG	TGG	TGC	AGG	AAG	AAG	CAG	CGG	GCA	GGT	GTG	ATG	CTG	2064
T	V	G	T	K	P	D	G	S	P	D	E	R	W	C	F	R	V	E	E	708

FIGURE 2 (continued)

ACC GTT GGC ACT AAG CCA GAT GGC AGC CCG GAT GAG CGC TGG TGC TTC AGG GTG GAG GAG	2124
V N W A S W E Q T L P T L C E D P S G A	728
GTG AAC TGG GCT TCA TGG GAG CAG ACG CTG CCT ACG CTG TGT GAG GAC CCG TCA GGG GCA	2184
G V P R T L E N P V L A S P P K E D E D	748
GGT GTC CCT CGA ACT CTC GAG AAC CCT GTC CTG GCT TCC CCT CCC AAG GAG GAT GAG GAT	2244
G A S E E N Y V P V Q L L Q S N *	765
GGT GCC TCT GAG GAA AAC TAT GTG CCC GTC CAG CTC CTC CAG TCC AAC TGA	2295

TGGCCCAGATGCAGCAGGAGGCCAGAGGACAGAGCAGAGGATCTTTCCAACCACATCTGCTGGCTCTGGGGTCCAGTG

AATTCTGGTGGCAAATATATATATTTTCACTAACTCAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAA

FIGURE 2 (continued)

Partial human VR2 alternate form

Input file frhob12c4.seq; Output File frhob12c4.tra
Sequence length 1489

G R F F Q K G Q G T C F Y F G E L' P L	19
GC GGC CGC TTC TTC CAG AAG GGC CAA GGG ACT TGC TTT TAT TTC GGT GAG CTA CCC CTC	57
S L A A C T K Q W D V V S Y L L E N P H	39
TCT TTG GCC GCT TGC ACC AAG CAG TGG GAT GTG GTA AGC TAC CTC CTG GAG AAC CCA CAC	117
Q P A S L Q A T D S Q G N T V L H A L V	59
CAG CCC GCC AGC CTG CAG GCC ACT GAC TCC CAG GGC AAC ACA GTC CTG CAT GCC CTA GTG	177
M I S D N S A E N I A L V T S M Y D G L	79
ATG ATC TCG GAC AAC TCA GCT GAG AAC ATT GCA CTG GTG ACC AGC ATG TAT GAT GGG CTC	237
L Q A G A R L C P T V Q L E D I R N L Q	99
CTC CAA GCT GGG GCC CGC CTC TGC CCT ACC GTG CAG CTT GAG GAC ATC CGC AAC CTG CAG	297
D L T P L K L A A K E G K I E I F R H I	119
GAT CTC ACG CCT CTG AAG CTG GCC GCC AAG GAG GGC AAG ATC GAG ATT TTC AGG CAC ATC	357
L Q R E F S G L S H L S R K F T E W C Y	139
CTG CAG CGG GAG TTT TCA GGA CTG AGC CAC CTT TCC CGA AAG TTC ACC GAG TGG TGC TAT	417
G P V R V S L Y D L A S V D S C E E N S	159
GGG CCT GTC CGG GTG TCG CTG TAT GAC CTG GCT TCT GTG GAC AGC TGT GAG GAG AAC TCA	477
V L E I I A F H C K S P H R H R M V V L	179
GTG CTG GAG ATC ATT GCC TTT CAT TGC AAG AGC CCG CAC CGA CAC CGA ATG GTC GTT TTG	537
E P L N K L L Q A K W D L L I P K F F L	199
GAG CCC CTG AAC AAA CTG CTG CAG GCG AAA TGG GAT CTG CTC ATC CCC AAG TTC TTC TTA	597
N F L C N L I Y M F I F T A V A Y H Q P	219
AAC TTC CTG TGT AAT CTG ATC TAC ATG TTC ATC TTC ACC GCT GTT GCC TAC CAT CAG CCT	657
T L K K Q A A P H L K A E V G N S M L L	239
ACC CTG AAG AAG CAG GCC GCC CCT CAC CTG AAA GCG GAG GTT GGA AAC TCC ATG CTG CTG	717
T G H I L I L L G G I Y L L V G Q L W Y	259
CAG GGC CAC ATC CTT ATC CTG CTA GGG GGG ATC TAC CTC CTC GTG GGC CAG CTG TGG TAC	777
F W R R H V F I W I S F I D S Y F E I L	279
TTC TGG CGG CGC CAC GTG TTC ATC TGG ATC TCG TTC ATA GAC AGC TAC TTT GAA ATC CTC	837
F L F Q A L L T V V S Q V L C F L A I E	299
TTC CTG TTC CAG GCC CTG CTC ACA GTG GTG TCC CAG GTG CTG TGT TTC CTG GCC ATC GAG	897
W Y L P L L V S A L V L G W L N L L Y Y	319
TGG TAC CTG CCC CTG CTT GTG TCT GCG CTG GTG CTG GGC TGG CTG AAC CTG CTT TAC TAT	957
T R G F Q H T G I Y S V M I Q K K A I S	339
ACA CGT GGC TTC CAG CAC ACA GGC ATC TAC AGT GTC ATG ATC CAG AAG AAA GCC ATC TCT	1017
V L E M E N G Y W W C R K K Q R A G V M	359
GTC CTG GAG ATG GAG AAT GGC TAT TGG TGG TGC AGG AAG AAG CAG CGG GCA GGT GTG ATG	1077

FIGURE 3

L	T	V	G	T	K	P	D	G	S	P	D	E	R	W	C	F	R	V	E	379
CTG	ACC	GTT	GGC	ACT	AAG	CCA	GAT	GGC	AGC	CCG	GAT	GAG	CGC	TGG	TGC	TTC	AGG	GTG	GAG	1137
E	V	N	W	A	S	W	E	Q	T	L	P	T	L	C	E	D	P	S	G	399
GAG	GTG	AAC	TGG	GCT	TCA	TGG	GAG	CAG	ACG	CTG	CCT	ACG	CTG	TGT	GAG	GAC	CCG	TCA	GGG	1197
A	G	V	P	R	T	L	E	N	P	V	L	A	S	P	P	K	E	D	E	419
GCA	GGT	GTC	CCT	CGA	ACT	CTC	GAG	AAC	CCT	GTC	CTG	GCT	TCC	CCT	CCC	AAG	GAG	GAT	GAG	1257
D	G	A	S	E	E	N	Y	V	P	V	Q	L	L	Q	S	N	*			437
GAT	GGT	GCC	TCT	GAG	GAA	AAC	TAT	GTG	CCC	GTC	CAG	CTC	CTC	CAG	TCC	AAC	TGA			1311
TGGCCCAGATGCAGCAGGAGGCCAGAGGACAGAGCAGAGGATCTTTCCAACCACATCTGCTGGCTCTGGGGTCCCAAGTG																				
AATTCTGGTGGCAAATATATATATTTTCACTAACTCAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAGGA																				
GCGGACGCGTGGGTGAC																				

FIGURE 3 (continued)

Partial rat VR2

Input file Flrxb147g11.seq; Output File Flrxb147g11.tra
Sequence length 1794

S T H A S A L S L A A C T K Q W D V V	19
G TCG ACC CAC GCG TCC GCT CTT TCT CTG GCT GCG TGC ACC AAG CAG TGG GAT GTG GTG	57
T Y L L E N P H Q P A S L E A T D S L G	39
ACC TAC CTC CTG GAG AAC CCA CAC CAG CCG GCC AGC CTG GAG GCC ACC GAC TCC CTG GGC	117
N T V L H A L V M I A D N S P E N S A L	59
AAC ACA GTC CTG CAT GCT CTG GTA ATG ATT GCA GAT AAC TCG CCT GAG AAC AGT GCC CTG	177
V I H M Y D G L L Q M G A R L C P T V Q	79
GTG ATC CAC ATG TAC GAC GGG CTT CTA CAA ATG GGG GCG CGC CTC TGC CCC ACT GTG CAG	237
L E E I S N H Q G L T P L K L A A K E G	99
CTT GAG GAA ATC TCC AAC CAC CAA GGC CTC ACA CCC CTG AAA CTA GCC GCC AAG GAA GGC	297
K I E I F R H I L Q R E F S G P Y Q P L	119
AAA ATC GAG ATT TTC AGG CAC ATT CTG CAG CGG GAA TTC TCA GGA CCG TAC CAG CCC CTT	357
S R K F T E W C Y G P V R V S L Y D L S	139
TCC CGA AAG TTT ACT GAG TGG TGT TAC GGT CCT GTG CGG GTA TCG CTG TAC_GAC CTG TCC	417
S V D S W E K N S V L E I I A F H C K S	159
TCT GTG GAC AGC TGG GAA AAG AAC TCG GTG CTG GAG ATC ATC GCT TTT CAT TGC AAG AGC	477
P N R H R M V V L E P L N K L L Q E K W	179
CGG AAC CGG CAC CGC ATG GTG GTT TTA GAA CCA CTG AAC AAG CTT CTG CAG GAG AAA TGG	537
D R L V S R F F F N F A C Y L V Y M F I	199
GAT CGG CTC GTC TCA AGA TTC TTC TTC AAC TTC GCC TGC TAC TTG GTC TAC ATG TTC ATC	597
F T V V A Y H Q P S L D Q P A I P S S K	219
TTC ACC GTC GTT GCC TAC CAC CAG CCT TCC CTG GAT CAG CCA GCC ATC CCC TCA TCA AAA	657
A T F G E S M L L L G H I L I L L G G I	239
GCG ACT TTT GGG GAA TCC ATG CTG CTG CTG GGC CAC ATT CTG ATC CTG CTT GGG GGT ATT	717
Y L L L G Q L W Y F W R R R L F I W I S	259
TAC CTC TTA CTG GGC CAG CTG TGG TAC TTT TGG CGG CGG CGC CTG TTT ATC TGG ATC TCA	777
F M D S Y F E I L F L L Q A L L T V L S	279
TTC ATG GAC AGC TAC TTT GAA ATC CTC TTT CTC CTT CAG GCT CTG CTC ACA GTG CTG TCC	837
Q V L R F M E T E W Y L P L L V L S L V	299
CAG GTG CTG CGC TTC ATG GAG ACT GAA TGG TAC CTA CCC CTG CTA GTG TTA TCC CTA GTG	897
L G W L N L L Y Y T R G F Q H T G I Y S	319
CTG GGC TGG CTG AAC CTG CTT TAC TAC ACA CGG GGC TTT CAG CAC ACA GGC ATC TAC AGT	957
V M I Q K V I L R D L L R F L L V Y L V	339
GTC ATG ATC CAG AAG GTC ATC CTT CGA GAC CTG CTC CGT TTC CTG CTG GTC TAC CTG GTC	1017
F L F G F A V A L V S L S R E A R S P K	359
TTC CTT TTC GGC TTT GCT GTA GCC CTA GTA AGC TTG AGC AGA GAG GCC CGA AGT CCC AAA	1077

FIGURE 4

A	P	E	D	N	N	S	T	V	T	E	Q	P	T	V	G	Q	E	E	E	379
GCC	CCT	GAA	GAT	AAC	AAC	TCC	ACA	GTG	ACG	GAA	CAG	CCC	ACG	GTG	GGC	CAG	GAG	GAG	GAG	1137
P	A	P	Y	R	S	I	L	D	A	S	L	E	L	F	K	F	T	I	G	399
CCA	GCT	CCA	TAT	CGG	AGC	ATT	CTG	GAT	GCC	TCC	CTA	GAG	CTG	TTC	AAG	TTC	ACC	ATT	GGT	1197
M	G	E	L	A	F	Q	E	Q	L	R	F	R	G	V	V	L	L	L	L	419
ATG	GGG	GAG	CTG	GCT	TTC	CAG	GAA	CAG	CTG	CGT	TTT	CGT	GGG	GTG	GTC	CTG	CTG	TTG	CTG	1257
L	A	Y	V	L	L	T	Y	V	L	L	L	N	M	L	I	A	L	M	S	439
TTG	GCC	TAC	GTC	CTT	CTC	ACC	TAC	GTC	CTG	CTG	CTC	AAC	ATG	CTC	ATT	GCT	CTC	ATG	AGC	1317
E	T	V	N	H	V	A	D	N	S	W	S	I	W	K	L	Q	K	A	I	459
GAA	ACT	GTC	AAC	CAC	GTT	GCT	GAC	AAC	AGC	TGG	AGC	ATC	TGG	AAG	TTG	CAG	AAA	GCC	ATC	1377
S	V	L	E	M	E	N	G	Y	W	W	C	R	R	K	K	H	R	E	G	479
TCT	GTC	TTG	GAG	ATG	GAG	AAT	GGT	TAC	TGG	TGG	TGC	CGG	AGG	AAG	AAA	CAT	CGT	GAA	GGG	1437
R	L	L	K	V	G	T	R	G	D	G	T	P	D	E	R	W	C	F	R	499
AGG	CTG	CTG	AAA	GTC	GGC	ACC	AGG	GGG	GAT	GGT	ACC	CCT	GAT	GAG	CGC	TGG	TGC	TTC	AGG	1497
V	E	E	V	N	W	A	A	W	E	K	T	L	P	T	L	S	E	D	P	519
GTG	GAG	GAA	GTA	AAT	TGG	GCT	GCT	TGG	GAG	AAG	ACT	CTT	CCC	ACC	TTA	TCT	GAG	GAT	CCA	1557
S	G	P	G	I	T	G	N	K	K	N	P	T	S	K	P	G	K	N	S	539
TCA	GGG	CCA	GGC	ATC	ACT	GGT	AAT	AAA	AAG	AAC	CCA	ACC	TCT	AAA	CCG	GGG	AAG	AAC	AGT	1617
A	S	E	E	D	H	L	P	L	Q	V	L	Q	S	P	*					555
GCC	TCA	GAG	GAA	GAC	CAT	CTG	CCC	CTT	CAG	GTC	CTC	CAG	TCC	CCC	TGA					1665

TGGCCCAGATGCAGCAGCAGGCTGGCAGGATGGAGTAGGGAATCTTCCCAGCCACACCAGAGGCTACTGAATTTTGGTG

GAAATATAAATATTTTTTTTTTGCATAAAAAAAAAAAAAAAGGGCGGCCGC

FIGURE 4 (continued)

GAP of: humanvr2.pep check: 5746 from: 1 to: 764

humanVR2 Flh21e11

to: humanvr1.pep check: 6877 from: 1 to: 839

humanVR1 _Fbh18547pat - fchrb87a6, 3909 bases, 4554 checksum.

Symbol comparison table:

/ddm_local/gcg/gcg_9.1/gcgcore/data/rundata/blosum62.cmp

CompCheck: 6430

Gap Weight: 12 Average Match: 2.912
Length Weight: 4 Average Mismatch: -2.003

Quality: 1530 Length: 850
Ratio: 2.003 Gaps: 10
Percent Similarity: 55.378 Percent Identity: 46.348

Match display thresholds for the alignment(s):

| = IDENTITY
: = 2
. = 1

humanvr2.pep x humanvr1.pep

```
1 .....MTSPSSSPVF 10
1 MKKWSSTDLGTAADPLQKDTCPDPLDGD PNSRPPPAKPQLPTAKSRTRLF 50
11 RLETLDGGQEDGSEADRGKLD FGSGLPPMESQFQGEDRKFA PQIRVNLNY 60
51 GKG DSEEA FPVDCPHEEGELDSCPTI.TVSPVITIQRPGDGPTGARLLSQ 99
61 RKGTGASQPD PNRFRDRRLFNAVSRGVPEDLAGLPEYLSKTSKYLTDSEY 110
100 DSVAASTEKTLRLYDRRSIFEAVAQNNCQDLESLLLFLQSKKKHLTDNEF 149
111 TEGSTGKTCLMKAVLNLKDGVNACILPLLQIDRDSGNPQPLVNAQCTDDY 160
150 KDPETGKTCLLKAMLNLDGQNTTIPLLLEIARQTD SLKELVNASYTDSY 199
161 YRGHSALHIAIEKRS LQCVKLLVENGANVHARACGRFFQKGQG.TCFYFG 209
200 YKGQTALHIAIERNMALVTLLVENGADVQAAAHGDFFKTKGRPGFYFG 249
210 ELPLSLAACTKQWDVVS YLLENPHQPASLQATDSQGNTVLHALVMISD NS 259
250 ELPLSLAACTNQLGIVKFLQNSWQTADISARDSVGNTVLHALVEVADNT 299
260 AENIALVTSMYDGLLQAGARLCPTVQLEDIRNLQDLT PLKLAAKEGKIEI 309
300 ADNTKFVTSMYNEILMLGAKLHPTLKLEELTNKKGMTPLALAAAGTGKIGV 349
```

FIGURE 5

310 FRHILQREFS..GLSHLSRKFTWCYGPVRVSLYDLASVDSCEENSVLEI 357
:||||| ||||| ||||| ||||| :|.||.|||||:
350 LAYILQREIQEPECRHLRSRKFTEWAYGVPVHSSLYDLSCIDTCEKNSVLEV 399
358 IAF.HCKSPHRHRMVVLEPLNKLLOAKWDLIPK.FFLNFLCNLIYMFIF 405
||: ..|.|| |...|||:|||| ||| : : |: ||| :|| ||
400 IAYSSSETPNRHDMLLVEPLNRLLODKWDRFVKRIFYFNFLVYCLYMIIF 449
406 TAVAYHQPTLKKQAAPHLKAEVGNSMLLTGHILILLGGIYLLVGQLWYFW 455
| ||:| | :|. .|| || .|||:| : ||
450 TMAAYYRPV..DGLPPFKMEKIGDYFRVTGEILSVLGGVYFFFRGIQYFL 497
456 RRHVFIIWISFIDSYEILFLFQALLTVVSQVLCFLAIEWYLPLLVSALVL 505
.| . |:|| |.|| |. . || | :. |. :| .| |
498 QRRPSMKTLEFVDSYSEMLFFLQSLFMLATVVLYFSHLKEYVASMVFSLAL 547
506 GWLNLLYYTRGFQHTGIYSVMIQKVILRDLLRFLLIYLVFLFGFAVALVS 555
|| |:||||||| |||.|||:|.||||| ||: |:|||||. |. |.
548 GWTNMLYYTRGFQQMGIIYAVMIEKMILRDLCRFMFVYIVFLFGFSTAVVT 597
556 LSQEAWRPEAPTGPNATESVQPMEGQEDEGNGAQYRGILEASLELFKFTL 605
| :: |. . | | . | : |||||
598 LIEDGKNDSLPESTSHRWRGPACRPPD....SSYNSLYSTCLELFKFTI 643
606 GMGELAFQEQLHFRGMVLLLLLAYVLLTYILLNMLIALMSETVNSVATD 655
|||:| | | |: . ::|||:||||| |||| :| :
644 GMDLEFTENYDFKAVFIILLLAYVILTYILLNMLIALMGETVNKIAQE 693
656 SWSIWKLQKAISVLEMENGYWWC.RKKQRAGVMLTVGTPDGSPDERWCF 704
|.|||||:||.::| : | || |. | :| || ||| | ||||
694 SKNIWKLQRAITILDEKSFLLKCMRKAFRSGKLLQVGYPDGKDDYRWCF 743
705 RVEEVNWSWEQTLPTLCEDPSGA.GVPRTLNPVLASPPKEDEDGASEE 753
||:|||| |. . : ||| || ||| . . | | .
744 RVDEVNWTWNTNVGIINEDPGNCEGVKRTLSFSLRSS....RVSGRHWK 789
754 NYVPVQLQSN..... 764
|: | ||.
790 NFALVPLLREASARDRQSAQPEEVYLRQFSGSLKPEDAIEVFKSPAASGEK 839

FIGURE 5 (cont'd)

GAP of: humanvr2.seq check: 8853 from: 1 to: 2809

humanVR2 21e11a, 2809 bases, 8853 checksum.

to: humanvr1.seq check: 4554 from: 1 to: 3909

humanVR1 Fbh18547pat - Import - complete

Symbol comparison table:

/ddm_local/gcg/gcg_9.1/gcgcore/data/rundata/nwsgapdna.cmp

CompCheck: 8760

Gap Weight: 50 Average Match: 10.000
Length Weight: 3 Average Mismatch: 0.000

Quality: 14359 Length: 3934
Ratio: 5.112 Gaps: 15
Percent Similarity: 55.316 Percent Identity: 55.316

Match display thresholds for the alignment(s):

| = IDENTITY
: = 5
. = 1

humanvr2.seq x humanvr1.seq

```
1 .....GGCTAGCCTGTCCTGACAGGGGAGAG 26
      | | | | | | | |
801 TGTCCACAGTAGTCCCCCCTTATCCACGGGTGTCACTTTCCATGGGGTTCA 850
      .
27 TTAAGCTCCCGTTCTCCACCGTGCCGGCTGGCCAGGTGGGCTGAGGGTGA 76
      | | | | | | | | | | | | | | | |
851 GTTATTTGCGGTCAACCACGGTCTGCCAATATTAAATGGAAAATTCTTCA 900
      .
77 CCGAGAGACCAGAACCTGCTTGCTGGAGCTTAGTGCTCAGAGCTGGGGAG 126
      | | | | | | | | | | | | | | | |
901 AACAGTTCCCAAGTTTTCCCTTGTCATTGTTCTGAGCAGTGTGATGAAG 950
      .
127 GGAGGTTCCGCCGCTCCTCTGCTGTCAGCGCCGGCAGCCCCTCCCGGCTT 176
      | | | | | | | | | | | | | | | |
951 AGTCTCTGCCGTGCCATCTGGGATGCAAACCGTCCCTGTGTCCCCCACGT 1000
      .
177 CACTTCCTCCCGCAGCCCCTGCTACTGAGAAGCTCCGGGATCCCAGCAGC 226
      | | | | | | | | | | | | | | | |
1001 CCAGGCCGTAGATGCTCCCCGCCGGTCAGTCACTTAGTCGTCAGATCGCC 1050
      .
227 CGCCACGCCCTGGC.....CTCAGCCTGCGGG 253
      | | | | | | | | | | | | | | | |
1051 CGTCCTGGTATCACAGTGCTTCTGTTTCAGGTTGCACACTGGGCCACAGAG 1100
```

FIGURE 6

254 GCTCCAGTCAGGCCAACACCGACGCGCAGCTGGGAGGAAG..... 293
 | | | | | | | | | | | | | | | | | | | | | |
 1101 GATCCAGCAAGGATGAAGAAATGGAGCAGCACAGACTTGGGGACAGCTGC 1150
 294ACAGGACCCTTGACATCTCCATCTGCACAGAGGTCCTG 331
 | | | | | | | | | | | | | | | | | | | | | |
 1151 GGACCCACTCCAAAAGGACACCTGCCCAGACCCCCTGGATGGAGACCCTA 1200
 332 GCTGGACCGAGCAGCCTCCTCCTCCTAGGATGACCTCACCTCCAGC..T 379
 | | | | | | | | | | | | | | | | | | | | | |
 1201 ACTCCAGGCCACCTCCAGCCAAGCCCCAGCTCCCCACGGCCAAGAGCCGC 1250
 380 CTCCAGTTTTTCAGGTTGGAGACATTAGATGGAGGCCAAGAAGATGGCTCT 429
 | | | | | | | | | | | | | | | | | | | | | |
 1251 ACCCGGCTCTTTGGGAAGGGTGACTCGGAGGAGGCTTTCCCGGTGGATTG 1300
 430 GAGGCGGACAGAGGAAAAGCTGGATTTTGGGAGCGGGCTGCCTCCCATGGA 479
 | | | | | | | | | | | | | | | | | | | | | |
 1301 CCCCCACGAGGAAGGTGAGTTGGACTCCTGCCCGACCATCACAGTCAGCC 1350
 480 GTCACAGTTCCAGGGCGAGGACCGGAAATTCGCCCTCAGATAAGAGTCA 529
 | | | | | | | | | | | | | | | | | | | | | |
 1351 CTGTTATCACCATCCAGAGGCCAGGAGACGGCCCCACCGGTGCCAGG..G 1398
 530 ACCTCAACTACCGAAAGGGAACAGGTGCCAGTCAGCCGGATCCAAACCGA 579
 | | | | | | | | | | | | | | | | | | | | | |
 1399 TGCTGTCCCAGGACTCTGTGCGCCGCAGCACCAGAGAAGACCCTCAGGCTC 1448
 580 TTTGACCGAGATCGGCTCTTCAATGCGGTCTCCCGGGGTGTCCCCGAGGA 629
 | | | | | | | | | | | | | | | | | | | | | |
 1449 TATGATCGCAGGAGTATCTTTGAAGCCGTGCTCAGAATAACTGCCAGGA 1498
 630 TCTGGCTGGACTTCCAGAGTACCTGAGCAAGACCAGCAAGTACCTCACCG 679
 | | | | | | | | | | | | | | | | | | | | | |
 1499 TCTGGAGAGCCTGCTGCTCTTCTGCAGAAGAGCAAGAAGCACCTCACAG 1548
 680 ACTCGGAATACACAGAGGGCTCCACAGGTAAGACGTGCCTGATGAAGGCT 729
 | | | | | | | | | | | | | | | | | | | | | |
 1549 ACAACGAGTTCAAAGACCCTGAGACAGGGAAGACCTGTCTGCTGAAAGCC 1598
 730 GTGCTGAACCTTAAGGACGGAGTCAATGCCTGCATTCTGCCACTGCTGCA 779
 | | | | | | | | | | | | | | | | | | | | | |
 1599 ATGCTCAACCTGCACGACGGACAGAACACCACCATCCCCCTGCTCCTGGA 1648
 780 GATCGACAGGGACTCTGGCAATCCTCAGCCCCCTGGTAAATGCCCAGTGCA 829
 | | | | | | | | | | | | | | | | | | | | | |
 1649 GATCGCGCGGCAAACGGACAGCCTGAAGGAGCTTGTCACGCCAGCTACA 1698
 830 CAGATGACTATTACCGAGGCCACAGCGCTCTGCACATCGCCATTGAGAAG 879
 | | | | | | | | | | | | | | | | | | | | | |
 1699 CGGACAGCTACTACAAGGGCCAGACAGCACTGCACATCGCCATCGAGAGA 1748
 880 AGGAGTCTGCAGTGTGTGAAGCTCCTGGTGGAGAATGGGGCCAATGTGCA 929
 | | | | | | | | | | | | | | | | | | | | | |
 1749 CGCAACATGGCCCTGGTGACCCTCCTGGTGGAGAACGGAGCAGACGTCCA 1798
 930 TGCCCGGGCCTGCGGCCGCTTCTTCCAGAAGGGCCAAG...GGACTTGCT 976
 | | | | | | | | | | | | | | | | | | | | | |
 1799 GGCTGCGGCCCATGGGGACTTCTTTAAGAAAACCAAAGGGCGGCCTGGAT 1848

FIGURE 6 (cont'd)

1665 CCACATCCTTATCCTGCTAGGGGGGATCTACCTCCTCGTGGGCCAGCTGT 1714
 | | | | | | | | | | | | | | | | | | | | | |
 2543 AGAGATCCTGTCTGTGTTAGGAGGAGTCTACTTCTTTTCCGAGGGATTC 2592
 1715 GGTACTTCTGGCGGGCCACGTGTTTCTGATCTCGTTCATAGACAGC 1764
 | | | | | | | | | | | | | | | | | | | | | |
 2593 AGTATTTCTGCAGAGGCGGCCGTCGATGAAGACCCTGTTTGTGGACAGC 2642
 1765 TACTTTGAAATCCTCTTCTGTTCCAGGCCCTGCTCACAGTGGTGTCCCA 1814
 | | | | | | | | | | | | | | | | | | | | | |
 2643 TACAGTGAGATGCTTTTCTTTCTGCAGTCACTGTTTCTGCTGGCCACCGT 2692
 1815 GGTGCTGTGTTTCTGGCCATCGAGTGGTACCTGCCCCTGCTTGTGTCTG 1864
 | | | | | | | | | | | | | | | | | | | | | |
 2693 GGTGCTGTACTTCAGCCACCTCAAGGAGTATGTGGCTTCCATGGTATTCT 2742
 1865 CGCTGGTGTCTGGGCTGGCTGAACCTGCTTTACTATACAGTGGCTTCCAG 1914
 | | | | | | | | | | | | | | | | | | | | | |
 2743 CCCTGGCCTTGGGCTGGACCAACATGCTCTACTACACCGCGGTTCAG 2792
 1915 CACACAGGCATCTACAGTGTCTGATCCAGAAGGTCTCCTGCGGGACCT 1964
 | | | | | | | | | | | | | | | | | | | | | |
 2793 CAGATGGGCATCTATGCCGTCATGATAGAGAAGATGATCCTGAGAGACCT 2842
 1965 GCTGCGCTTCCTTCTGATCTACTTAGTCTTCTTTTCGGCTTCGCTGTAG 2014
 | | | | | | | | | | | | | | | | | | | | | |
 2843 GTGCCGTTTCATGTTTGTCTACATCGTCTTCTTGTTCGGGTTTTCCACAG 2892
 2015 CCCTGGTGAGCCTGAGCCAGGAGGCTTGGCGCCCCGAAGCTCCTACAGGC 2064
 | | | | | | | | | | | | | | | | | | | | | |
 2893 CGGTGGTGACGCTGATTGAAGACGGGAAGAATGACTCCCTGCCGTCTGAG 2942
 2065 CCCAATGCCACAGAGTCAGTGCAGCCCATGGAGGGACAGGAGGACGAGGG 2114
 | | | | | | | | | | | | | | | | | | | | | |
 2943 TCCA.....CGTCGCACAGGTGGCGGGGGCCTGCCTGCAGGCC 2980
 2115 CAACGGGGCCAGTACAGGGGTATCCTGGAAGCCTCCTTGGAGCTCTTCA 2164
 | | | | | | | | | | | | | | | | | | | | | |
 2981 CCCCAGATAGCTCCTACAACAGCCTGTACTCCACCTGCCTGGAGCTGTTCA 3030
 2165 AATTCACCATCGGCATGGGCGAGCTGGCCTTCCAGGAGCAGCTGCACTTC 2214
 | | | | | | | | | | | | | | | | | | | | | |
 3031 AGTTCACCATCGGCATGGGCGACCTGGAGTTCAGTGAAGTATGACTTC 3080
 2215 CGCGGCATGGTGCTGCTGCTGCTGGCCTACGTGCTGCTCACCTACAT 2264
 | | | | | | | | | | | | | | | | | | | | | |
 3081 AAGGCTGTCTTCATCATCTGCTGCTGGCCTATGTAATTCTCACCTACAT 3130
 2265 CCTGCTGCTCAACATGCTCATCGCCCTCATGAGCGAGACCGTCAACAGTG 2314
 | | | | | | | | | | | | | | | | | | | | | |
 3131 CCTCTGCTCAACATGCTCATCGCCCTCATGGGTGAGACTGTCAACAAGA 3180
 2315 TCGCCACTGACAGCTGGAGCATCTGGAAGCTGCAGAAAGCCATCTCTGTC 2364
 | | | | | | | | | | | | | | | | | | | | | |
 3181 TCGCACAGGAGAGCAAGAACATCTGGAAGCTGCAGAGAGCCATCACCATC 3230

FIGURE 6 (cont'd)

2365 CTGGAGATGGAGAATGGCTATTGGTGGTGCAGGAAGAAG...CAGCGGGC 2411
 ||||| | ||||| ||| ||||| || ||||| || |||
 3231 CTGGACACGGAGAAGAGCTTCCTTAAGTGCATGAGGAAGGCCTTCCGCTC 3280
 2412 AGGTGTGATGCTGACCGTTGGCACTAAGCCAGATGGCAGCCCGGATGAGC 2461
 ||| | ||||| || || | || ||||| || |||
 3281 AGGCAAGCTGCTGCAGGTGGGGTACACACCTGATGGCAAGGACGACTACC 3330
 2462 GCTGGTGCTTCAGGGTGGAGGAGGTGAACTGGGCTTCATGGGAGCAGACG 2511
 | ||||| ||||| ||||| ||||| || | ||| |
 3331 GGTGGTGCTTCAGGGTGGACGAGGTGAACTGGACCACCTGGAACACCAAC 3380
 2512 CTGCCTACGCTGTGTGAGGACCCG...TCAGGGGCAGGTGTCCCTCGAAC 2558
 || | | || ||||| || || || ||| || |||
 3381 GTGGGCATCATCAACGAAGACCCGGGCAACTGTGAGGGCGTCAAGCGCAC 3430
 2559 TCTCGAGAACCCTGTCTTG...GCTTCCCCTCCCAAGGAGGATGAGGAT 2604
 || | | || || || || || || || ||
 3431 CCTGAGCTTCTCCCTGCGGTCAAGCAGAGTTTCAGGCAGACACTGGAAGA 3480
 2605 GGTGCCTCTGAGGAAACTATGTGCCCCGTCCAGCTCCTCCAGTCCAACCTG 2654
 | | | | || || || || || || || || || ||
 3481 ACTTTGCCCTGGTCCCCCTTTAAGAGAGGCCAAGTGCTCGAGATAGGCAG 3530
 2655 ATGGCCCAGATGCAGCAGGAGGCCAGAGGACAGAGCAGAGGATCTTTCCA 2704
 || ||| || || | || || || || || || || ||
 3531 TCTGCTCAGCCCGAGGAAGTTTATCTGCGACAGTTTTTCAGGGTCTCTGAA 3580
 2705 ACCACATCTGCTGGCTCTGGGGTCCCAGTGAATTCTGGTGGCAAATATAT 2754
 ||| | | || || || || || || || || || ||
 3581 GCCA.....GAGGACGCTGAGGTCTTCAAGAGTCCTGCCGCTTCCGGGGA 3625
 2755 ATTTTCACTAACTCAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAA 2804
 | | | ||| || | || | || | || | || | || |
 3626 GAAGTGAGGACGTACGCAGACAGCACTGTCAACACTGGGCCTTAGGAGA 3675
 2805 AAAAA..... 2809
 3676 CCCCCTTGCCACGGGGGGCTGCTGAGGGAACACCAGTGCTCTGTCTCAGCAG 3725

FIGURE 6 (cont'd)

CLUSTAL W (1.74) multiple sequence alignment

```

humanVR2      MTSPSSSPVFRLETLDDGGQEDGSEADRGKLDGSGGLPPMESQFQGEDRKFAPIQIRVNLNY
ratVR2      -----

humanVR2      RKG TGASQPD PNRFRDRDLFNAVSRGVPEDLAGLPEYLSKTSKYLT DSEYTEGSTGKTCL
ratVR2      -----

humanVR2      MKAVLN LKDG VNA CILPLLQIDRDSGNPQPLVNAQCTDDYYRGHSALHIAIEKRSLQC VK
ratVR2      -----

humanVR2      LLVENGANVHARACGRFFQKGQGTCTFYFGEPLPSLAACKQWDVVSYLLENPHQPASLQA
ratVR2      -----STHASALSLAACKQWDVVSYLLENPHQPASLEA
                      *****:*****:

humanVR2      TDSQGNTVLHALVMISDNSAENIALVTSMYDGLLQAGARLCPTVQLEDIRNLQDLTPLKL
ratVR2      TDSLGN TVLHALVMIADNSPENSALVIHMYDGLLQMGARLCPTVQLEEISNHQGLTPLKL
                      *** *****:***.*** ***** *****:***.***

humanVR2      AAKEGKIEIFRHILQREFSG-LSHL SRKFTWCYGPVRVSLYDLASVDSCEENSVLEIIA
ratVR2      AAKEGKIEIFRHILQREFSGPYQPLSRKFTWCYGPVRVSLYDLSSVDSWEKNSVLEIIA
                      ***** *****:***.*** *****

humanVR2      FHCKSPHRHRMVVLEPLNKL LQAKWDL LIPKFFLNFLCNLIYMFIFTAVAYHQPTLKKQA
ratVR2      FHCKSPNRHRMVVLEPLNKL LQEKWDLVSRFFNFACYLVMFIFTVAVYHQPSLDQPA
                      *****:***** ***** *:***:*** * *:*****.*****:*.

humanVR2      APHLKAEVGN SMLLTGHILIL LGGIYLLVGQLWYFWRRHVFIWISFIDS YFEILFLFQAL
ratVR2      IPSSKATFGESMLLLGHILIL LGGIYLLVGQLWYFWRRLFIWISFMDSYFEILFLFQAL
                      * ** *:*** *****:*****:*****:*****:***

humanVR2      LTVVSQVLCFLAIEWYLPLLVSALVLGWLNL LYYTRGFQHTGIYSVMIQKVILRDLLRFL
ratVR2      LTVLSQVLRFMETEWYLP LLVLSLVLGWLNL LYYTRGFQHTGIYSVMIQKVILRDLLRFL
                      ***:*** *: ***** :*****:*****:*****:*****

humanVR2      LIYLVFLFGFAVALVSLSQEAWRPEAPTGP NATESVQPMEGQEDEGNGAQYRGILEASLE
ratVR2      LVYLVFLFGFAVALVSLSR EARS PKAPEDNNSTVTEQPTVGQEEEP--APYRSILDASLE
                      *:*****:*** *:***. *: * ** *:***

humanVR2      LFKFTIGMGELAFQEQ LHF RGMV LLL LAYVLLTYILLINMLIALMSETVNSVATDSWSI
ratVR2      LFKFTIGMGELAFQEQ LRF RGMV LLL LAYVLLTYVLLINMLIALMSETVNHVADNSWSI
                      *****:***:*****:*****:***** ** :***

humanVR2      WKLQKAISVLEMENGYW WCR-KKQ RAGV MLTVG TKPDGSPDERWC FRVEEVN WASWEQTL
ratVR2      WKLQKAISVLEMENGYW WCRKKHREGRL LKVGTRGDGTPDERWC FRVEEVN WAAWEKTL
                      ***** **:* *:***: *:*****:***:***

humanVR2      PTLCEDPSGAGVPR TLENPVLASPPKEDE DGASEENYVPVQLLQSN
ratVR2      PTLSEDPSPGIGITGN KKNPTSK-PGK---NSASEDHLPLQVLQSP
                      ***.*****:*. . :***. * * :.*****:***:***

```

FIGURE 7

GAP of: ratvr2.pep check: 9190 from: 1 to: 554

ratVR2 Flrxb147g11

to: humanvr2.pep check: 5746 from: 1 to: 764

humanVR2 Flh21e11

Symbol comparison table: /usr/local/gcg_9.1/gcgcore/data/rundata/blosum62.cmp
CompCheck: 6430

Gap Weight:	12	Average Match:	2.912
Length Weight:	4	Average Mismatch:	-2.003
Quality:	2182	Length:	766
Ratio:	3.939	Gaps:	4
Percent Similarity:	81.703	Percent Identity:	79.167

Match display thresholds for the alignment(s):

| = IDENTITY
: = 2
. = 1

ratvr2.pep x humanvr2.pep

1STHASALSIACTKQWDVVITYLLENPHQPASLEATDSLGNITVLH 44
|||||
201 GQGTCTFYFGEPLPSLAIACTKQWDVVSYLLENPHQPASLQATDSQGNITVLH 250
45 ALVMIADNSPENSALVIHMYDGLLQMGARLCPTVQLEEISNHQGLTPLKL 94
|||||
251 ALVMISDNSAENIALVTSMYDGLLQAGARLCPTVQLEDIRNLQDLTPLKL 300
95 AAKEGKIEIFRHILQREFSGPYQPLSRKFTWCYGPVRVSLYDLSSVDSW 144
|||||
301 AAKEGKIEIFRHILQREFSG.LSHLSRKFTWCYGPVRVSLYDLASVDSC 349
145 EKNSVLEIIAFHCKSPNRHRMVLEPLNKLLOEKWDRLVSRFFNFACYL 194
|.|||||
350 EENSIVLEIIAFHCKSPHRHRMVLEPLNKLLOAKWDLIPKFFLNFLCNL 399
195 VYMFIFTVVAYHQPSLDQPAIPSSKATFGESMLLGHILILLGGIYLLG 244
:|||||
400 IYMFIFTAVAYHQPTLKKQAAPHLKAEVGNMMLTGHILILLGGIYLLVG 449
245 QLMYFWRRRLFIWISFMDSYFEILFLQALLTVLSQVLRFMETEWYLPIL 294
|||||
450 QLMYFWRRRHVFIWISFIDSIFEILFLQALLTVVSQVLCFLAIEWYLPIL 499
295 VLSVLGWLNLNLYYTRGFQHTGIYSVMIOKVILRDLLRFLLVYLVFLFGF 344
|.|||||
500 VSALVLGWLNLNLYYTRGFQHTGIYSVMIOKVILRDLLRFLLIYLVFLFGF 549
345 AVALVSLSRREARSPKAPEDNNSTVTEQPTVGQERE..PAPYRSILDASLE 392
|||||
550 AVALVSLSQEAWRPEAPTGNATESVQPMEGQDEGNGAQYRGILEASLE 599

FIGURE 8

393 LFKFTIGMGELAFQEQLRFRGVVLLLLLAYVLLTYVLLLNMLIALMSETV 442
 |||||:|||||
 600 LFKFTIGMGELAFQEQLHFRGMVLLLLLAYVLLTYILLNMLIALMSETV 649
 443 NHVADNSWSIWKLQKAISVLEMENGYWWCRRKKHREGRLKVGTRGDGTP 492
 | | . |||||:| ||: ||.
 650 NSVATDSWSIWKLQKAISVLEMENGYWWC.RKKQRAGVMLTVGTPKPDGSP 698
 493 DERWCFRVEEVNWAANEKTLPTLSEDPSGPGITGNKKNPT....SKPGKN 538
 |||||.||.|||| |: .|| |..
 699 DERWCFRVEEVNWAWEQTLPTLCEDPGAGVPRTLENPVLASPPKEDED 748
 539 SASEEDHLPLOVLQSP 554
 |||.:.|.|||
 749 GASEENYVPVQLQSN 764

FIGURE 8 (continued)

1401 CTGTCCCAGGACTCTGTGCGCCGCCAGCACCCGAGAAGACCCTCAGGCTCTA 1450
 ||||| ||| ||| | ||||| ||| |||||
 369 TCATCCCAGGACTCCGTCTCCGCTGG...TGAGAAGCCCCCGAGGCTCTA 415
 1451 TGATCGCAGGAGTATCTTTGAAGCCGTTGCTCAGAATAACTGCCAGGATC 1500
 ||||| ||||| ||| ||| ||| ||||| ||||| ||||| |||||
 416 TGATCGCAGGAGCATCTTCGATGCTGTGGCTCAGAGTAACTGCCAGGAGC 465
 1501 TGGAGAGCCTGCTGCTCTTCCTGCAGAAGAGCAAGAAGCACCTCACAGAC 1550
 ||||| ||||| ||||| ||||| ||||| ||||| ||| ||| |||
 466 TGGAGAGCCTGCTGCCCTTCCTGCAGAGGAGCAAGAAGCGCCTGACTGAC 515
 1551 AACGAGTTCAAAGACCCTGAGACAGGGAAGACCTGTCTGCTGAAAGCCAT 1600
 | ||||| ||||| ||||| ||||| ||||| ||||| ||||| |||||
 516 AGCGAGTTCAAAGACCCAGAGACAGGAAAGACCTGTCTGCTAAAAGCCAT 565
 1601 GCTCAACCTGCACGACGGACAGAACACCACCATCCCCCTGCTCCTGGAGA 1650
 ||||| ||||| | || ||||| ||||| | ||||| |||||
 566 GCTCAATCTGCACAATGGGCAGAATGACACCATCGCTCTGCTCCTGGACG 615
 1651 TCGCGCGGCAAACGGACAGCCTGAAGGAGCTTGTCAACGCCAGCTACACG 1700
 | || ||| | || ||||| ||||| || ||||| ||||| |||||
 616 TTGCCCCGAAGACAGACAGCCTGAAGCAGTTTGTCAATGCCAGCTACACA 665
 1701 GACAGCTACTACAAGGGCCAGACAGCACTGCACATCGCCATCGAGAGACG 1750
 ||||| ||||| ||||| ||||| ||||| ||||| ||||| |||||
 666 GACAGCTACTACAAGGGCCAGACAGCACTGCACATTGCCATTGAACGGCG 715
 1751 CAACATGGCCCTGGTGACCCTCCTGGTGGAGAACGGAGCAGACGTCCAGG 1800
 ||||| | ||||| ||||| ||||| ||||| ||||| |||||
 716 GAACATGACGCTGGTGACCCTCCTGGTGGAGAATGGAGCAGATGTCCAGG 765
 1801 CTGCGGCCCATGGGGACTTCTTTAAGAAAACCAAAGGGCGGCCTGGATTC 1850
 ||||| | ||||| ||||| ||||| ||||| ||||| ||||| |||
 766 CTGCGGCTAACGGGGACTTCTTCAAGAAAACCAAAGGGAGGCCTGGCTTC 815
 1851 TACTTCGGTGAAGTTCCTGCTGCAGAACTCCTGGCAGACGGCCGACATCAGCG 1900
 |||| | |||| | ||||| ||||| ||||| ||||| ||||| |||||
 816 TACTTTGGTGAGCTGCCCCTGTCCCTGGCTGCGTGCACCAACCAGCTGGC 865
 1901 CATCGTGAAGTTCCTGCTGCAGAACTCCTGGCAGACGGCCGACATCAGCG 1950
 ||| ||||| ||||| ||||| ||||| ||||| ||||| ||||| |||||
 866 CATTGTGAAGTTCCTGCTGCAGAACTCCTGGCAGCCTGCAGACATCAGCG 915
 1951 CCAGGGACTCGGTGGGCAACACGGTGTGCACGCCCTGGTGGAGGTGGCC 2000
 || ||||| ||||| ||||| ||||| ||||| ||||| ||||| |||||
 916 CCCGGGACTCAGTGGGCAACACGGTGTTCATGCCCTGGTGGAGGTGGCA 965
 2001 GACAACACGGCCGACAACACGAAGTTTGTGACGAGCATGTACAATGAGAT 2050
 || ||||| | ||||| ||||| ||||| ||||| ||||| ||||| |||||
 966 GATAACACAGTTGACAACACCAAGTTCGTGACAAGCATGTACAACGAGAT 1015
 2051 TCTGATGCTGGGGGCCAAACTGCACCCGACGCTGAAGCTGGAGGAGCTCA 2100
 |||| ||||| ||||| ||||| ||||| ||||| ||||| ||||| |||||
 1016 CTTGATCCTGGGGGCCAAACTCCACCCACGCTGAAGCTGGAAGAGATCA 1065

FIGURE 9 (cont'd)

2101 CCAACAAGAAGGGAATGACGCCGCTGGCTCTGGCAGCTGGGACCGGGAAG 2150
 ||||| ||||| | ||||| ||||| ||||| ||||| |||||

1066 CCAACAGGAAGGGGCTCACGCCACTGGCTCTGGCTGCTAGCAGTGGGAAG 1115

2151 ATCGGGGTCTTGGCCTATATTCTCCAGCGGGAGATCCAGGAGCCCGAGTG 2200
 ||||| ||||| ||||| ||||| ||||| ||||| ||||| |||||

1116 ATCGGGGTCTTGGCCTACATTCTCCAGAGGGAGATCCATGAACCCGAGTG 1165

2201 CAGGCACCTGTCCAGGAAGTTACCCGAGTGGGCCTACGGGCCCGTGCACT 2250
 | | ||||| ||||| ||||| ||||| ||||| ||||| |||||

1166 CCGACACCTATCCAGGAAGTTACCCGAATGGGCCTATGGGCCAGTGCACT 1215

2251 CCTCGCTGTACGACCTGTCCTGCATCGACACCTGCGAGAAGAACTCGGTG 2300
 ||||| ||||| ||||| ||||| ||||| ||||| ||||| |||||

1216 CCTCCCTTTATGACCTGTCCTGCATTGACACCTGTGAAAAGAACTCGGTT 1265

2301 CTGGAGGTGATCGCCTACAGCAGCAGCGAGACCCCTAATCGCCACGACAT 2350
 ||||| ||||| ||||| ||||| ||||| ||||| ||||| |||||

1266 CTGGAGGTGATCGCTTACAGCAGCAGTGAACCCCTAACCGTCATGACAT 1315

2351 GCTCTTGGTGGAGCCGCTGAACCGACTCCTGCAGGACAAGTGGGACAGAT 2400
 ||||| ||||| ||||| ||||| ||||| ||||| ||||| |||||

1316 GCTTCTCGTGAACCCCTGAACCGACTCCTACAGGACAAGTGGGACAGAT 1365

2401 TCGTCAAGCGCATCTTCTACTTCAACTTCCTGGTCTACTGCCTGTACATG 2450
 | ||||| ||||| ||||| ||||| ||||| ||||| ||||| |||||

1366 TTGTCAAGCGCATCTTCTACTTCAACTTCTTCGTCTACTGCTTGTATATG 1415

2451 ATCATCTTCACCATGGCTGCCTACTACAGGCCCGTGGATGGCTTGCCCTCC 2500
 ||||| ||||| ||||| ||||| ||||| ||||| ||||| |||||

1416 ATCATCTTCACCGCGGCTGCCTACTATCGGCCTGTGGAAGGCTTGCCCCC 1465

2501 CTTTAAGATGGAAAAA...TTGGAGACTATTTCCGAGTTACTGGAGAGA 2547
 || |||| || |||| | |||| ||||| ||||| ||||| |||||

1466 CTATAAGCTGAAAAACACCGTTGGGGACTATTTCCGAGTCACCGGAGAGA 1515

2548 TCCTGTCTGTGTTAGGAGGAGTCTACTTCTTTTCCGAGGGATTCAATAT 2597
 || ||||| ||||| ||||| ||||| ||||| ||||| ||||| |||||

1516 TCTTGTCTGTGTCAGGAGGAGTCTACTTCTTCTTCCGAGGGATTCAATAT 1565

2598 TTCCTGCAGAGGGCGGCCGTCGATGAAGACCCTGTTTGTGGACAGCTACAG 2647
 ||||| ||||| ||||| ||||| ||||| ||||| ||||| |||||

1566 TTCCTGCAGAGGGCGACCATCCCTCAAGAGTTGTTTGTGGACAGCTACAG 1615

2648 TGAGATGCTTTTCTTTCTGCAGTCACTGTTTCATGCTGGCCACCGTGGTGC 2697
 ||||| ||||| ||||| ||||| ||||| ||||| ||||| |||||

1616 TGAGATACTTTTCTTTGTACAGTCGCTGTTTCATGCTGGTGTCTGTGGTAC 1665

2698 TGTACTIONCAGCCACCTCAAGGAGTATGTGGCTTCCATGGTATTCTCCCTG 2747
 ||||| ||||| ||||| ||||| ||||| ||||| ||||| |||||

1666 TGTACTIONCAGCCAACGCAAGGAGTATGTGGCTTCCATGGTGTCTCCCTG 1715

2748 GCCTTGGGCTGGACCAACATGCTCTACTACACCCGCGGTTTCCAGCAGAT 2797
 ||||| ||||| ||||| ||||| ||||| ||||| ||||| |||||

1716 GCCATGGGCTGGACCAACATGCTCTACTATACCCGAGGATTCCAGCAGAT 1765

FIGURE 9 (cont'd)

2463 CTTCTGAGGGATGCAAGCACTCGAGATAGACATGCCACCCAGCAGGAAGA 2512
 3548 AGTTTATCTGCGACAGTTTTTCAGGGTCTCTGAAGCCAGAGGACGCTGAGG 3597
 2513 AGTTCAACTGAAGCATTATACGGGATCCCTTAAGCCAGAGGATGCTGAGG 2562
 3598 TCTTCAAGAGTCCTGCCGCTTCCGGGGAGAAAGTGA.GGACGTCACGCAGA 3646
 2563 TTTTCAAGGATTCCATGGTCCCAGGGGAGAAATAATGGACACTATGCAGG 2612
 3647 CAGCACTGTCAACACTGGGCCTTAGGAGACCCCGTTGCCACGGGGGGCTG 3696
 2613 GATCAATG.....CGGGGTCTTTGGGTGGTCTG 2640
 3697 CTGAGGGAACACCAAGTGCTCTGTGCAGCAGCCTGGCCTGGTCTGTGCCTGC 3746
 2641 CTTAGGGAAC.CAGCAGGGTTGACGTTATCTGGGTCCACTCTGTGCCTGC 2689
 3747 CCA.GCATGTTCCCAAATCTGTGCTGGACAAGCTGTGGGAAGCGTTCTTG 3795
 2690 CTAGGCACATTCTTAGGACTTCGGCGGGCCTGCTGTGGGAA.CTGGGAGG 2738
 3796 GAAGCATGGGGAGTGATGTACATCCAACCGTCACTGTCCCCAAGTGAATC 3845
 2739 TGTGTGGGAATTGAGATGTGTATCCAACCATGA...TCTCCAAACATTTG 2785
 3846 TCCTAACAGACTTTTCAGGTTTTTACTCACTTTACTAAAAAAAAAAAAAAAAA 3895
 2786 GCTTTCAACTCTTTATGGACTTTATTAAACAGAGTGAATGGCAAATCTCT 2835
 3896 AGGGCGGCCGCTTA 3909
 2836 ACTTGGACACAT.. 2847

FIGURE 9 (cont'd)

GAP of: humanvr1.pep check: 6877 from: 1 to: 839

humanVR1_Fbh18547pat - fchrb87a6, 3909 bases, 4554 checksum.

to: ratvr1.pep check: 5764 from: 1 to: 838

ratVR1 | AF029310 Rattus norvegicus vanilloid receptor subtype 1 mRNA,
complete
cds.

Symbol comparison table:

/ddm_local/gcg/gcg_9.1/gcgcore/data/rundata/blosum62.cmp

CompCheck: 6430

Gap Weight: 12 Average Match: 2.912
Length Weight: 4 Average Mismatch: -2.003

Quality: 3734 Length: 840
Ratio: 4.456 Gaps: 3
Percent Similarity: 89.247 Percent Identity: 86.022

Match display thresholds for the alignment(s):

| = IDENTITY
: = 2
. = 1

humanvr1.pep x ratvr1.pep

```
1 MKKWSSTDLGTAADPLQKDTCPDPLDGDPNRPPPAKPQLPTAKSRTLRF 50
|.. .| | . | |...| || | ||| :|| | | : | :|||
1 MEQRASLDSEESPPQENSCLDPPDRDPNCKPPPVPKPHIFTTRSRTLRF 50

51 GKGDSEEAFFVDCPHEEGELDSCPTITVSPVITIQRPGDGPTGARLLSQD 100
| | | | | | | | | | | | | | | | | | | | | | | | | | | |
51 GKGDSEEA SPLDCPYEEGGLASCPIITVSSVLTIQRPDGPASVRPSSQD 100

101 SVAASTEKTRLRYDRRSIFEAVAQNNCQDLESLLLFLQSKKHLTDNEFK 150
| | . | | | | | | | | | | | | | | | | | | | | | | |
101 SVSAG.EKPPRLYDRRSIFDAVAQSNCOELESLLPFLQRSKKRLTDSEFK 149

151 DPETGKTCLLKAMLNLHDGQNTTIPLLLIEIARQTDLSKELVNASYTDYY 200
| | | | | | | | | | | | | | | | | | | | | | | | | |
150 DPETGKTCLLKAMLNLHNGQNDTIALLLDVARKTDSLKQFVNASYTDYY 199

201 KGQTALHIAIERRNMALVTLLVENGADVQAAAHGDFFKKTKGRPGFYFGE 250
| | | | | | | | | | | | | | | | | | | | | | | | | |
200 KGQTALHIAIERRNMTLVTLVENGADVQAAANGDFFKKTKGRPGFYFGE 249

251 LPLSLAACTNQLGIVKFLQNSWQTADISARDSVGNTVLHALVEVADNTA 300
| | | | | | | | | | | | | | | | | | | | | | | | | |
250 LPLSLAACTNQLAIVKFLQNSWQPADISARDSVGNTVLHALVEVADNTV 299

301 DNTKFVTSMYNEILMLGAKLHPTLKLEELTNKKGMTPLALAAGTGKIGVL 350
| | | | | | | | | | | | | | | | | | | | | | | | | |
300 DNTKFVTSMYNEILILGAKLHPTLKLEEITNRKGLTPLALAASSGKIGVL 349

351 AYILQREIQEPECRHLRSRKFTWAYGPVHSSLYDLSCIDTCEKNSVLEVI 400
| | | | | | | | | | | | | | | | | | | | | | | | | |
350 AYILQREIHEPECRHLRSRKFTWAYGPVHSSLYDLSCIDTCEKNSVLEVI 399
```

FIGURE 10

```

401 AYSSSETPNRHDMLLVEPLNRLQLQDKWDRFVKRIFYFNFLVYCLYMIIFT 450
|||||:|||||:|||||:|||||:|||||:|||||:|||||:|||||:|||||:
400 AYSSSETPNRHDMLLVEPLNRLQLQDKWDRFVKRIFYFNFFVYCLYMIIFT 449

451 MAAYYRPVDGLPPFKMEK.IGDYFRVTGEILSVLGGVYFFFRGIQYFLQR 499
|||||:|||||:|||||:|||||:|||||:|||||:|||||:|||||:
450 AAAYYRPVEGLPPYKLNKNTVGDYFRVTGEILSVSGGVYFFFRGIQYFLQR 499

500 RPSMKTLFVDSYSEMLFFLQSLFMLATVVLYFSLKEYVASMVFSALGW 549
|||:|.|||||.|||||.|||||.|||||.|||||.|||||.|||||:|
500 RPSLKSFLVDSYSEILFFVQSLFMLVSVVLYFSQRKEYVASMVFSLAMGW 549

550 TNMLYYTRGFQQMGIYAVMIEKMILRDLCRFMFVYIVFLFGFSTAVVTLI 599
|||||:|||||:|||||:|||||:|||||:|||||:|||||:|||||:
550 TNMLYYTRGFQQMGIYAVMIEKMILRDLCRFMFVYLVFLFGFSTAVVTLI 599

600 EDGKNDSLPESTSHRWRGPACRPPDSSYNSLYSTCLELFKFTIGMGDLE 649
|||||.||| ||| |: || |: | .|||||:|||||:|||||:|||||:
600 EDGKNNSLPMESTPHKCRGSACK.PGNSYNSLYSTCLELFKFTIGMGDLE 648

650 FTENYDFKAVFIILLLAYVILTYILLNMLIALMGETVNKIAQESKNIWK 699
|||||:|||||:|||||:|||||:|||||:|||||:|||||:|||||:
649 FTENYDFKAVFIILLLAYVILTYILLNMLIALMGETVNKIAQESKNIWK 698

700 LQRAITILDTEKSFLKCMRKAFRSGKLLQVGYTPDGKDDYRWCFRVDEVN 749
|||||:|||||:|||||:|||||:|||||:|||||:|||||:|||||:
699 LQRAITILDTEKSFLKCMRKAFRSGKLLQVGFTPDGKDDYRWCFRVDEVN 748

750 WTTWNTNVGIINEDPGNCEGVKRTLSFSLRSSRVSGRHWKNFALVPLLRE 799
|||||:|||||:|||||:|||||:|||||:|||||:|||||:|||||:
749 WTTWNTNVGIINEDPGNCEGVKRTLSFSLRSGRVSGRNWKNFALVPLLRD 798

800 ASARDRQSAQPEEVYLRQFSGSLKPEDA EVFKSPAASGEK 839
|| ||| . | ||| |: :.|||||:|||||:|||||:
799 ASTRDRHATQQEEVQLKHYTGSLKPEDA EVFKDSMVPGEK 838

```

FIGURE 10 (cont'd)

CLUSTAL W (1.74) multiple sequence alignment

```

humanVR2.alt -----
humanVR2      MTSPSSSPVFRLETLDGGQEDGSEADRGKLDGSGLPMPESQFQGEDRKFAPOIRVNLNY

humanVR2.alt -----
humanVR2      RKG TGASQDPNRFDRDLFNAVSRGVPEDLAGLPEYLSKTSKYLTDSEYTEGSTGKTCL

humanVR2.alt -----
humanVR2      MKAVLNLKDGVNACILPLLQIDRDSGNPQPLVNAQCTDDYYRGHSALHIAIEKRSLQCVK

humanVR2.alt -----GRFFQKGQGTCTFYFGELPLSLAACTKQWDVVSYLENPHQPASLQA
humanVR2      LLVENGANVHARACGRFFQKGQGTCTFYFGELPLSLAACTKQWDVVSYLENPHQPASLQA
                *****

humanVR2.alt TDSQGN TVLHALVMISDN SAENIALVTSMYDGLLQAGARLCPTVQLEDIRNLQDLTPLKL
humanVR2      TDSQGN TVLHALVMISDN SAENIALVTSMYDGLLQAGARLCPTVQLEDIRNLQDLTPLKL
                *****

humanVR2.alt AAKEGKIEIFRHILQREFSGLSHLSRKFTWCYGPVRVSLYDLASVDSCEENSVLEIIAF
humanVR2      AAKEGKIEIFRHILQREFSGLSHLSRKFTWCYGPVRVSLYDLASVDSCEENSVLEIIAF
                *****

humanVR2.alt HCKSPHRHRMVVLEPLNKLLOAKWDLIPKFFINFLCNLIYMFIFTAVAYHQPTLKKQAA
humanVR2      HCKSPHRHRMVVLEPLNKLLOAKWDLIPKFFINFLCNLIYMFIFTAVAYHQPTLKKQAA
                *****

humanVR2.alt PHLKAEVGN SMLLTGHI LILLGGIYLLVGQLWYFWRRHVFIWISFIDSYFEILFLFQALL
humanVR2      PHLKAEVGN SMLLTGHI LILLGGIYLLVGQLWYFWRRHVFIWISFIDSYFEILFLFQALL
                *****

humanVR2.alt TVVSQVLCFLAIEWYLP LLVSALVLGWLNL LYYTRGFQHTGIYSVMIQ-----
humanVR2      TVVSQVLCFLAIEWYLP LLVSALVLGWLNL LYYTRGFQHTGIYSVMIQVILRDLLRFL
                *****

humanVR2.alt -----
humanVR2      IYLVFLFGFAVALVSLSQEAWRPEAPTGNATESVQPMEGQEDEGNGAQYRGILEASLEL

humanVR2.alt -----
humanVR2      FKFTIGMGELAFQEQ LHFGRMVLLLLL LAYVLLTYILLNMLIALMSETVNSVATDSWSIW

humanVR2.alt --KKAISVLEMENGYWWCRKKQ RAGVMLTVGTPDGSPDERWC FRVEEVN WASWEQTLPT
humanVR2      K LQKAISVLEMENGYWWCRKKQ RAGVMLTVGTPDGSPDERWC FRVEEVN WASWEQTLPT
                ;*****

humanVR2.alt LCEDPSGAGVPRTLENPVLASPPKEDEDGASEENYVPVQLLQSN
humanVR2      LCEDPSGAGVPRTLENPVLASPPKEDEDGASEENYVPVQLLQSN
                *****

```

FIGURE 11




FIGURE 12

Protein Family / Domain Matches, HMMer versión 2

Searching for complete domains

hmmpfam - search a single seq against HMM database

HMMER 2.1.1 (Dec 1998)

Copyright (C) 1992-1998 Washington University School of Medicine

HMMER is freely distributed under the GNU General Public License (GPL).

HMM file: /prod/ddm/seqanal/PFAM/pfam4.2/Pfam
Sequence file: /usr/ns-home/docs/seqanal/orfanal/oa-script.18670.seq

Query: hVR-1

Scores for sequence family classification (score includes all domains):

Model	Description	Score	E-value	N
ank	Ank repeat	51.5	1.9e-11	3

Parsed for domains:

Model	Domain	seq-f	seq-t	hmm-f	hmm-t	score	E-value
ank	1/3	201	233	1	33	34.4	2.6e-06
ank	2/3	248	283	1	33	13.2	2
ank	3/3	333	361	1	33	3.4	26

Alignments of top-scoring domains:

ank: domain 1 of 3, from 201 to 233: score 34.4, E = 2.6e-06

->nGnTPLHlAarygnvevvklLLehGAdvnartk<-
+G+T+LH+A + n+ +v lL+e+GAdv a+

hVR-1 201 KGQTALHIAIERNMALVTLLVENGADVQAAAH 233

ank: domain 2 of 3, from 248 to 283: score 13.2, E = 2

->nGnTPLHlAarygnvevvklLLe...hGAdvnartk<-
G PL lAa ++++ +vk+LL+++ + Ad+ ar+

hVR-1 248 FGELPLSLAACTNQLGIVKFLlQnswQTADISARDS 283

ank: domain 3 of 3, from 333 to 361: score 3.4, E = 26

->nGnTPLHlAarygnvevvklLLehGAdvnartk<-
+G TPL lAa +g++ v ++ L+ ++

hVR-1 333 KGMTPLALAAGTGKIGVLAYILQ-----REIQEP 361

FIGURE 13




FIGURE 14

Protein Family / Domain Matches, HMMer version 2

Searching for complete domains

hmmpfam - search a single seq against HMM database

HMMER 2.1.1 (Dec 1998)

Copyright (C) 1992-1998 Washington University School of Medicine

HMMER is freely distributed under the GNU General Public License (GPL).

HMM file: /prod/ddm/seqanal/PFAM/pfam3.3/Pfam
Sequence file: /tmp/orfana1.579.aa

Query: Flh21c11

Scores for sequence family classification (score includes all domains):

Model	Description	Score	E-value	N
ank	PF00023 Ank repeat	53.7	4e-12	3

Parsed for domains:

Model	Domain	seq-f	seq-t	hmm-f	hmm-t	score	E-value
ank	1/3	162	194	1	33	38.3	1.7e-07
ank	2/3	208	243	1	33	6.4	4.3
ank	3/3	293	328	1	33	8.8	2.1

Alignments of top-scoring domains:

ank: domain 1 of 3, from 162 to 194: score 38.3, E = 1.7e-07

```

*--> nGnTFLHIAarygnvevkvlllehgAdvnartk<--*
      4G+++LH+A ++ ++++vklle+GA+v+ar
Flh21c11 162  EGHSALEHTATEKRSLOQVLLVNGANVHARAC 194
  
```

ank: domain 2 of 3, from 208 to 243: score 6.4, E = 4.3

```

*--> nGnTFLHIAarygnvovvkvllle...hgAdvnartk<--*
      G FL IAs + +++vv +LL++++ A+ E++
Flh21c11 208  FGELESLAACKTKQMDVSVGLSphQPASLOATDS 243
  
```

ank: domain 3 of 3, from 293 to 328: score 8.8, E = 2.1

```

*--> nGnTFLHIAarygnvevkvllle...hgAdvnartk<--*
      + +TEL IAs+g+e+ + L+++ G + +r
Flh21c11 293  QQLATPLKLAKEOKLEIFRHILQraCSGLSHLSRK? 328
  
```

FIGURE 15

>hVR2.altFL (full-length) >predicted)
MTSPSSSPVFRLETLDGGQEDGSEADRGKLDFGSGLPPMESQFQGEDRKFAPQIRVNLNY
RKG TGASQPD PNRFD RDLFNAVSRGVPEDLAGLPEYLSKTSKYLT DSEYTEGSTGKTCL
AKAVLNLKDGVNACILPLLQIDRDSGNPQPLVNAQCTDDYYRGHSALHIAIEKRSLQCVK
LLVENGANVHARACGRFFQKGQGTCTFYFGELPLSLAACTKQWDVVSYLENPHQPASLQATDSQGNTVLHALVM
[SDNSAENIALVTSMYDGLLQAGARLCPTVQLEDIRNLQDLTPLKLAKEGKIEIFRHIL
REFSGLSHLSRKFTWCYGPVRVSLYDLASVDSCEENSVLEIIAFHCKSPHRHRMVVLE
?LNKLLQAKWDL LIPKFFLNFLCNLIYMFIFTAVAYHQPTLKKQAAPHLKAEVGNSMLLT
?HILILLGGIYLLVGQLWYFWRRHVFIWISFIDSYFEILFLFQALLTVVSQVLCFLAIEW
?LPLLVSALVLGWLNL LYYTRGFQHTGIYSVMIQKKAISVLEMENGYWWCRKKQRAGVML
VGTKPDGSPDERWCFRVEEVN WASWEQTLPTLCEDPSGAGVPRTLENPVLASPPKEDED
ASEENYVPVQLLQSN

Figure 16

CLUSTAL W (1.74) multip sequence alignment

```

humanVR2      MTSPSSSPVFRLETLDGGQEDGSEADRGKLDGSGGLPPMESQFQGEDRKFAPOIRVNLNY
hVR2.altFL    MTSPSSSPVFRLETLDGGQEDGSEADRGKLDGSGGLPPMESQFQGEDRKFAPOIRVNLNY
*****

humanVR2      RKG TGASQPD PNR FDR DRL FNA VSR GVP EDLAGL PEYLSKTSKYLTDSEYTEGSTGKTCL
nVR2.altFL    RKG TGASQPD PNR FDR DRL FNA VSR GVP EDLAGL PEYLSKTSKYLTDSEYTEGSTGKTCL
*****

humanVR2      MKAVLNLKDG VNA CIL PLLQIDR DSGNPQPLVNAQCTDDYYRGHSALHIAIEKRSLQCVK
iVR2.altFL    MKAVLNLKDG VNA CIL PLLQIDR DSGNPQPLVNAQCTDDYYRGHSALHIAIEKRSLQCVK
*****

humanVR2      LLVENGANV HARACGRFFQKGQGT CFYFGELPLSLAACTKQWDVVS YLLENPHQPASLQA
iVR2.altFL    LLVENGANV HARACGRFFQKGQGT CFYFGELPLSLAACTKQWDVVS YLLENPHQPASLQA
*****

humanVR2      TDSQGN TVLHALVMISD NSAENIALV TSMYDGLLQAGARLCPTVQLEDIRNLQDLTPLKL
VR2.altFL     TDSQGN TVLHALVMISD NSAENIALV TSMYDGLLQAGARLCPTVQLEDIRNLQDLTPLKL
*****

humanVR2      AAKEGKIEIFRHILQREFSGLSHLSRKFTWCYGPVRVSLYDLASVDSCEENSVLEIIAF
VR2.altFL     AAKEGKIEIFRHILQREFSGLSHLSRKFTWCYGPVRVSLYDLASVDSCEENSVLEIIAF
*****

humanVR2      HCKSPHRHRMVVLEPLNKL LQAKWDL LIPKFFLNFLCNLIYMFIFTAVAYHQPTLKKQAA
VR2.altFL     HCKSPHRHRMVVLEPLNKL LQAKWDL LIPKFFLNFLCNLIYMFIFTAVAYHQPTLKKQAA
*****

humanVR2      PHLKAEVGNSMLLTGHILIL LGGIYLLVGQLWYFWRRHVFIWISFIDSYFEILFLFQALL
VR2.altFL     PHLKAEVGNSMLLTGHILIL LGGIYLLVGQLWYFWRRHVFIWISFIDSYFEILFLFQALL
*****

humanVR2      TVVSQVLCFLAIEWYLPLLVSALVLGWLNL LYYTRGFQHTGIYSVMIQK VILRDLLRFL
VR2.altFL     TVVSQVLCFLAIEWYLPLLVSALVLGWLNL LYYTRGFQHTGIYSVMIQK-----
*****

humanVR2      IYLVFLFGFAVALVSLSQEAWRPEAPTGP NATESVQPMEGQEDEGNGAQYRGILEASLEL
VR2.altFL     -----

humanVR2      FKFTIGMGELAFQEQ LHF RGMV LLLL LAYVLLTY ILLL NMLIALMSETVNSVATDSWSIW
VR2.altFL     -----

humanVR2      KLQKAISVLEMENGYWWCRKKQRAGVMLTVG TKPDGSPDERWC FRVEEVN WASWEQTLPT
VR2.altFL     ---KAISVLEMENGYWWCRKKQRAGVMLTVG TKPDGSPDERWC FRVEEVN WASWEQTLPT
*****

humanVR2      LCEDPSGAGVPRTLENPVLASPPKEDEDGASEENYVPVQLLQSN
VR2.altFL     LCEDPSGAGVPRTLENPVLASPPKEDEDGASEENYVPVQLLQSN
*****

```

FIGURE 17

Declaration, Petition and Power of Attorney
for Continuation-in-Part Patent Application

As a below named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below next to my name,

I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled

NOVEL MEMBERS OF THE CAPSAICIN/VANILLOID RECEPTOR FAMILY OF PROTEINS
AND USES THEREOF

the specification of which

(check one)

X is attached hereto.

___ was filed on _____ as

Application Serial No. _____

and was amended on _____
(if applicable)

I hereby state that I have reviewed and understand the contents of the above-identified specification, including the claims, as amended by any amendment referred to above.

This application in part discloses and claims subject matter disclosed in my earlier filed application(s), as follows:

X Serial No. 60/108,322, filed November 13, 1998 ;
Serial No. 60/114,078, filed December 28, 1998, as to which I claim priority
benefit under Title 35, United States Code, §119(e).

X Serial No. 09/258,633, filed February 26, 1999 ;
Serial No. 09/421,134, filed October 19, 1999, as to which I claim priority
benefit under Title 35, United States Code, §120.

I acknowledge the duty to disclose to the United States Patent and Trademark Office all information known to me to be material to patentability as defined in Title 37, Code of Federal Regulations, §1.56, including all information known to me to be material to patentability as defined in Title 37, Code of Federal Regulations, §1.56 which became available between the filing date of the prior application(s) and the national or PCT international filing date of the continuation-in-part application.

As to the subject matter of this application which is common to said earlier application, I do not know and do not believe that the same was ever known or used in the United States of America before my or our invention thereof or patented or described in any printed publication in any country before my or our invention thereof, or more than one year prior to said earlier application, or in public use or on sale in the United States of America more than one year prior to said earlier application; that the common subject matter has not been patented or made the subject of an inventor's certificate issued before the date of said earlier application in any country foreign to the United States of America on an application filed by me or my legal representatives or assigns more than twelve months prior to said earlier application; and

As to applications for patents or inventor's certificate or PCT international application(s) designating at least one country other than the United States of America, on the common subject matter, filed in or designating any country foreign to the United States of America, prior to said earlier application by me or my legal representatives or assigns,

Check one:

☒ no such applications have been filed.

☐ such applications have been filed as follows

EARLIEST FOREIGN APPLICATION(S), IF ANY, FILED WITHIN 12 MONTHS
(6 MONTHS FOR DESIGN) PRIOR TO SAID EARLIER U.S. APPLICATION

Country	Application Number	Date of Filing (month,day,year)	Priority Claimed Under 35 USC 119
			_ Yes No _
			_ Yes No _
			_ Yes No _
			_ Yes No _
			_ Yes No _

ALL FOREIGN APPLICATION(S), IF ANY, FILED MORE THAN 12 MONTHS
(6 MONTHS FOR DESIGN) PRIOR TO SAID EARLIER U.S. APPLICATION

As to the subject matter of this application which is not common to said earlier application, I do not know and do not believe that the same was ever known or used in the United States of America before my or our invention thereof or patented or described in any printed publication in any country before my or our invention thereof, or more than one year prior to this application, or in public use or on sale in the United States of America more than one year prior to this application; that said non-common subject matter has not been patented or made the subject of an inventor's certificate issued before the date of this application in any country foreign to the United States of America on an application filed by me or my legal representatives or assigns more than twelve months prior to this application; and

As to applications for patents or inventor's certificate or PCT international application(s) designating at least one country other than the United States of America, on said non-common subject matter, filed in or designating any country foreign to the United States of America, prior to this application by me or my legal representatives or assigns,

Check one:

☒ no such applications have been filed.

☐ such applications have been filed as follows

EARLIEST FOREIGN APPLICATION(S), IF ANY, FILED WITHIN 12 MONTHS
(6 MONTHS FOR DESIGN) PRIOR TO THIS U.S. APPLICATION

Country	Application Number	Date of Filing (month,day,year)	Priority Claimed Under 35 USC 119
			_ Yes No _
			_ Yes No _
			_ Yes No _
			_ Yes No _
			_ Yes No _

ALL FOREIGN APPLICATION(S), IF ANY, FILED MORE THAN 12 MONTHS
(6 MONTHS FOR DESIGN) PRIOR TO THIS U.S. APPLICATION

CLAIM FOR BENEFIT OF U.S. PROVISIONAL APPLICATION(S)

I hereby claim the benefit under 35 U.S.C. §119(e) of any United States provisional application(s) listed below.

60/108,322
(Application Serial No.)

November 13, 1998
(Filing Date)

60/114,078
(Application Serial No.)

December 28, 1998
(Filing Date)

CLAIM FOR BENEFIT OF U.S. PATENT APPLICATION(S)

I hereby claim the benefit under 35 U.S.C. §120 of any United States patent application(s) listed below.

09/258,633
(Application Serial No.)

February 26, 1999
(Filing Date)

09/421,134
(Application Serial No.)

October 19, 1999
(Filing Date)

POWER OF ATTORNEY: As a named inventor, I hereby appoint the following attorneys and/or agents to prosecute this application and transact all business in the Patent and Trademark Office connected therewith.

W. Hugo Liepmann	Reg. No. 20,407	Catherine J. Kara	Reg. No. 41,106
James E. Cockfield	Reg. No. 19,162	Faustino A. Lichauco	Reg. No. 41,942
Thomas V. Smurzynski	Reg. No. 24,798	Jeanne M. DiGiorgio	Reg. No. 41,710
Ralph A. Loren	Reg. No. 29,325	Megan E. Williams	Reg. No. 43,270
Giulio A. DeConti, Jr.	Reg. No. 31,503	Nicholas P. Triano III	Reg. No. 36,397
Ann Lamport Hammitte	Reg. No. 34,858	Peter C. Lauro	Reg. No. 32,360
Elizabeth A. Hanley	Reg. No. 33,505	Reza Mollaaghababa	Reg. No. 43,810
Amy E. Mandragouras	Reg. No. 36,207	Timothy J. Douros	Reg. No. 41,716
Anthony A. Laurentano	Reg. No. 38,220	John L. Welch	Reg. No. 28,129
Jane E. Remillard	Reg. No. 38,872	DeAnn F. Smith	Reg. No. 36,683
Jeremiah Lynch	Reg. No. 17,425	William D. DeVaul	Reg. No. 42,483
Kevin J. Canning	Reg. No. 35,470	David J. Ridders	Reg. No. 43,882
David A. Lane, Jr.	Reg. No. 39,261	Chi Suk Kim	Reg. No. 42,728

all of: LAHIVE & COCKFIELD, LLP, 28 State Street, Boston, MA 02109

and to: Jean M. Silveri	Reg. No. 39,030	Cynthia Kanik	Reg. No. 37,320
Mark F. Boshar	Reg. No. 35,456	Theodore Allen	Reg. No. 41,578

of: Millennium Pharmaceuticals, Inc., 75 Sidney Street, Cambridge, MA 02139

Send Correspondence to Amy E. Mandragouras at Customer Number 000959 whose address is:

Lahive & Cockfield, LLP, 28 State Street, Boston, MA 02109

Direct Telephone Calls to: (name and telephone number)

Amy E. Mandragouras, (617) 227-7400

Wherefore I petition that letters patent be granted to me for the invention or discovery described and claimed in the attached specification and claims, and hereby subscribe my name to said specification and claims and to the foregoing declaration, power of attorney, and this petition.

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

Full name of sole or first inventor Rory A.J. Curtis	
Inventor's signature <i>Rory Curtis</i>	Date 11/11/99
Residence 31 Constitution Drive, Southborough, Massachusetts 01772	
Citizenship U.S.	
Post Office Address (if different)	

In re the application of: Rory A.J. Curtis

Serial No.: N/A

Filed: Herewith

For: NOVEL MEMBERS OF THE CAPSAICIN/VANILLOID
RECEPTOR FAMILY OF PROTEINS AND USES THEREOF

Attorney Docket No.: MNI-062CP2DV1

Group Art Unit:

Examiner:

Assistant Commissioner for Patents
Washington, D.C. 20231

CERTIFICATION UNDER 37 CFR 1.10

Date of Deposit: June 2, 2000

Mailing Label Number: EL 266 739 353 US

I hereby certify that this 37 CFR 1.53(b) request and the documents referred to as attached therein are being deposited with the United States Postal Service on the date indicated above in an envelope as "Express Mail Post Office to Addressee" service under 37 CFR 1.10 and addressed to the Assistant Commissioner for Patents, Box Patent Application, Washington, D.C. 20231.

Larry Taylor
Name of Person Mailing Paper

Larry Taylor
Signature of Person Mailing Paper

ASSOCIATE POWER OF ATTORNEY

Sir:

The undersigned attorney has the power of attorney in the subject application. She hereby grants an associate power to:

Maria C. Laccotripe
LAHIVE & COCKFIELD, LLP
28 State Street
Boston, MA 02109

As demonstrated by Appendix A (submitted herewith) Maria C. Laccotripe has passed the Patent Bar Examination and has been granted limited recognition under 37 C.F.R. § 10.9(b).

Please continue to forward all written and telephonic communications to Amy E. Mandragouras at the address and telephone number listed below.

Respectfully submitted,

Catherine J. Kara
Catherine J. Kara, Ph.D.
Registration No. 41,106
Attorney for Applicant

LAHIVE & COCKFIELD, LLP
28 State Street
Boston, MA 02109
Tel. (617) 227-7400
Dated: June 2, 2000


**BEFORE THE OFFICE OF ENROLLMENT AND DISCIPLINE
UNITED STATE PATENT AND TRADEMARK OFFICE**

LIMITED RECOGNITION UNDER 37 CFR § 10.9(b)

Maria C. Laccotripe is hereby given limited recognition under 37 CFR § 10.9(b) as an employee of Lahive and Cockfield to prepare and prosecute patent applications wherein the patent applicant is the client of Lahive and Cockfield, and the attorney or agent of record in the applications is a registered practitioner who is a member of Lahive and Cockfield. This limited recognition shall expire on the date appearing below, or when whichever of the following events first occurs prior to the date appearing below: (i) Maria C. Laccotripe ceases to lawfully reside in the United States, (ii) Maria C. Laccotripe's employment with Lahive and Cockfield ceases or is terminated, or (iii) Maria C. Laccotripe ceases to remain or reside in the United States on an H-1 visa.

This document constitutes proof of such recognition. The original of this document is on file in the Office of Enrollment and Discipline of the U.S. Patent and Trademark Office.

Expires: October 5, 2000



**Harry I. Moatz, Acting Director
Office of Enrollment and Discipline**

003090-7748960